

Doctoral School in Health Sciences  
Doctoral Programme in Drug Research

**EXPERIMENTAL NEUROTROPHIC THERAPIES OF PARKINSON'S DISEASE:  
EFFECTS ON NIGROSTRIATAL DOPAMINE SYSTEM**

**Juho-Matti Renko**

Division of Pharmacology and Pharmacotherapy  
Faculty of Pharmacy  
University of Helsinki  
Finland

DOCTORAL DISSERTATION

To be presented, with the permission  
of the Faculty of Pharmacy, University of Helsinki, for public examination  
in Viikki Biocenter 2, auditorium 1041, on 14 January 2021, at 13:00.

**Supervisors**

Professor Raimo Tuominen, MD, PhD  
Division of Pharmacology and Pharmacotherapy  
Faculty of Pharmacy  
University of Helsinki  
Finland

Docent Merja Voutilainen, PhD  
Institute of Biotechnology  
University of Helsinki  
Finland

**Reviewers**

Professor (hc) Seppo Kaakkola, MD, PhD  
Department of Neurology  
University of Helsinki  
Finland

Docent Sanna Janhunen, PhD  
University of Helsinki  
Finland

**Opponent**

Professor Edgar Kramer, PhD  
Peninsula Medical School  
Institute of Translational and Stratified Medicine  
Faculty of Health and Human Sciences  
University of Plymouth  
United Kingdom

© Juho-Matti Renko 2020

Dissertationes Scholae Doctoralis Ad Sanitatem Investigandam Universitatis Helsinkiensis

ISSN 2342-3161 (print)

ISSN 2342-317X (online)

ISBN 978-951-51-6936-5 (print)

ISBN 978-951-51-6937-2 (online)

Helsinki, Finland 2020

Press: Painosalama Oy, Turku



*“Kites rise highest against the wind, not with it.”*  
— Sir Winston Churchill

# TABLE OF CONTENTS

## ABSTRACT

## TIIVISTELMÄ

## LIST OF ORIGINAL PUBLICATIONS

## ABBREVIATIONS

<b>1</b>	<b>INTRODUCTION .....</b>	<b>1</b>
<b>2</b>	<b>REVIEW OF THE LITERATURE .....</b>	<b>3</b>
2.1	Neuroplasticity and neurotrophic therapies .....	3
2.1.1	Neuroplasticity.....	3
2.1.2	Neurotrophic factors.....	4
2.1.3	GDNF family ligands .....	6
2.1.4	CDNF/MANF family of neurotrophic factors.....	15
2.1.5	Retrograde signaling of neurotrophic factors .....	23
2.1.6	Neurotrophic factor mimetics, inducers and variants .....	24
2.2	Basal ganglia circuitry and dopaminergic system.....	27
2.2.1	Organization of the basal ganglia circuits .....	28
2.2.2	Functions of the basal ganglia circuits .....	30
2.2.3	Striatal interneurons .....	32
2.2.4	Striatal microcircuits regulating the direct and indirect pathways.....	33
2.2.5	Dopamine cell groups and projection pathways.....	33
2.2.6	Dopamine receptors and modulatory effects .....	36
2.2.7	Dopamine lifecycle .....	37
2.3	Parkinson's disease.....	43
2.3.1	Epidemiology and risk factors .....	43
2.3.2	Pathology of Parkinson's disease .....	45
2.3.3	Symptoms and diagnosis of Parkinson's disease .....	54
2.3.4	Biomarkers of Parkinson's disease .....	55
2.3.5	Current treatments for Parkinson's disease .....	58
2.4	Animal models of Parkinson's disease .....	62
2.4.1	Pharmacological models .....	62
2.4.2	Neurotoxin-based models .....	62
2.4.3	Genetic models .....	65
2.4.4	Alpha-synuclein pathology spreading model.....	66
2.4.5	Behavioral and histological assessment in animal models .....	66
2.5	Preclinical studies of neurotrophic factors for Parkinson's disease.....	67
2.5.1	Effects of neurotrophic factors on dopaminergic function of intact nigrostriatal system.....	67
2.5.2	Effects of neurotrophic factors in animal models of Parkinson's disease .....	69
2.6	Neurotrophic factors in clinical trials for Parkinson's disease .....	72
2.6.1	Clinical studies with GDNF.....	72
2.6.2	Clinical studies with NRTN.....	74
2.6.3	The first clinical study with CDNF .....	75
<b>3</b>	<b>AIMS OF THE STUDY .....</b>	<b>76</b>
<b>4</b>	<b>MATERIALS AND METHODS.....</b>	<b>77</b>
4.1	Animals.....	77
4.2	Drugs, toxins and neurotrophic therapies.....	77
4.2.1	Neurotrophic factors.....	77
4.2.2	Small molecule RET agonists .....	77

4.2.3	Neurotoxin .....	78
4.2.4	Drugs .....	78
4.3	Stereotaxic surgeries .....	78
4.3.1	Neurotrophic factor injections and guide cannula implantation .....	79
4.3.2	6-OHDA injections .....	79
4.3.3	Infusions of RET agonists and GDNF with osmotic minipumps .....	79
4.4	Microdialysis experiments .....	80
4.5	Enzymes activity measurements.....	81
4.6	Behavioral tests.....	82
4.6.1	Amphetamine-induced rotational behavior .....	82
4.6.2	Cylinder test.....	83
4.7	Perfusion and brain sectioning .....	83
4.8	Immunohistochemistry .....	83
4.9	Histological analyses.....	85
4.9.1	Fiber density in the striatum.....	85
4.9.2	Cell counts in the SNpc .....	85
4.10	Statistical analyses.....	86
<b>5</b>	<b>RESULTS .....</b>	<b>87</b>
5.1	Injection of neurotrophic factors into the intact rat brain .....	87
5.1.1	Effects of exogenous neurotrophic factors on dopamine release and turnover in the striatum (I).....	87
5.1.2	Effects of exogenous neurotrophic factors on dopamine neurochemistry (I).....	88
5.1.3	Effects of exogenous neurotrophic factors on GABA release in the GPe (unpublished data).....	89
5.1.4	Distribution of CDNF after nigral injection (II) .....	89
5.2	Activation of neuronal pro-survival signaling pathways by RET agonists .....	92
5.2.1	RET agonists phosphorylate RET and induce intracellular signaling cascades (III and IV) .....	93
5.2.2	RET agonists promote the survival of cultured midbrain dopamine neurons (III and IV).....	95
5.3	Delivery of RET agonists into the striatum of hemiparkinsonian rats .....	96
5.3.1	BT13 protects against motor dysfunction in 6-OHDA model of Parkinson's disease (III).....	96
5.3.2	BT44 shows neurorestorative potential in 6-OHDA model of Parkinson's disease (IV) .....	97
<b>6</b>	<b>DISCUSSION.....</b>	<b>101</b>
6.1	Effects of exogenously administered neurotrophic factors on dopamine release in the striatum .....	101
6.2	GABA release in the globus pallidus after exogenously administered neurotrophic factors .....	103
6.3	Effects of exogenously administered neurotrophic factors on dopamine synthesis and metabolism ....	104
6.4	Spreading properties of neurotrophic factors after administration into the basal ganglia .....	105
6.5	Potential of neurotrophic factor mimetics as a therapeutic strategy for Parkinson's disease .....	107
6.6	Methodological considerations .....	109
6.6.1	Considerations relating to 6-OHDA lesion models .....	109
6.6.2	Relevance of the behavioral tests in 6-OHDA lesion models .....	111
6.6.3	Validity of the disease models .....	113
6.7	General discussion.....	114
6.7.1	Importance of biomarker development .....	116
6.7.2	Issues with the clinical studies.....	117
6.7.3	Future directions .....	118
<b>7</b>	<b>CONCLUSIONS.....</b>	<b>120</b>
	<b>ACKNOWLEDGEMENTS .....</b>	<b>121</b>
	<b>REFERENCES .....</b>	<b>123</b>
	<b>APPENDIX: Original publications I-IV</b>	

# ABSTRACT

Parkinson's disease (PD) is a chronic neurodegenerative disorder characterized by progressive loss of nigrostriatal dopamine neurons and propagating Lewy body pathology. Dopamine depletion in the striatum gives rise to the cardinal motor symptoms of PD. Current PD medications are based on replenishing striatal dopamine and provide symptomatic relief to the motor deficits. However, troublesome adverse effects and diminished efficacy complicate their long-term use. There is a great unmet medical need for a therapy that could slow or halt the progression of the disease. Neurotrophic factors (NTFs) are secreted proteins that promote neuronal growth, differentiation and survival. They are able to prevent the progression of neurodegeneration and restore aberrant neuronal function in a variety of preclinical models of PD. Nonetheless, outcomes from clinical trials have been disappointing. The purpose of this work was to characterize the effects of cerebral dopamine neurotrophic factor (CDNF), mesencephalic astrocyte-derived neurotrophic factor (MANF) and novel small molecule receptor tyrosine kinase RET agonists (BT13 and BT44) on nigrostriatal dopamine system and support their preclinical development as potential neurotrophic therapies of PD.

To further clarify the functional effects of glial cell line-derived neurotrophic factor (GDNF), CDNF and MANF in the normal rat brain, microdialysis measurements were performed after a bolus injection of NTFs into the striatum. We saw augmented stimulus-evoked dopamine release and elevated dopamine turnover in the striatum of MANF-injected rats. GDNF injection increased *in vivo* tyrosine hydroxylase (TH) and catechol-*O*-methyltransferase activity and decreased monoamine oxidase A activity. These data are relevant when considering exogenously administered NTFs as a potential therapeutic approach for PD, since they have to be compatible with the existing dopaminergic medications of the patients.

We also investigated the distribution properties of <sup>125</sup>I-labeled and unlabeled CDNF after a nigral injection to intact rats. CDNF readily diffused into the brain areas surrounding the injection site and colocalized with TH-immunoreactive neurons in the substantia nigra. We did not detect active transportation of CDNF to distal brain areas. This characterization provides valuable insights into the selection of optimal delivery site and protocol for CDNF therapy.

Our *in vitro* assays showed that RET agonists BT13 and BT44 were able to induce RET phosphorylation and activate downstream pro-survival signaling cascades Akt and ERK. They also supported the survival of cultured midbrain dopamine neurons from wild-type, but not from RET knockout, mice. The functional effects of BT13 and BT44 were evaluated in a unilateral 6-hydroxydopamine rat model of PD, where both compounds alleviated amphetamine-induced turning behavior. BT44 also showed potential to restore striatal TH-immunoreactive fibers. As blood-brain barrier penetrating compounds, BT13 and BT44 serve as promising leads that can be further developed into a disease-modifying therapy for PD.

# TIIVISTELMÄ

Parkinsonin tauti on etenevä hermorappeumasairaus, jolle on ominaista aivojen mustatumakkeen dopamiinihermosolujen tuhoutuminen ja Lewyn kappaleiden esiintyminen aivoissa. Dopamiinihermosolujen tuhoutuminen johtaa aivojuovion dopamiinivajeeseen, joka saa aikaan Parkinsonin taudille ominaiset liikehäiriöt. Taudin nykyinen lääkitys perustuu dopamiinivajeen korjaamiseen ja on luonteeltaan oireita lievittävää. Pitkäaikaiskäytössä lääkkeet menettävät tehoaan ja johtavat hankaliin haittavaikutuksiin. Taudin kulkua hidastavalle hoidolle onkin suuri tarve. Hermokasvutekijät ovat hermosolujen kasvua, erilaistumista ja selviytymistä edistäviä proteiineja. Niiden on osoitettu estävän dopamiinihermosolujen tuhoutumista ja korjaavan niiden häiriintynyttä toimintaa useissa kokeellisissa Parkinson-malleissa. Kliinisissä kokeissa testattujen hermokasvutekijähoitojen teho on kuitenkin jäänyt puutteelliseksi. Tämän väitöstyön tarkoituksena oli selvittää CDNF:n (dopamiinisolujen hermokasvutekijä) ja MANF:n (keskiaivojen astrosyyttiperäinen hermokasvutekijä) sekä uusien, RET-tyrosiinikinaasia aktivoivien pienmolekyylien BT13:n ja BT44:n vaikutuksia aivojen dopamiinijärjestelmään ja samalla tukea niiden prekliinistä kehitystä mahdolliseksi Parkinsonin taudin kulkuun vaikuttavaksi hoidoksi.

Ensimmäisessä osatyössä selvitimme aivojuovioon annostellun GDNF:n (gliasolulinjaperäinen hermokasvutekijä), CDNF:n ja MANF:n aiheuttamia toiminnallisia muutoksia terveen rotan aivoissa käyttämällä hyväksi mikrodialyysimenetelmää. Havaitimme MANF:n lisäävän stimuloitua dopamiinin vapautumista ja dopamiinin aineenvaihduntaa aivojuoviossa. GDNF injektio puolestaan lisäsi dopamiinia syntetisoivan tyrosiinihydroksylaasientsyymin aktiivisuutta. Lisäksi se vaikutti dopamiinia metaboloivien entsyymien toimintaan lisäämällä katekoli-O-metyylitransferaasin ja vähentämällä monoamiinioksidaasin aktiivisuutta. Nämä löydökset auttavat sovittamaan mahdollisia hermokasvutekijähoitoja yhteen Parkinson-potilaiden käyttämien dopaminergisten lääkkeiden kanssa.

Toisessa osatyössä tutkimme CDNF:n leviämisominaisuuksia terveen rotan aivoissa mustatumakkeeseen kohdistuneen annostelun jälkeen. CDNF levisi laajalle injektiokohtaa ympäröivään aivokudokseen ja oli havaittavissa mustatumakkeen dopamiinihermosoluissa. Merkkejä CDNF:n aktiivisesta kuljetuksesta kaukaisemmille aivoalueille ei havaittu. Tämä tutkimus tarjoaa arvokasta lisätietoa CDNF:n optimaalisen annostelutavan määrittämiseksi.

Kolmannessa ja neljännessä osatyössä osoitimme BT13:n ja BT44:n aktivoivan soluja suojaavat Akt- ja ERK-signaalointireitit sekä edistävän viljeltyjen dopamiinihermosolujen selviytymistä. Kun BT13:n ja BT44:n vaikutuksia tutkittiin Parkinsonin taudin eläinmallissa rotilla, molempien yhdisteiden havaittiin lievittävän kokeessa mitattua liikehäiriötä. BT44 osoitti myös viitteitä dopaminergisia hermosäikeitä korjaavasta vaikutuksesta. Veri-aivoesteen läpäisevät BT13 ja BT44 ovatkin lupaavia johtolankamolekyylijä, joita edelleen optimoimalla voisi olla mahdollista kehittää uusi Parkinsonin taudin etenemistä hidastava hoito.

# LIST OF ORIGINAL PUBLICATIONS

This thesis is based on the following publications:

- I Renko J-M, Bäck S, Voutilainen MH, Piepponen TP, Reenilä I, Saarma M, Tuominen RK (2018) Mesencephalic Astrocyte-Derived Neurotrophic Factor (MANF) Elevates Stimulus-Evoked Release of Dopamine in Freely-Moving Rats. *Mol. Neurobiol.* 55(8):6755-6768
- II Albert K, Renko J-M, Mätlik K, Airavaara M, Voutilainen MH (2019) Cerebral Dopamine Neurotrophic Factor Diffuses Around the Brainstem and Does Not Undergo Anterograde Transport After Injection to the Substantia Nigra. *Front. Neurosci.* 13:590
- III Mahato AK\*, Renko J-M\*, Kopra J, Visnapuu T, Korhonen I, Pulkkinen N, Bespalov M, Ronken E, Piepponen TP, Voutilainen MH, Tuominen RK, Karelson M, Sidorova YA, Saarma M (2019) GDNF receptor agonist supports dopamine neurons in vitro and protects their function in animal model of Parkinson's disease. bioRxiv 540021, *Preprint manuscript*
- IV Renko J-M\*, Mahato AK\*, Visnapuu T, Karelson M, Voutilainen MH, Saarma M, Tuominen RK, Sidorova YA (2020) Neuroprotective potential of a small molecule RET agonist in cultured dopamine neurons and hemiparkinsonian rats. *Manuscript under review*

\* Equal contribution

The publications are referred to in the text by their roman numerals. Reprints were made with the permission of the copyright holders.

# ABBREVIATIONS

3-MT	3-Methoxytyramine
6-OHDA	6-Hydroxydopamine
AADC	Aromatic amino acid decarboxylase
AAV	Adeno-associated virus
AD	Alzheimer's disease
ANOVA	Analysis of variance
ARTN	Artemin
ATP	Adenosine triphosphate
$\alpha$ -syn	$\alpha$ -Synuclein
BBB	Blood-brain barrier
BDNF	Brain-derived neurotrophic factor
CDNF	Cerebral dopamine neurotrophic factor
CNS	Central nervous system
COMT	Catechol- <i>O</i> -methyltransferase
CSF	Cerebrospinal fluid
DAT	Dopamine transporter
DOPAC	3,4-Dihydroxyphenylacetic acid
ENS	Enteric nervous system
ER	Endoplasmic reticulum
$^{18}\text{F}$ -DOPA	6- $^{18}\text{F}$ -fluoro-L-dopa
FGF	Fibroblast growth factor
GABA	$\gamma$ -Aminobutyric acid
GDNF	Glial cell line-derived neurotrophic factor
GFL	GDNF family ligand
GFP	Green fluorescent protein
GFR $\alpha$	GDNF family receptor $\alpha$
GPe	Globus pallidus external segment
GPi	Globus pallidus internal segment
GPI	Phosphatidylinositol
GRP78	Glucose-regulated protein 78 kDa
HPLC	High-performance liquid chromatography
HVA	Homovanillic acid
i.p.	Intraperitoneally
ir	Immunoreactive
L-DOPA	L-3,4-Dihydroxyphenylalanine
LPS	Lipopolysaccharide
LTD	Long-term depression
LTP	Long-term potentiation
MANF	Mesencephalic astrocyte-derived neurotrophic factor

MAO	Monoamine oxidase
MAPK	Mitogen activated protein kinase
MCI	Mild cognitive impairment
MPP+	1-Methyl-4-phenylpyridinium
MPTP	1-Methyl-4-phenyl-1,2,3,6- tetrahydropyridine
MRI	Magnetic resonance imaging
MSN	Medium spiny neuron
NAcc	Nucleus accumbens
nAChR	Nicotinic acetylcholine receptor
NCAM	Neural cell adhesion molecule
NGF	Nerve growth factor
NHP	Non-human primate
NRTN	Neurturin
NTF	Neurotrophic factor
PBS	Phosphate-buffered saline
PD	Parkinson's disease
PET	Positron emission tomography
PG	Propylene glycol
PNS	Peripheral nervous system
PSPN	Persephin
PV	Parvalbumin
REGWF	Ryan-Einot-Gabriel-Welsch F
RET	Rearranged during transfection
RMS	Rostral migratory stream
ROS	Reactive oxygen species
s.c.	Subcutaneous
SEM	Standard error of the mean
SGZ	Subgranular zone
SN	Substantia nigra
SNpc	Substantia nigra pars compacta
SNr	Substantia nigra pars reticulata
SPECT	Single-photon emission computed tomography
STN	Subthalamic nucleus
SVZ	Subventricular zone
TGF- $\beta$	Transforming growth factor $\beta$
TH	Tyrosine hydroxylase
UPDRS	Unified Parkinson's disease rating scale
UPR	Unfolded protein response
VMAT2	Vesicular monoamine transporter 2
VTA	Ventral tegmental area



# 1 INTRODUCTION

Parkinson's disease (PD) is a debilitating neurological disorder with increasing prevalence in aging populations. The cardinal motor symptoms of PD arise from the progressive degeneration of nigrostriatal dopamine neurons and the resultant dopamine depletion in the dorsal striatum (Kalia and Lang 2015; McGregor and Nelson 2019). This causes imbalance in the excitability of the direct and indirect striatal projection pathways within the basal ganglia leading to inadequate activation of the motor cortex and impaired motor functions. The underlying pathological processes of PD are incompletely understood, but they seem to be associated with neuroinflammation and the presence of intracellular protein inclusions called Lewy bodies.

There is no cure for PD. The current treatments provide symptomatic relief mainly to the motor symptoms by increasing dopaminergic activity within the dorsal striatum (Armstrong and Okun 2020). With the progression of the disease, however, the therapies gradually lose their effects and start to be accompanied by troublesome adverse effects such as motor fluctuations, dyskinesias and psychiatric symptoms. The dopaminergic treatments have minor effects on disabling non-motor symptoms of PD and, importantly, they are unable to halt the neurodegenerative processes underlying the disease. Thus, a disease-modifying therapy remains an urgent unmet medical need for PD.

Our brain has an innate capacity to reorganize its structure, neuronal connections and functions in response to intrinsic and extrinsic stimuli (Cramer et al. 2011). These dynamic adaptations, commonly called neuroplasticity, have a pivotal role in the proper function of the nervous system. Neurotrophic factors (NTFs) are small secreted proteins that regulate almost all aspects of neuroplasticity including neurogenesis, neuronal development and maintenance and, importantly, the survival and recovery of neurons (Paratcha and Ledda 2008). Therefore, NTFs serve as promising candidates for developing disease-modifying therapies for neurodegenerative disorders such as PD.

There are four major families of NTFs, two of which, GDNF family ligands and CDNF/MANF family of NTFs, are of special interest in this thesis. Glial cell line-derived neurotrophic factor (GDNF) is the founding member of GDNF family ligands and was identified in 1993 based on its survival-promoting effects on cultured midbrain dopamine neurons (Lin et al. 1993). Thereafter, intensive research efforts have uncovered the signaling mechanisms of GDNF as well as its robust survival promoting and regenerative effects on midbrain dopamine neurons both *in vitro* and in animal models of PD.

Cerebral dopamine neurotrophic factor (CDNF) and mesencephalic astrocyte-derived neurotrophic factor (MANF) form structurally and functionally distinct family of NTFs (Lindholm et al. 2007; Petrova et al. 2003). They have also shown strong neurotrophic properties on midbrain dopamine neurons promoting their survival and repair *in vitro* and in animal models of PD. Their precise mechanism of action, however, has remained undetermined thus far. Being endoplasmic reticulum (ER)-resident proteins, CDNF and MANF have been suggested to regulate ER stress and

subsequent unfolded protein response (UPR). As inflammation and ER stress responses are closely linked with each other, their neuroprotective effects may be due to regulation of neuroinflammation in the brain.

Following promising outcomes in preclinical experiments, intracranial GDNF application proceeded into clinical trials with PD patients. These trials suggest a favorable safety profile for the NTF therapies together with some promising efficacy outcomes, yet none of the randomized, placebo-controlled trials met their primary endpoints (Nutt et al. 2003; Lang et al. 2006; Whone et al. 2019). The varying success in the clinical trials has raised questions about the optimal dosing paradigm and protein therapy related challenges. NTFs are, for example, unable to cross the blood-brain barrier (BBB), and thus require intracranial delivery increasing treatment-related risks and costs. A small molecule that mimics the effects of NTFs on degenerating dopamine neurons and is suitable for systemic administration would be an attractive drug candidate for PD.

In the present study, we pursued new insights into the biological effects of intrastrially injected CDNF and MANF on dopamine synthesis, release and metabolism in the normal rat brain in order to gain better understanding of their therapeutic applicability for degenerative brain diseases. We also further elucidated the spreading properties of CDNF in the rat brain after intranigral injection which is highly relevant information in terms of finding the optimal sites of administration. Lastly, we investigated two novel small molecule RET agonists, their signaling properties and neuroprotective effects on nigrostriatal dopamine neurons in a neurotoxin-induced rat model of PD. These agonists serve as promising lead compounds that may open avenues in the search of a BBB-penetrating neurotrophic therapy for PD.

The literature review will first give an overview of neuroplasticity and NTFs, their structure, expression and signaling mechanisms, focusing on GDNF family ligands, CDNF/MANF family of NTFs and NTF mimetics. Then, the organization and functions of the basal ganglia circuitry will be reviewed together with dopamine projection pathways, modulatory effects and lifecycle in the brain. Finally, PD and preclinical and clinical studies of NTFs in PD will be described.

## 2 REVIEW OF THE LITERATURE

### 2.1 Neuroplasticity and neurotrophic therapies

In this thesis, neurotrophic therapies refer to a concept of utilizing endogenous NTFs or similarly acting small molecule compounds (i.e. NTF mimetics) to induce neuroplasticity changes in the central nervous system (CNS). Neurotrophic therapies are considered as potential disease-modifying treatments for various neurological disorders.

#### 2.1.1 Neuroplasticity

The brain has traditionally been considered as a static organ, without turnover of neurons or capacity for repair after neuronal insults. However, it has a fundamental ability to reorganize its structure, connections and function in response to intrinsic and extrinsic stimuli (Cramer et al. 2011). These long-lasting and bidirectional alterations are of crucial importance for neuronal development and brain functions, e.g. learning. Developmental and environmental changes, diseases and therapeutic interventions induce adaptive neuroplastic changes which take place at many levels. During normal development, newborn neurons that are not functionally integrated into neural circuits are eliminated through selective apoptosis (Castrén and Hen 2013). It has been estimated, that up to 60% of the neurons originally generated die in most neuronal populations (Oppenheim 1991). This naturally occurring cell death is regulated by NTFs that promote the survival and maturation of neurons.

At cellular level, the morphology of mature neurons can be modified through arborization and pruning of axonal and dendritic branches and spines (Castrén and Hen 2013). The synapse-containing branches are more likely stabilized whereas spines lacking a synapse are readily pruned. The number of synapses is regulated dynamically through synaptogenesis and elimination of inactive synapses. At functional level, long-term potentiation (LTP) increases the synaptic strength, whereas long-term depression (LTD) suppresses it in inactive synapses. Active information transfer in LTP enhances the synaptic function and protects the synapse from pruning.

Growth factors, especially NTFs, have been shown to play a key role in regulating the dynamic adaptations in the neuronal connections and functions which are crucial for the proper function of the nervous system (Paratcha and Ledda 2008; Bothwell 2014; Lu et al. 2014; Levy et al. 2018). NTFs control almost all processes relating to the neuroplasticity including neural stem cell proliferation, migration and differentiation of neuroblasts, growth and survival of neurons, neurite outgrowth, formation of synapses and LTP. In addition, numerous transcriptional and epigenetic mechanisms regulate the expression of effector genes involved in neuroplasticity (Castrén and Hen 2013).

Brain resident innate immune cells, microglia, are also prominent players in the regulation of brain homeostasis and neuronal plasticity (Block and Hong 2005; Sierra et al. 2010; Kettenmann et al. 2013; Wake et al. 2013). Reactive microglia are primarily responsible for immune reactions in the

CNS in response to an infection or neuronal injury. Resting microglia, instead, are responsible for the surveillance and elimination of presynaptic terminals and dendritic spines in a process termed synaptic pruning. Microglia also release various substances such as adenosine triphosphate (ATP), NTFs and cytokines which regulate neuroinflammation, programmed cell death, neurogenesis, neuronal repair and synaptic connectivity. In addition, microglia phagocytose apoptotic cells and newborn neural progenitor cells. These functions contribute to the maturation, plasticity and homeostasis of neuronal circuits during development and adulthood.

New neurons are continuously supplied to the hippocampus and olfactory bulb in most mammals throughout life conferring plasticity to these neuronal circuits (Falk and Frisén 2005; Steiner et al. 2019). The newborn hippocampal neurons derive from local stem cells residing in the subgranular zone (SGZ) of the dentate gyrus, whereas neurons added to the olfactory bulb are derived from a neurogenic niche in the subventricular zone (SVZ) of the lateral ventricle wall from where they migrate along the rostral migratory stream (RMS). Innovative research strategies have demonstrated that substantial neurogenesis occurs also in the germinal areas of the adult human brain. The incorporation of 5-bromo-3'-deoxyuridine (BrdU), a synthetic nucleotide injected to cancer patients for diagnostic purposes, into the DNA of the dividing cells (Eriksson et al. 1998) or retrospective analysis of radioactive  $^{14}\text{C}$  incorporated into the genomic DNA as a consequence of increased atmospheric levels of  $^{14}\text{C}$  produced by nuclear weapon tests during the Cold War (Spalding et al. 2005) has enabled these research achievements. Measuring the level of nuclear-bomb-test-derived  $^{14}\text{C}$  in the neuronal DNA Spalding and coworkers revealed extensive neurogenesis of hippocampal cells also in adult humans (Spalding et al. 2013). Similarly, retrospective  $^{14}\text{C}$ -dating has revealed that a substantial number of newborn interneurons continuously integrates into the striatum (Ernst et al. 2014). In humans, neuroblasts arising from the SVZ provide a source for striatal neurons instead of migrating to the olfactory bulb via RMS.

### **2.1.2 Neurotrophic factors**

NTFs are endogenous proteins secreted by a variety of tissues in the body (reviewed in Huang and Reichardt 2001; Airaksinen and Saarma 2002). Importantly, NTFs support the survival of neurons and help them to recover from injuries making them a promising therapeutic strategy not only for the management of neurodegenerative disorders, such as PD, Alzheimer's disease (AD) and amyotrophic lateral sclerosis (ALS), but also for the treatment of neuronal traumas, e.g. spinal cord injury. There appears to be a shortage of NTFs in neurodegenerative diseases; for example in PD patients, decreased expression of GDNF, brain-derived neurotrophic factor (BDNF) and ciliary neurotrophic factor (CNTF) have been reported in the surviving neurons of the substantia nigra (SN) (Mogi et al. 1999; Parain et al. 1999; Howells et al. 2000; Chauhan et al. 2001). Apart from their important effects on the development and maintenance of neurons, NTFs also exert several essential functions outside the nervous system in non-neural tissues.

#### **2.1.2.1 Neurotrophic hypothesis**

Mature neurons have highly polarized morphology; they consist of a cell body containing the nucleus and other organelles, dendrites receiving synaptic input from neighboring cells, and one

long axon, which ends in presynaptic terminals. According to the classical neurotrophic hypothesis the target tissue of neuronal innervation secretes limiting quantities of NTFs which are transported retrogradely along the axon to the cell body where they suppress apoptosis of the innervating neuron (Hamburger 1939, 1934; Hamburger and Levi-Montalcini 1949). In this way, NTF secretion by the target organ ensures the balance between the size of the target and the number of innervating neurons. During development, most neuronal populations are initially overproduced. Neuronal target fields, however, produce NTFs in amounts that are not sufficient for all neurons. The lack of target-derived NTFs drives the neurons without proper connections with the target organ into programmed cell death, thus, regulating the innervation density of the organ. The main caveat of the abovementioned hypothesis is that it has been demonstrated to hold true almost exclusively in the peripheral nervous system (PNS). In the CNS, the relationship between target-derived NTFs and neuronal survival appears to be more complex because neighboring cells also can provide trophic support to neurons through paracrine secretion and a neuron might be able to secrete NTFs itself through an autocrine loop (Cerchia 2006). In addition, the classical neurotrophic hypothesis has been broadened since some neuronal populations seem to depend on several different NTFs regulating concurrently or sequentially the target organ innervation (Davies 1996).

#### 2.1.2.2 Characteristics and families of NTFs

Structurally mature NTFs proteins consist of approximately 100-160 amino acids and contain typically several conserved disulphide bridges between cysteine residues enabling closely related conformations within different NTF families (Ibáñez 1998; Airaksinen et al. 1999; Airaksinen and Saarma 2002; Lindholm and Saarma 2010; Bothwell 2014). Like most other secreted polypeptides, NTFs are synthesized and packaged into secretory vesicles in the rough ER. NTFs can be secreted from various neuronal and non-neuronal cells in the CNS and PNS. NTFs are commonly produced in the form of a precursor protein. The signal sequence is cleaved either intracellularly in secretory vesicles or by extracellular proteases producing mature NTFs. Mature NTFs usually form non-covalently associated dimers that bind to transmembrane receptor tyrosine kinases which in turn initiate intracellular signaling cascades resulting in trophic effects (Ibáñez 1998).

NTFs known today can be divided into four major families (Bothwell 2014; Ibáñez and Andressoo 2017; Lindahl et al. 2017):

1. neurotrophins including nerve growth factor (NGF), BDNF, neurotrophin-3 (NT-3) and neurotrophin-4 (NT-4)
2. GDNF family ligands (GFLs) including GDNF, neurturin (NRTN), artemin (ARTN), persephin (PSPN), and a distant member growth and differentiation factor-15 (GDF-15, also known as macrophage-inhibiting cytokine-1, MIC-1)
3. neurotrophic cytokines (=neurokines) including CNTF, cardiotrophin-1, leukemia inhibitory factor, neuropoietin, oncostatin M, cardiotrophin-like cytokine, interleukin 6 (IL-6), IL-11 and IL-27
4. CDNF/MANF family of NTFs.

Two of these families, GFLs and unconventional CDNF/MANF family of NTFs are in the focus of this thesis and will be discussed in more detail in the following chapters. Also, other trophic and growth factors, e.g. members of transforming growth factor  $\beta$  (TGF- $\beta$ ) superfamily, vascular endothelial growth factors, insulin-like growth factors and fibroblast growth factors (FGFs) are shown to have neurotrophic activities but those fall out of the scope of this thesis (Grothe and Timmer 2007; Zacchigna et al. 2008).


### **2.1.3 GDNF family ligands**

GDNF was identified and isolated in 1993 based on its survival-promoting effects on ventral midbrain dopamine neurons (Lin et al. 1993). NRTN, ARTN and PSPN, the other three family members structurally similar to GDNF, were characterized within five years after the discovery of GDNF (Kotzbauer et al. 1996; Baloh et al. 1998; Milbrandt et al. 1998). GDF-15 is a distant member of the GFLs which functions as a peripheral signal downregulating food intake, energy expenditure and body weight in response to tissue damage and stress (Bootcov et al. 1997; Böttner et al. 1999; Hsiao et al. 2000). GFLs are distant members of TGF- $\beta$  superfamily. Together with their receptors they form one of the major neurotrophic networks in the nervous system regulating the development, maintenance and function of a variety of neurons and glial cells (Ibáñez and Andressoo 2017). GFLs, their receptors and binding preferences are summarized in Figure 2.1.

#### **2.1.3.1 Synthesis, structure and secretion of GFLs**

GFLs are synthesized in the form of a precursor protein preproGFL (Lin et al. 1993; Lonka-Nevalaita et al. 2010). The pre-sequence guides GFLs to the ER for secretion. During secretion GFLs form disulfide-bonded homodimers and can be modified by N-linked glycosylation (Lin et al. 1993; Lonka-Nevalaita et al. 2010; Piccinini et al. 2013). Proteolytic cleavage of proGFLs into mature GFL proteins takes place extracellularly by furin, PACE4, PC5A, PC5B and PC7. After secretion GFLs bind to heparan-sulphate side chains of extracellular matrix proteoglycans which limits their diffusion in brain parenchyma and increases their local concentration.

GDNF, the founding member GFLs, is a ~20 kD, N-glycosylated protein consisting of 211 amino acids (Lin et al. 1993). The mature 134 amino acids long GDNF is formed after the cleavage of the pre- (19 amino acids) and pro- (58 amino acids) signaling sequences. The primary structure of GDNF contains seven conserved cysteine residues with the same relative spacing as in the other TGF- $\beta$  superfamily members. The tertiary structure of GDNF is stabilized with three disulphide bridges that are formed between the cysteine residues. The one remaining cysteine connects two GDNF molecules with each other via a disulphide bridge forming GDNF homodimer.



**Figure 2.1. GDNF family ligands (GFLs) and their receptors.** Glial cell line-derived neurotrophic factor (GDNF), neurturin (NRTN), artemin (ARTN), persephin (PSPN) and a distant member growth and differentiation factor-15 (GDF-15) bind preferentially to their cognate co-receptors GDNF family receptor  $\alpha$  (GFR $\alpha$ ) 1-4 and GDNF family receptor alpha-like (GFRAL), respectively, although cross-binding occurs too. Thereafter, ligand-receptor complex binds to the common transmembrane receptor tyrosine kinase RET or neural cell adhesion molecule (NCAM), which initiate intracellular signaling cascades. GFLs can also directly bind to and signal through syndecan-3. Glycosyl phosphatidylinositol (GPI) -anchor attaches GFR $\alpha$  receptors to the plasma membrane (PM) and localizes them into lipid rafts. Primary phosphorylated tyrosine residues of RET, which serve as docking sites for the intracellular adaptor proteins, and the  $\text{Ca}^{2+}$ -binding site in the middle of the four cadherin-like domains of the extracellular RET are depicted in the figure. High affinity binding is indicated with solid lines and low affinity binding with dashed lines. Figure drawn by the author, inspired by Kramer and Liss (2015).

### 2.1.3.2 RET-mediated signaling

Mature GFL homodimers mediate their biological effects via a multicomponent receptor complex consisting of GDNF family receptor  $\alpha$  (GFR $\alpha$ ) and receptor tyrosine kinase RET (rearranged during transfection) (Airaksinen and Saarma 2002). A GFL first binds to its cognate GFR $\alpha$  co-receptor which dimerizes. Subsequently, GFL-GFR $\alpha$  complex recruits RET as a signal transducing receptor and triggers its homodimerization and autophosphorylation of its intracellular tyrosine kinase domain. Apart from RET, two alternative signaling receptors have been identified: neural cell adhesion molecule (NCAM) (Chao et al. 2003; Paratcha et al. 2003) and syndecan-3 (Bespalov et al. 2011). A summary of the known GFL signaling mechanisms is illustrated in Figure 2.2.

RET was identified as the primary signaling receptor for GFLs in 1996 (Durbec et al. 1996a; Trupp et al. 1996; Vega et al. 1996; Worby et al. 1996). RET is a canonical single spanning transmembrane receptor tyrosine kinase. The extracellular region consists of four cadherin-like domains (CLD 1-4) with one  $\text{Ca}^{2+}$ -ion between CLD2 and CLD3 that is required for ligand binding to RET (Anders et al. 2001; Knowles et al. 2006). The cytoplasmic tyrosine kinase domain contains several tyrosine phosphorylation sites regulating the catalytic activities of RET (Myers et al. 1995; Tahira et al. 1990). In most cases, the phosphorylated residues are Tyr<sup>905</sup>, Tyr<sup>1015</sup>, Tyr<sup>1062</sup> and Tyr<sup>1096</sup> which serve as docking sites for signal-transducing adaptor proteins (Coulpier et al. 2002; Arighi et al. 2005). Alternative splicing of RET mRNA can produce three isoforms with varying length of the C-

terminal tail (RET9, RET43 and RET51). RET9 and RET51 are the major isoforms and highly conserved in different species of vertebrates (Carter et al. 2001).

Phosphorylation of the intracellular tyrosine residues of RET activates downstream signaling cascades including Ras/mitogen activated protein kinase (MAPK), phosphatidylinositol 3-kinase (PI3K)/Akt, phospholipase C- $\gamma$  (PLC $\gamma$ ) and c-Jun N-terminal kinase (JNK) pathways (Takahashi 2001). Ras/MAPK pathway leads to the activation of transcription factor cAMP response element-binding protein (CREB) and regulates cellular proliferation, growth, differentiation, survival and neuritogenesis (Hayashi et al. 2000). PI3K/Akt pathway activates transcription factor NF $\kappa$ B and is primarily responsible for cellular survival, growth, migration and proliferation.

RET is unable to bind GFLs in absence of the ligand-binding co-receptor GFR $\alpha$  (Jing et al. 1996; Treanor et al. 1996; Paratcha and Ledda 2008). The GFR $\alpha$ :s are a family of four glycosyl phosphatidylinositol (GPI) -anchored extracellular proteins, GFR $\alpha$ 1-4, that serve as preferential receptors for GDNF, NRTN, ARTN and PSPN, respectively (Jing et al. 1996; Treanor et al. 1996; Baloh et al. 1997; Buj-Bello et al. 1997; Jing et al. 1997; Klein et al. 1997; Sanicola et al. 1997; Suvanto et al. 1997; Baloh et al. 1998; Naveilhan et al. 1998; Thompson et al. 1998; Trupp et al. 1998; Widenfalk et al. 1998; Worby et al. 1998; Masure et al. 2000). In addition, GDNF family receptor alpha-like (GFRAL) was recently discovered as a co-receptor for GDF-15 (Emmerson et al. 2017; Hsu et al. 2017; Mullican et al. 2017; Yang et al. 2017). It mediates GDF-15–GFRAL signaling via RET in the same way as the other GFL–GFR $\alpha$  complexes. Apart from the high-affinity binding to the cognate co-receptors, some low-affinity cross-reactivities between different GFLs and GFR $\alpha$ :s have been demonstrated *in vitro* as summarized in Figure 2.1 and Table 2.1.

The GPI-anchor links GFR $\alpha$ :s to the plasma membrane and localizes them into lipid rafts, special subdomains of the plasma membrane (Tansey et al. 2000; Paratcha et al. 2001; Paratcha and Ibáñez 2002; Tsui et al. 2015). These cholesterol and sphingolipid-rich microdomains accumulate signaling proteins and thereby increase their interactions with each other. During receptor activation in *cis*, GFL homodimer first binds with high affinity to one of the GPI-anchored GFR $\alpha$  receptors which dimerizes (Airaksinen and Saarma 2002; Jing et al. 1996; Treanor et al. 1996) (Figure 2.2.A). Subsequently, GFL-GFR $\alpha$  complex recruits RET into the lipid raft. The relocation of RET into the lipid rafts potentiates downstream signal transduction through the receptor complex (Tansey et al. 2000). Through phosphorylated Tyr<sup>1062</sup> residue, for example, RET associates with the adaptor protein FRS2 inside the lipid rafts and with SHC outside the rafts (Paratcha et al. 2001). Inside the lipid rafts, FRS2 first recruits Grb2 and Sos proteins which then leads to the activation of Ras/MAPK pathway (Melillo et al. 2001). Outside the lipid rafts, SHC can recruit either Grb2 and Gab proteins leading to the activation of PI3K/Akt pathway, or Grb2 and Sos proteins activating Ras/MAPK pathway (Besset et al. 2000; Hayashi et al. 2000).

The GPI-anchor of GFR $\alpha$  receptor can also be cleaved by membrane-associated phospholipases or proteases which releases GFR $\alpha$  into the extracellular space and leads to the activation of RET in *trans* (Yu et al. 1998; Paratcha et al. 2001; Ledda et al. 2002). Soluble GFR $\alpha$  binds GFL in the extracellular space with high affinity after which the complex binds to and activates RET outside the lipid rafts where RET associates with SHC adaptor protein. Subsequently, RET is recruited into




the lipid raft where it activates FRS2. The fact that GFR $\alpha$  receptors are much more widely expressed than RET, together with their ability to activate RET in *trans*, suggest non-cell-autonomous functions for soluble GFR $\alpha$  during neuronal development and regeneration. Secretion of soluble GFR $\alpha$  by target tissues of RET-expressing neurons may act as a long-range cue that guides the growing axons.

It should be noted, that RET signaling is under negative control which regulates the magnitude and duration of RET activation. A prevalent mechanism to control the downregulation of RET is through ligand-induced ubiquitination which leads to proteasomal degradation of the active receptor (Carniti et al. 2003; Scott et al. 2005). In response to ligand-mediated activation, RET can also be internalized from the plasma membrane into early endosomes via clathrin-mediated endocytosis (Richardson et al. 2006; Richardson et al. 2012; Crupi et al. 2015). The endocytosed GFL-GFR $\alpha$ -RET complex contributes to the retrograde GFL signaling, but finally receptor internalization terminates the extracellular signal transduction.

### 2.1.3.3 NCAM-mediated signaling

GFLs can also signal through NCAM independently of RET (Paratcha et al. 2003) (Figure 2.2.B). NCAM plays an important role in neurodevelopment, regeneration and synaptic plasticity by mediating cell adhesion to other cells and components of the extracellular matrix. GFLs, but not other NTFs, can bind directly to NCAM via its third Ig domain (Paratcha et al. 2003; Sjöstrand et al. 2007; Nielsen et al. 2009). However, high-affinity binding of a GFL and downstream signaling through NCAM require the presence of GFR $\alpha$  co-receptor (Paratcha et al. 2003). Binding of GFL-GFR $\alpha$  complex to NCAM activates intracellular Src family kinase Fyn and focal adhesion kinase (FAK) which, in turn, activate MAP kinases and cellular responses such as neurite outgrowth of hippocampal, cortical and midbrain neurons, survival of cultured dopamine neurons and stimulation of Schwann cell migration (Chao et al. 2003; Paratcha et al. 2003; Iwase et al. 2005; Cao et al. 2008a; Nielsen et al. 2009). GDNF has been shown to function as a chemoattractant factor for neuronal precursors migrating along the RMS and NCAM-mediated signaling seems to play a key role in this guidance process (Paratcha et al. 2006). A recent study suggested that NCAM can function as an alternative receptor also for ARTN (Ilieva et al. 2019). ARTN was shown to induce neuritogenesis by binding directly to NCAM and activating NCAM-associated signaling pathways in primary cerebellar neuron cultures.



**Figure 2.2. Signaling mechanisms of GDNF family ligands (GFLs).** (A) Signaling through receptor tyrosine kinase RET can occur in *cis* or in *trans*. During activation of RET in *cis*, GFL homodimer first binds to a glycosyl phosphatidylinositol (GPI)-anchored GDNF family receptor  $\alpha$  (GFR $\alpha$ ) which dimerizes (1). Then, GFL-GFR $\alpha$  complex recruits two RET molecules into the lipid raft of plasma membrane (PM) (2). This triggers dimerization of RET and autophosphorylation of its intracellular tyrosine residues which serve as docking sites for intracellular adaptor proteins (3). In *trans* signaling, the GPI-anchor of GFR $\alpha$  is cleaved by membrane-associated phospholipases (scissors) which releases soluble GFR $\alpha$  (sGFR $\alpha$ ) into extracellular space. GFL binds with high affinity to sGFR $\alpha$  (1). GFL-sGFR $\alpha$  complex activates RET outside of the lipid rafts (2). Subsequently, the receptor complex is recruited into the lipid raft (3). Inside the lipid rafts, phosphorylated RET associates with the adaptor protein FRS2 leading to the activation of Ras/MAPK pathway. Outside the rafts, RET interacts with SHC leading to the activation of PI3K/Akt pathway. GDNF can also activate intracellular Src family kinases (SFKs) signaling via GFR $\alpha$ 1 independently of RET. (B) Neural cell adhesion molecule (NCAM) is an alternative signaling receptor for GFLs in cells lacking RET. GFL binds with high affinity to GFR $\alpha$  which dimerizes (1). Then, GFL-GFR $\alpha$  complex binds to transmembrane NCAM (2) leading to signal transduction via intracellular Src family kinase Fyn and focal adhesion kinase (FAK), and ultimately, the activation of MAP kinases (3). (C) Extracellular matrix-bound GFLs can also signal through a transmembrane heparan sulfate (HS) proteoglycan syndecan-3, independently of GFR $\alpha$ , RET or NCAM. Immobilized GFLs bind to the HS side chains of syndecan-3. The cytoplasmic domain activates intracellular SFKs and cortactin-mediated signaling cascades regulating the cytoskeleton. One syndecan-3 can bind several GFLs simultaneously. Figure drawn by the author.

#### 2.1.3.4 Signaling via syndecan-3

The distribution of GDNF and NRTN in the brain parenchyma is restricted by binding to heparan sulfate proteoglycans of the extracellular matrix (Lin et al. 1994; Hamilton et al. 2001; Rickard et al. 2003; Piltonen et al. 2009; Bespalov et al. 2011). Bespalov and his colleagues showed that immobilized, extracellular matrix-bound GDNF, NRTN and ARTN can also signal through a novel receptor, the transmembrane heparan sulfate proteoglycan syndecan-3, that is expressed on neuronal cells (Bespalov et al. 2011) (Figure 2.2.C). Immobilized GFLs bind with high affinity to the heparan sulfate side chains of syndecan-3 which activates intracellular SFK-mediated signaling cascades. As syndecan-3 has several heparan sulfate chains, one syndecan-3 can simultaneously bind multiple GFL homodimers acting as a high affinity and high capacity receptor for GFLs. GFL–syndecan-3 interaction promotes hippocampal neurite outgrowth and migration of cortical  $\gamma$ -aminobutyric acid (GABA) containing neurons, and thus may play a distinctive role in the embryonic development of these brain areas. In addition to direct intracellular signal transduction, syndecan-3, and other heparan sulfate proteoglycans, may modulate conventional signaling through GFR $\alpha$ -RET or NCAM by concentrating and presenting diffusible GFLs to the receptors. GFLs may have a dual mode of action: as diffusible soluble proteins, they prefer signaling via conventional receptors GFR $\alpha$ -RET and NCAM, whereas extracellular matrix-bound GFLs seem to signal through syndecan-3 independently of GFR $\alpha$ , RET or NCAM.

#### 2.1.3.5 Expression and functions of GFLs, GFR $\alpha$ s and RET

Following the initial observation of the effects of GDNF on midbrain dopamine neurons, several studies have shown its neurotrophic actions in other neuronal populations of the CNS and PNS. For example, GDNF supports the survival of noradrenergic neurons of the locus coeruleus (Arenas et al. 1995), basal forebrain cholinergic neurons (Williams et al. 1996), facial and spinal cord motor neurons (Henderson et al. 1994; Oppenheim et al. 1995; Yan et al. 1995) and peripheral sympathetic, parasympathetic and sensory neurons (Buj-Bello et al. 1995; Ebendal et al. 1995; Trupp et al. 1995). The biological effects on such a broad spectrum of neuronal populations suggest widespread expression of GDNF and its receptors throughout the CNS, PNS and non-neuronal tissues.

Although all GFLs activate the same downstream signaling pathways through RET, the selectivity in their biological effects is thought to be due to differential expression patterns of GFLs and their cognate GFR $\alpha$  co-receptors. GFL, GFR $\alpha$  and RET expression levels have been investigated using Northern blotting, *in situ* hybridization, RT-PCR and immunohistochemical techniques. The neuronal and non-neuronal expression of GFLs, GFR $\alpha$ s and RET in developing and adult rodents together with their physiological main functions are summarized in Table 2.1. In general, the levels and temporospatial expression patterns of GFLs are strictly regulated (Mogi et al. 2001). GFLs and their receptors are more prominently expressed during embryonic development as compared to adult animals (Golden et al. 1999). In the mature brain, GFR $\alpha$  receptors show wider expression than GFLs or RET (Nosrat et al. 1997; Trupp et al. 1997; Ortega-de San Luis and Pascual 2016). When RET and GFR $\alpha$  are expressed in the same tissue, such as in the SN or spinal cord, the receptors are able to interact in *cis*. However, the expression pattern of GFR $\alpha$  co-receptors does

not always match with that of RET which suggests in *trans* mode of interaction between the receptors or that GFLs convey RET-independent signaling via alternative receptors as described above.

Only GDNF and NRTN are expressed in the CNS of adult rodents which makes them to be of specific interest in regards of potential disease-modifying approaches for neurodegenerative diseases. Attempts to reveal the cellular basis of the relatively abundant expression of GDNF in postnatal striatum have shown that in the normal brain, neurons are the principal source of GDNF (Oo et al. 2005). More precisely, GDNF is mainly synthesized by striatal interneurons (Bizon et al. 1999; Hidalgo-Figueroa et al. 2012). The vast majority (approximately 95%) of the GDNF-expressing cells in the striatum are parvalbumin (PV) -positive GABAergic interneurons and the remaining 5% are either cholinergic or somatostatin-positive interneurons.

Upregulation of GDNF has been reported in several nigrostriatal injury models (Batchelor et al. 1999; Liberatore et al. 1997; Yurek and Fletcher-Turner 2002, 2001). Upon injury, glial cells become the predominant source of GDNF as well as other growth factors like NGF, NT-3 and FGF (Bresjanac and Antauer 2000; Nakagawa and Schwartz 2004; Nakagawa et al. 2005; Chen et al. 2006). The switch in the production of NTFs from neurons to glial cells may be part of the local mechanisms that aim to protect neurons and promote their regeneration. The cross-talk between damaged neurons and glia inducing the glial GDNF expression seems to be mediated via pro-inflammatory cytokines IL-1 $\beta$ , IL-6, interferon- $\gamma$  (IFN- $\gamma$ ), TNF- $\alpha$  and TNF- $\beta$ , FGF (Appel et al. 1997; Verity et al. 1998; Verity et al. 1999; Kuno et al. 2006; Saavedra et al. 2007) and endothelin-1 (Koyama et al. 2003a, 2003b).

Endogenous GDNF does not affect the normal embryonic development of brainstem noradrenergic and midbrain dopaminergic neurons (Moore et al. 1996; Sánchez et al. 1996). However, the physiological role of endogenous GDNF in the maintenance of normal adult brain catecholaminergic neurons has remained poorly studied because of neonatal mortality of GDNF full knockout mice due to renal and ENS agenesis. Interestingly, there is an ongoing debate whether GDNF is vital or dispensable for the survival of catecholaminergic neurons of the substantia nigra pars compacta (SNpc) and the locus coeruleus (Enterría-Morales et al. 2020; Kopra et al. 2015; Kumar et al. 2015; Pascual et al. 2008; Pascual and López-Barneo 2015). In adult mice, GDNF overexpression from the native locus seems to exert trophic effects on the nigrostriatal dopamine system and enhance its function (Kumar et al. 2015).

Table 2.1. GDNF family ligands (GFL), GDNF family ligand receptors (GFR $\alpha$ ) and RET: expression in rodents and physiological main functions.

	Expression	Main functions	References
GDNF	<p><b>CNS:</b> cortical areas, STR, NAcc, Hc, thalamic nuclei, hypothalamic nuclei, OB, olfactory tubercle, septum, midbrain, PAG, cerebellum, pons (incl. pedunculopontine nucleus and locus coeruleus), medulla, spinal cord, pituitary gland, pineal gland*, wall of the 4<sup>th</sup> ventricle</p> <p><b>PNS:</b> sensory neurons, sympathetic neurons, Schwann cells</p> <p><b>Peripheral tissues:</b> skeletal muscle, gonads, prostate, lung, liver, adrenal gland, stomach, intestine, spleen, heart, thymus, thyroid, salivary glands, skin, kidney*, bone*, teeth*, ear*, nose*, tongue*, eye*, urogenital system*, digestive system*, limb buds*, cartilage*, blood*</p>	<ul style="list-style-type: none"> <li>Originally discovered as a potent survival-promoting factor for embryonic midbrain dopamine neurons in culture</li> <li>Essential for the prenatal development of ENS and kidneys</li> <li>Essential for the development of peripheral sensory, sympathetic and parasympathetic neurons</li> <li>Plays a crucial role in spermatogenesis</li> <li>Supports the development and survival of spinal motoneurons, hindbrain noradrenergic neurons, midbrain dopamine neurons, basal forebrain cholinergic neurons and cerebellar Purkinje cells</li> <li>May promote the insertion and stabilization of postsynaptic receptors in the neuromuscular junction</li> <li><i>GDNF</i><sup>-/-</sup> genotype leads to perinatal death due to renal agenesis and the absence of enteric neurons</li> </ul>	<p>(Lin et al. 1993), (Scharr et al. 1993), (Henderson et al. 1994), (Springer et al. 1994), (Arenas et al. 1995), (Bowenkamp et al. 1995), (Buj-Bello et al. 1995), (Choi-Lundberg and Bohn 1995), (Ebendal et al. 1995), (Li et al. 1995), (Mount et al. 1995), (Oppenheim et al. 1995), (Springer et al. 1995), (Tomac et al. 1995), (Trupp et al. 1995), (Yan et al. 1995), (Hellmich et al. 1996), (Moore et al. 1996a), (Nosrat et al. 1996), (Pichel et al. 1996), (Sánchez et al. 1996), (Suvanto et al. 1996), (Vega et al. 1996), (Sainio et al. 1997), (Trupp et al. 1997), (Widenfalk et al. 1997), (Enomoto et al. 1998), (Golden et al. 1998), (Heuckeroth et al. 1998), (Fundin et al. 1999), (Golden et al. 1999), (Baudet et al. 2000), (Enomoto et al. 2000), (Garcès et al. 2000), (Meng et al. 2000), (Mikaels et al. 2000), (Worley et al. 2000), (Young et al. 2001), (Kramer et al. 2006), (Naughton et al. 2006), (Wang et al. 2010), (Savitt et al. 2012), (Ortega-de San Luis and Pascual 2016)</p>
Neurturin (NRTN)	<p><b>CNS:</b> cortical areas, STR, Hc, thalamic nuclei, hypothalamic nuclei, ventral midbrain, cerebellum, pituitary gland, septum*, brainstem nuclei*, pineal gland*</p> <p><b>PNS:</b> sensory neurons, sympathetic neurons, retina</p> <p><b>Peripheral tissues:</b> gonads, prostate, kidney, heart, bladder, urethra, skin, GI-tract, liver, lung, thymus, exocrine glands, skeletal muscle*, teeth*, digestive system*, sensory organs*</p>	<ul style="list-style-type: none"> <li>Originally identified on the basis of survival-promoting effects on sympathetic neurons in culture</li> <li>Essential for the normal development and survival of parasympathetic neurons</li> <li>Essential for the proper development, maintenance and function of ENS</li> <li>Contributes to food digestion by ensuring proper intestinal motility and secretion of pancreatic enzymes and saliva</li> <li>Promotes the survival and neurite outgrowth of spinal motoneurons and embryonic basal forebrain cholinergic neurons and survival of sensory neurons and midbrain dopaminergic neurons</li> <li><i>NRTN</i><sup>-/-</sup> mice are viable and fertile, no gross developmental defects except ptosis due of lack of parasympathetic innervation of the lacrimal gland</li> </ul>	<p>(Kotzbauer et al. 1996), (Klein et al. 1997), (Widenfalk et al. 1997), (Golden et al. 1998), (Heuckeroth et al. 1998), (Hörger et al. 1998), (Bilak et al. 1999), (Forgie et al. 1999), (Fundin et al. 1999), (Golden et al. 1999), (Heuckeroth et al. 1999), (Jomary et al. 1999), (Rossi et al. 1999), (Taraviras et al. 1999), (Åkerud et al. 1999), (Baudet et al. 2000), (Enomoto et al. 2000), (Hiltunen et al. 2000), (Laurikainen et al. 2000), (Rossi et al. 2000), (Golden et al. 2003), (Rossi et al. 2003), (Cho et al. 2004b), (Mabe et al. 2006)</p>
Artemin (ARTN) also termed enovin and neublastin	<p><b>CNS:</b> <i>not detected</i></p> <p><b>PNS:</b> nerve roots of DRG*, immature Schwann cells*, sympathetic neurons*, along the routes of sympathetic neuroblast migration and along sympathetic axonal projections*</p> <p><b>Peripheral tissues:</b> blood vessels (smooth muscle cells), esophagus*, stomach*, pancreas*</p>	<ul style="list-style-type: none"> <li>Originally identified as the ligand for the orphan GFR<math>\alpha</math>3–RET receptor</li> <li>Vascular-derived NTF crucial for the migration and axonal outgrowth of sympathetic neuroblasts and development of target tissue sympathetic innervation</li> <li>Promotes the survival of sensory and midbrain dopamine neurons</li> <li>Modulates the sensitivity of sensory neurons to noxious stimuli</li> <li><i>ARTN</i><sup>-/-</sup> mice are viable and fertile, no gross developmental defects except ptosis due of lack of sympathetic innervation to the superior tarsus muscle</li> </ul>	<p>(Baloh et al. 1998b), (Nishino et al. 1999), (Baudet et al. 2000), (Rosenblad et al. 2000), (Andres et al. 2001), (Enomoto et al. 2001), (Honma et al. 2002), (Wang et al. 2008), (McIlvried et al. 2010), (Nivlet et al. 2016)</p>
Persephin (PSPN)	<p><b>CNS:</b> cortical areas*, Hc*, STR*, diencephalon*, midbrain*, pons*, medulla*, cerebellum*, spinal cord*, astrocytes*</p> <p><b>PNS:</b> sympathetic neurons (SCG)*, sensory neurons (DRG)*, sciatic nerve*, optic nerve*, motoneurons*</p> <p><b>Peripheral tissues:</b> fat tissue, adrenal gland, heart, kidney, liver, skin, spleen*, skeletal muscle*, bone*, testicle*</p> <p><i>In general, PSPN is expressed at very low levels in adult rodents</i></p>	<ul style="list-style-type: none"> <li>Originally discovered as the result of its homology to GDNF and NRTN</li> <li>Regulates the function of thyroid C cells, their calcitonin production and bone formation in newborn and juvenile mice</li> <li>May act as a circulating growth factor due to inability to bind heparan sulfate side chains of the extracellular matrix</li> <li>Can modulate glutamate-mediated excitotoxicity in the CNS and supports the survival of motoneurons and sympathetic neurons</li> <li><i>In vitro</i>, promotes the survival and neurite outgrowth of embryonic midbrain dopamine and basal forebrain cholinergic neurons</li> <li><i>PSPN</i><sup>-/-</sup> mice are viable and fertile, no gross developmental defects</li> </ul>	<p>(Enokido et al. 1998), (Heuckeroth et al. 1998), (Jaszai et al. 1998), (Milbrandt et al. 1998), (Tomac et al. 2002), (Åkerud et al. 2002), (Golden et al. 2003), (Lindfors et al. 2006), (Bespalov et al. 2011)</p>

GFRα	GFRα1 also termed GDNFR-α, TrnR1 and RETL1	<p><b>CNS:</b> cortical areas, STR, NAcc, Hc, OB, thalamic nuclei, hypothalamic nuclei, OT, amygdala, septum, habenular nuclei, ventral pallidum, midbrain (incl. SNpc, SNr, VTA and dorsal Raphe nucleus), PAG, cerebellum, pons (incl. pedunculopontine nucleus and locus coeruleus), spinal cord</p> <p><b>PNS:</b> sensory neurons, Schwann cells, ENS, motoneurons, sympathetic neurons*, parasympathetic neurons*, retina*</p> <p><b>Peripheral tissues:</b> gonads, prostate, GI-tract, liver, kidney, bladder, urethra, heart, lung, spleen, digestive system*, skin*, bone*, teeth*, skeletal muscle*, inner ear*, endocrine glands*, salivary glands*</p>	<ul style="list-style-type: none"> <li>• Cognate co-receptor for GDNF</li> <li>• Shows low-affinity cross-reactivity with NRTN and ARTN <i>in vitro</i></li> <li>• Association with NCAM potentiates high-affinity binding of GDNF to NCAM</li> <li>• Interaction with NCAM downregulates homophilic NCAM binding and cell adhesion</li> <li>• Ligand-induced cell adhesion molecule in the presence of GDNF</li> <li>• <i>GFRα1<sup>-/-</sup></i> mice die early postnatally due to renal agenesis and deficits in the ENS</li> </ul>	<p>(Baloh et al. 1997), (Jing et al. 1997), (Klein et al. 1997), (Nosrat et al. 1997), (Sanicola et al. 1997), (Trupp et al. 1997), (Widenfalk et al. 1997), (Baloh et al. 1998b), (Cacalano et al. 1998), (Enomoto et al. 1998), (Glazner et al. 1998), (Golden et al. 1998), (Masure et al. 1998), (Yu et al. 1998b), (Fundin et al. 1999), (Golden et al. 1999), (Baudet et al. 2000), (Bennett et al. 2000), (Enomoto et al. 2000), (Hiltunen et al. 2000), (Mikaels et al. 2000), (Rossi et al. 2000), (Garcès et al. 2001), (Sarabi et al. 2001), (Paratcha et al. 2003), (Sarabi et al. 2003), (Cho et al. 2004a), (Ledda et al. 2007), (Omodaka et al. 2014), (Ortega-de San Luis and Pascual 2016)</p>
	GFRα2 also termed GDNFR-β, TrnR2, RETL2 and NTNR-α	<p><b>CNS:</b> cortical areas, Hc, OB, thalamic nuclei, hypothalamic nuclei, OT, amygdala, septum, nucleus basalis of Meynert, midbrain (incl. SNpc, SNr, VTA and dorsal Raphe nucleus), PAG, cerebellum, brainstem (incl. locus coeruleus), spinal cord, pineal gland, pituitary gland, STR*, habenular nuclei*</p> <p><b>PNS:</b> sensory neurons, sympathetic neurons, parasympathetic neurons, Schwann cells, ENS, retina, motoneurons*</p> <p><b>Peripheral tissues:</b> gonads, GI-tract, heart, lung, spleen, thyroid gland, kidney, placenta, pancreas, urogenital system*, skin*, bone*, teeth*, skeletal muscle*, endocrine glands*, salivary glands*, sensory organs*</p>	<ul style="list-style-type: none"> <li>• Cognate co-receptor for NRTN</li> <li>• Shows low-affinity cross-reactivity with GDNF <i>in vitro</i></li> <li>• Association with NCAM potentiates high-affinity binding of NRTN to NCAM <i>in vitro</i></li> <li>• <i>GFRα2<sup>-/-</sup></i> mice are viable and fertile, but have ptosis and grow poorly due to deficits in the enteric and parasympathetic nervous system</li> </ul>	<p>(Baloh et al. 1997), (Jing et al. 1997), (Klein et al. 1997), (Sanicola et al. 1997), (Widenfalk et al. 1997), (Golden et al. 1998), (Horger et al. 1998), (Masure et al. 1998), (Naveilhan et al. 1998), (Trupp et al. 1998), (Yu et al. 1998b), (Fundin et al. 1999), (Golden et al. 1999), (Jomary et al. 1999), (Rossi et al. 1999), (Baudet et al. 2000), (Bennett et al. 2000), (Enomoto et al. 2000), (Hiltunen et al. 2000), (Mikaels et al. 2000), (Rossi et al. 2000), (Garcès et al. 2001), (Paratcha et al. 2003), (Cho et al. 2004b), (Mabe et al. 2006), (Omodaka et al. 2014), (Ishida et al. 2016)</p>
	GFRα3	<p><b>CNS:</b> OB, Hc, cerebellum, <i>expressed at low levels in the CNS</i></p> <p><b>PNS:</b> sensory neurons, sympathetic neurons, Schwann cells, sympathetic neuroblasts*, peripheral nerves*, retina*</p> <p><b>Peripheral tissues:</b> epidermis, thymus, heart, lung, intestine, pancreas, spleen, ovary, kidney, adrenal medulla*, skeletal muscle*, salivary gland*, liver*</p>	<ul style="list-style-type: none"> <li>• Cognate co-receptor for ARTN</li> <li>• Shows low-affinity cross-reactivity with GDNF <i>in vitro</i></li> <li>• <i>GFRα3<sup>-/-</sup></i> mice are viable and fertile, no gross abnormalities other than ptosis</li> </ul>	<p>(Baloh et al. 1998a), (Masure et al. 1998), (Naveilhan et al. 1998), (Nomoto et al. 1998), (Trupp et al. 1998), (Widenfalk et al. 1998), (Worby et al. 1998), (Yu et al. 1998b), (Fundin et al. 1999), (Nishino et al. 1999), (Baudet et al. 2000), (Bennett et al. 2000), (Hiltunen et al. 2000), (Orozco et al. 2001), (Honma et al. 2002), (Omodaka et al. 2014), (Wong et al. 2015), (Nivlet et al. 2016)</p>
	GFRα4	<p><b>CNS:</b> cortical areas, Hc, OB, habenular nuclei, ventral midbrain (incl. SNpc and VTA), cerebellum, spinal cord, pituitary gland</p> <p><b>PNS:</b> sympathetic and parasympathetic ganglia</p> <p><b>Peripheral tissues:</b> thyroid gland, parathyroid gland, adrenal medulla, heart, testicle</p>	<ul style="list-style-type: none"> <li>• Cognate co-receptor for PSPN</li> <li>• Shows low-affinity cross-reactivity with NRTN <i>in vitro</i></li> <li>• Association with NCAM potentiates high-affinity binding of PSPN to NCAM <i>in vitro</i></li> <li>• <i>GFRα4<sup>-/-</sup></i> mice are viable and fertile, no gross developmental defects</li> </ul>	<p>(Enokido et al. 1998), (Lindahl et al. 2000), (Masure et al. 2000), (Åkerud et al. 2002), (Paratcha et al. 2003), (Lindfors et al. 2006)</p>
RET		<p><b>CNS:</b> dopamine neurons of midbrain (incl. SN and VTA), serotonergic neurons of dorsal Raphe nucleus, cholinergic neurons of basal forebrain, OB, thalamic and hypothalamic nuclei, amygdala, septum, PAG, cerebellum, pons (incl. locus coeruleus), cranial and spinal motoneurons, <i>low levels in the STR</i></p> <p><b>PNS:</b> ENS, sympathetic neurons, parasympathetic neurons, sensory neurons, retina</p> <p><b>Peripheral tissues:</b> testicle, salivary gland, GI-tract, adrenal medulla, thyroid gland, heart, lymphoid organs, kidney*, inner ear*, teeth*</p>	<ul style="list-style-type: none"> <li>• Transmembrane signaling receptor for GFL-GFRα complex</li> <li>• <i>Ret</i> gene originally discovered as an oncogene: gain-of-function mutations cause dominant cancer syndromes MEN2A, MEN2B and FMTC</li> <li>• Pivotal for the normal development and maintenance of ENS: loss-of-function mutations cause Hirschsprung's disease (intestinal obstruction/ megacolon)</li> <li>• Essential for kidney organogenesis and spermatogenesis</li> <li>• Essential for the development of sympathetic and parasympathetic neurons</li> <li>• Promotes the survival, axonal guidance and maturation of motoneurons</li> <li>• Regulates long-term maintenance of the nigrostriatal dopamine system</li> <li>• <i>RET<sup>-/-</sup></i> mice die at birth due to renal agenesis and deficits in the ENS</li> </ul>	<p>(Takahashi et al. 1985), (Pachnis et al. 1993), (Schuchardt et al. 1994), (Tsuzuki et al. 1995), (Durbec et al. 1996b), (Schuchardt et al. 1996), (Trupp et al. 1996), (Nosrat et al. 1997), (Trupp et al. 1997), (Glazner et al. 1998), (Golden et al. 1998), (Yu et al. 1998b), (Golden et al. 1999), (Taraviras et al. 1999), (Bennett et al. 2000), (Enomoto et al. 2000), (Garcès et al. 2000), (Hiltunen et al. 2000), (Lindahl et al. 2000), (Enomoto et al. 2001), (Garcès et al. 2001), (Golden et al. 2003), (Jain et al. 2004), (Shakya et al. 2005), (Kramer et al. 2006), (Plaza-Menacho et al. 2006), (Kramer et al. 2007), (Mijatovic et al. 2007), (Baudet et al. 2008), (Jijiwa et al. 2008), (Uesaka et al. 2008), (The Human Protein Atlas 2020)</p>

CNS, central nervous system; PNS, peripheral nervous system; ENS, enteric nervous system; DRG, dorsal root ganglia; SCG, superior cervical ganglia; STR, striatum; NAcc, nucleus accumbens; Hc, hippocampus; GI, gastrointestinal; OB, olfactory bulb; OT, olfactory tubercle; PAG, periaqueductal gray; SNpc, substantia nigra pars compacta; SNr, substantia nigra pars reticulata; VTA, ventral tegmental area; FMTC, familial medullary thyroid carcinoma; MEN2, multiple endocrine neoplasia type 2

\* expression reported only during embryonic or early postnatal development, but not in adult rodent

### 2.1.4 CDNF/MANF family of neurotrophic factors

More recently discovered NTFs, MANF and its paralog CDNF are structurally and functionally distinct from the classical NTFs (Petrova et al. 2003; Lindholm et al. 2007; Lindahl et al. 2017). MANF (also known as arginine-rich, mutated in early stage tumors; ARMET) and CDNF (also known as ARMET-like 1; ARMETL1) are both present in vertebrates, but invertebrates, including *D. melanogaster* and *C. elegans*, express only one homologous protein that is more closely related to MANF (DmMANF) (Palgi et al. 2009). Together CDNF and MANF form an evolutionary conserved family of NTFs that shows classical neurotrophic properties such as promoting the survival and repair of midbrain dopamine neurons *in vitro* and in animal models of PD, protecting cardiac myocytes and cortical neurons against ischemia, and enhancing peripheral nerve recovery in a sciatic nerve transection model (Petrova et al. 2003; Lindholm et al. 2007; Tadimalla et al. 2008; Airavaara et al. 2009; Voutilainen et al. 2009; Airavaara et al. 2010; Voutilainen et al. 2011; Airavaara et al. 2012; Glembotski et al. 2012b; Cheng et al. 2013a; Cordero-Llana et al. 2015a; Liu et al. 2018; Zhang et al. 2018a; Zhang et al. 2018b).


During embryonic development, MANF signaling seems to be necessary for the proper neuronal differentiation, neuroblast migration and neurite outgrowth in the developing mouse brain (Tseng et al. 2018, 2017). In line, studies in *D. melanogaster* and *C. elegans* suggest an important role for DmMANF in the maturation and survival of dopamine neurons and in the maintenance of ER homeostasis (Palgi et al. 2009; Richman et al. 2018). In zebrafish, MANF knockdown causes a decrease in brain dopamine level and impairs the development of tyrosine hydroxylase (TH) expressing neurons (Chen et al. 2012). As for CDNF, knockout mice show age-dependent aberrant functions of the brain dopamine system such as slower dopamine reuptake via dopamine transporter (DAT) and increased amphetamine-induced dopamine release in the striatum (Lindahl et al. 2020). However, the number of dopamine neurons in the SN and dopamine level in the striatum remain unaltered. Interestingly, CDNF deficiency leads to age-dependent loss of enteric neurons in the submucosal plexus due to increased autophagy and neuronal degeneration. To date, the structure and expression pattern of CDNF and MANF are well-studied but their cell surface receptors and signaling pathways remain to be unraveled.

#### 2.1.4.1 Structure and secretion of MANF and CDNF

CDNF and MANF are structurally unrelated to other NTFs. The amino acid sequences of mature human CDNF and MANF are 59% identical (Petrova et al. 2003; Lindholm et al. 2007; Lindahl et al. 2017). Their primary structure contains an N-terminal signal sequence (pre-sequence) that directs them to the ER during protein synthesis and allow access to the secretory pathway. Cleavage of the pre-sequence results in a mature protein that can be secreted via the canonical COPII-mediated ER-Golgi pathway (Apostolou et al. 2008; Oh-hashii et al. 2012; Norisada et al. 2016). Secreted CDNF and MANF seem to exert their trophic actions both in an autocrine and paracrine fashion but their binding to a transmembrane receptor has remained obscure until today (Apostolou et al. 2008; Lindahl et al. 2017). Although CDNF and MANF can be secreted, they are mostly retained intracellularly within the ER unlike classical NTFs that are exclusively secretory

proteins (Lindholm and Saarma 2010). In the ER, they seem to support protein folding and mitigate ER stress.

Human MANF is a glycosylated protein consisting of 179 amino acids and having a molecular mass of 20 kDa (Figure 2.3A) (Petrova et al. 2003). Its primary structure contains a 21-amino acid-long signal sequence which is cleaved to form 158 amino acids long mature protein. The secondary structure of the protein is dominated by  $\alpha$ -helices and random coils and virtually lacks  $\beta$ -sheets. Human CDFN is 187 amino acids long, glycosylated protein with a calculated molecular mass of 18 kDa (Lindholm et al. 2007; Apostolou et al. 2008; Sun et al. 2011). CDFN has a signal sequence of 26 amino acids, the cleavage of which forms a 161-amino acid-long mature protein. Similarly to MANF, the secondary structure of CDFN contains mostly  $\alpha$ -helices. A characteristic feature of the primary structure of CDFN and MANF is eight conserved cysteine residues (Lindholm et al. 2007; Petrova et al. 2003). Four disulphide bridges are formed between the cysteine residues stabilizing the tertiary structure of the mature proteins.



**Figure 2.3. Schematic structure of human MANF and CDFN.** (A) Schematic picture of the primary structure of human MANF and CDFN. The N-terminal (saposin-like) domain of MANF and CDFN is indicated in blue, and the C-terminal (SAP-like) domain in orange. Vertical yellow bars depict conserved cysteine residues, and they are numbered according to the sequence of mature protein. The formation of disulphide bridges is shown as black connecting lines below the sequence of MANF. The ER retention signals of MANF and CDFN (RTDL and KTEL, respectively) are marked with pink. (B) Solution structure of human MANF. MANF has two domains: saposin-like N-terminal domain containing five  $\alpha$ -helices ( $\alpha$ 1- $\alpha$ 5) and  $3_{10}$  helix in the linker region, and SAP-like C-terminal domain containing two  $\alpha$ -helices ( $\alpha$ 6 and  $\alpha$ 7). Yellow sticks show the formation of disulphide bridges in the tertiary structure. The CXXC motif ( $^{127}$ CKGC $^{130}$ ) and ER retention signal ( $^{155}$ RTDL $^{158}$ ) are indicated in the C-terminal domain. N - amino-terminus; C - carboxy-terminus. Adapted with permission from Elsevier: Neurobiology of Disease, Lindahl et al. (2017), © Elsevier 2016.



The tertiary structures of CDNF and MANF closely resemble each other. They consist of two domains connected to each other with a flexible linker region (Lindholm et al. 2007; Parkash et al. 2009; Hellman et al. 2011; Latge et al. 2015). N-terminal domain has a globular conformation containing five  $\alpha$ -helices ( $\alpha$ 1- $\alpha$ 5),  $3_{10}$  helix in the linker region and a hydrophobic core. Three out of four disulphide bridges are located in the N-terminal domain as depicted in Figure 2.3B. C-terminal domain is partially unstructured; it contains two  $\alpha$ -helices ( $\alpha$ 6 and  $\alpha$ 7) and one internal disulphide bridge between the two cysteine residues in CXXC motif ( $^{132}\text{CRAC}^{135}$  in CDNF and  $^{127}\text{CKGC}^{130}$  in MANF). This CXXC motif in the loop connecting the helices  $\alpha$ 6 and  $\alpha$ 7 seems to be important for the functionality of the proteins. Mätlik and coworkers showed that mutating the CKGC motif abolishes the survival-promoting activity of MANF when overexpressed intracellularly in cultured sympathetic or sensory neurons, or when applied extracellularly in a rat model of focal cerebral ischemia (Mätlik et al. 2015). At the very end of the C-terminus, CDNF and MANF have conserved four amino acid sequences ( $^{158}\text{KTEL}^{161}$  in CDNF and  $^{155}\text{RTDL}^{158}$  in MANF) which resemble the canonical ER retention signal KDEL (Glembotski et al. 2012; Henderson et al. 2013; Oh-hashii et al. 2012; Raykhel et al. 2007). Human has three KDEL receptors (KDEL1-3) that function in the Golgi. KDELs recognize KDEL-like sequences in the C-terminus of proteins and mediate their retrograde trafficking from the Golgi back to the ER. Henderson and colleagues suggested that KDELs, accumulated to the plasma membrane in ER-stressed cells, could mediate the cell surface binding of MANF (possibly CDNF too) through the interaction between the C-terminal RTDL sequence and KDELs (Henderson et al. 2013).

MANF retention to the ER is also dependent on another ER resident protein, glucose-regulated protein 78 kDa (GRP78; also known as binding immunoglobulin protein, BiP), the expression of which closely resembles the expression of MANF in mouse tissues (Mizobuchi et al. 2007; Glembotski et al. 2012; Oh-hashii et al. 2012). MANF interacts directly with GRP78 and the interaction has been shown to be calcium dependent. Depletion of ER calcium storages e.g. by thapsigargin, an inhibitor of SERCA (sarco/endoplasmic reticulum calcium transporting ATPase), triggers MANF secretion independently of the RTDL sequence. Taken together, if the ER calcium concentration is normal, both GRP78 and KDEL seem to participate in retaining MANF in the ER. Upon ER calcium depletion, MANF dissociates from GRP78 leading to its increased secretion although the interaction with KDEL remains unaffected. Thus, the RTDL sequence is anticipated to function as a weak retention motif fine-tuning the secretion of MANF. Overexpression of GRP78 and KDEL1 in HEK293 cells similarly decreased the secretion of MANF and CDNF suggesting that the mechanisms regulating the secretion of CDNF and MANF are fundamentally similar (Glembotski et al. 2012; Oh-hashii et al. 2012; Henderson et al. 2013; Norisada et al. 2016).

#### 2.1.4.2 Expression of MANF and CDNF

MANF is widely expressed both in neuronal and non-neuronal tissues starting from early stages of embryonic development until adulthood (Lindholm et al. 2008; Wang et al. 2014). Immunohistochemical studies show that in the rodent brain the expression of MANF is mainly neuronal (Lindholm et al. 2008; Shen et al. 2012; Wang et al. 2014). In the CNS of the adult mouse, MANF expression is abundant in several brain regions including the olfactory bulb, cortical areas, hippocampus, several hypothalamic and thalamic nuclei, pons, medulla, cerebellar Purkinje cells,

spinal cord and neural progenitors of the SVZ (Lindholm et al. 2008; Wang et al. 2014; Tseng et al. 2018). In the striatum and SN, only low to intermediate levels of MANF were detected both from the adult brain and during embryonic development. In the SNpc, MANF immunofluorescence co-localized with the TH-immunoreactive (ir) cells indicating MANF expression in dopamine neurons (Lindholm et al. 2008). However, MANF-positive neurons were more prevalent in the SN pars reticulata than in pars compacta. Wang and colleagues suggested that in the rat brain MANF expression is developmentally regulated with the expression being highest within the first two weeks after birth and decreasing upon aging (Wang et al. 2014).

Apart from the CNS, MANF is broadly expressed in peripheral tissues of adult and embryonic mice (Mizobuchi et al. 2007; Lindholm et al. 2008; Lindahl et al. 2014). The expression is ample especially in secretory cells and tissues, including choroid plexus, pancreatic exocrine acinar cells and endocrine  $\beta$  cells, salivary glands, liver and testis, suggesting an important role of MANF in regulating protein homeostasis in cells with a high rate of protein production. Lower levels of MANF are detected in the lung, skeletal muscles, kidney and heart (Lindholm et al. 2008). The indispensable role of MANF for the survival and normal function of pancreas was shown in MANF knockout mice (Lindahl et al. 2014). MANF deficiency caused progressive loss of functional  $\beta$  cells and chronic UPR activation in pancreatic islets, resulting in insufficient insulin secretion, increased blood glucose level and diabetic phenotype. Importantly, MANF is widely expressed in spleen and different types of immune cells (Cheng et al. 2014; Liu et al. 2015b; Neves et al. 2016). For example, MANF levels in peripheral leukocytes and synovial tissues are highly up-regulated in response to arthritis or other inflammatory diseases (Chen et al. 2015). Recently, MANF expression has also been detected in human blood serum (Galli et al. 2016) as well as in rodent and human optic nerves and retina where it promotes the survival of retinal ganglion and photoreceptor cells (Gao et al. 2017a; Gao et al. 2017b; Lu et al. 2018).

Similarly to MANF, CDFN transcripts are widely expressed both in the human and mouse brain as well as in peripheral non-neuronal tissues (Lindholm et al. 2007). The expression levels of CDFN, however, appear to be generally lower as compared to MANF. In immunohistochemical analyses, CDFN immunoreactivity was detected in somas where it co-localized with neuronal markers NeuN and Tuj1 indicating that the expression of CDFN is neuronal (Lindholm et al. 2007; Zhou et al. 2016). In addition, CDFN was shown to co-localize with glial fibrillary acidic protein (GFAP), a marker for astrocytes, in rat primary hippocampal cell culture. In the adult mouse brain, CDFN-positive cells were observed in the brain stem, including locus coeruleus, in the cortical layers II-VI, in CA1 and CA3 pyramidal regions and dentate gyrus of the hippocampus and in the Purkinje cells of the cerebellum (Lindholm et al. 2007). In the SN, however, CDFN immunoreactivity was detected only in solitary cells which did not include TH-positive dopamine neurons. CDFN staining was weak also in the striatum. In postnatal P1 and P10 mouse brains, CDFN signal was detected in the hippocampus, thalamus, striatum and SN. Relatively high CDFN levels were detected in the adult mouse heart, skeletal muscles and testis. Overall, the expression pattern of CDFN and MANF seems to partly overlap but the intensity of their expression differs.

### 2.1.4.3 Two proposed mechanisms of action

As described above, CDFN and MANF possess a two-domain protein structure (Parkash et al. 2009; Hoseki et al. 2010; Hellman et al. 2011; Latge et al. 2015). The domains are connected via a short polypeptide linker which allows them to move flexibly in relation to each other. The independent orientation of the domains may be an important feature in the mechanism of action of CDFN and MANF since the domains seem to exert distinct functions.

#### **Anti-apoptotic effect of MANF and CDFN**

The C-terminal domain of CDFN and MANF is homologous to the SAP (SAF-A/B, Acinus, and PIAS) domain of Ku70 (C-Ku70) (Hellman et al. 2011; Hoseki et al. 2010; Latge et al. 2015; Parkash et al. 2009). Ku70 inhibits the cytoplasmic pro-apoptotic Bax (Bcl-2-associated X protein) via its SAP domain thereby preventing apoptosis (Sawada et al. 2003; Wolter et al. 1997). During apoptosis, Ku70 dissociates from Bax which is translocated into mitochondria where it promotes release of cytochrome *c* into the cytosol. Cytochrome *c*, in turn, activates caspases resulting in cell death (Green 2000). Owing to the structural homology with the SAP domain of Ku70, the C-terminal domain of CDFN or MANF may prevent apoptosis by regulating Bax and subsequent caspase activation (Hellman et al. 2011; Mei and Niu 2014). Direct interaction between CDFN or MANF and Bax, however, has not been detected suggesting that the neuroprotective activity against Bax-dependent apoptosis is not mediated by a direct interaction of the proteins (Mätlik et al. 2015).

The N-terminal domain of CDFN and MANF has a homologous structure with saposin-like proteins (SAPLIPs), a diverse family of lipid and membrane -interacting proteins with various cellular functions (Parkash et al. 2009; Hoseki et al. 2010; Hellman et al. 2011; Latgé et al. 2013; Latge et al. 2015). SAPLIPs are globular proteins containing four or five  $\alpha$ -helices and two or three disulphide bridges (Bruhn 2005). Structurally N-CDFN and N-MANF are closest to the membrane-lytic proteins granulysin and NK-lysin which contain several positively charged amino acid residues responsible for their binding to the negatively charged phospholipid heads of biological membranes (Parkash et al. 2009). Similarly, positively charged lysine and arginine residues on the surface of helices  $\alpha$ 3,  $\alpha$ 4 and  $\alpha$ 5 in N-CDFN and N-MANF may increase the membrane binding properties of the proteins. Lipid-binding could potentially mediate the initial cellular interaction and internalization of the proteins. Nevertheless, in spite of the structural propensity, lipid interactions of CDFN and MANF have not been demonstrated so far (Lindahl et al. 2017).


#### **Regulation of ER stress by MANF and CDFN**

Different cellular perturbations, for example the expression of mutated proteins or viral infections, can lead to the abnormal accumulation of unfolded or misfolded proteins in the lumen of ER. The misfolded proteins alter the function of ER and cause disturbances in cellular proteostasis referred to as ER stress (reviewed in Ron and Walter 2007; Dufey et al. 2014). In order to cope with protein folding alterations, ER stress triggers the activation of conserved cellular signaling cascades collectively known as UPR. UPR leads to adaptive responses that suppress protein translation and translocation into the ER, enhance protein folding capacity by increasing the production of molecular chaperones and degrading misfolded proteins. Altogether, these events aim to

attenuate unfolded protein load in the ER and restore the homeostasis. UPR related cytosolic signaling cascades are activated by dissociation of GRP78 from the luminal domain of three different ER transmembrane sensor proteins: inositol-requiring enzyme 1 (IRE1), protein kinase RNA-like ER kinase (PERK) and activating transcription factor 6 (ATF6). If ER stress becomes prolonged or too intense to be resolved with adaptive changes, UPR triggers apoptotic cell death mechanisms. ER stress and UPR play an important part in the pathophysiology of many neurodegenerative diseases, such as PD or AD, which are characterized by the aggregation of misfolded proteins in the brain (Hetz and Saxena 2017).

CDNF and MANF have been shown to localize in the ER suggesting their function as ER stress and protein homeostasis regulating proteins (Apostolou et al. 2008; Cheng et al. 2013b; Mizobuchi et al. 2007; Tadimalla et al. 2008; Voutilainen et al. 2015). The promoter region of *Manf* gene contains two ER stress-responsive elements (ERSE I and ERSE II) which are recognized by ER stress-inducible transcription factors activated ATF6 and spliced X-Box-Binding Protein 1 (XBP1s) enhancing MANF transcription (Lee et al. 2003; Yamamoto et al. 2004; Mizobuchi et al. 2007; Oh-Hashi et al. 2013; Wang et al. 2018a). On the contrary, *Cdnf* gene has not been reported to contain such elements. MANF and CDNF expression levels are upregulated upon ER stress in various cell types *in vitro* and *in vivo* (Apostolou et al. 2008; Glembotski et al. 2012; Hartley et al. 2013; Mizobuchi et al. 2007; Tadimalla et al. 2008; Zhang et al. 2018). In physiological conditions, MANF and CDNF mostly localize to the luminal side of the ER, whereas under ER stress their secretion is enhanced which is an unconventional response for an ER stress-induced protein.

The ER stress regulation of MANF and CDNF is proposed to be mediated through GRP78 and its downstream UPR signaling pathways (Lindahl et al. 2017). The hypothetical mechanism of MANF in the regulation of UPR is depicted in Figure 2.4. In a rat model of focal cerebral ischemia, for example, intraventricularly administered MANF was able to rescue the neuronal loss in the hippocampus and cortex and downregulate UPR by repressing the elevated levels of GRP78, phosphorylated IRE1 and cleaved caspase-3 (Yang et al. 2014). Similarly, exogenously administered CDNF protected against A $\beta$ -induced synaptotoxicity by suppressing the increase of GRP78, phosphorylated JNK and cleaved caspase-3 in primary hippocampal cells (Zhou et al. 2016). In knockout mice, the lack of MANF led to chronic UPR activation in pancreatic islets and significant upregulation of GRP78, ATF4, CHOP and ATF6 (Lindahl et al. 2014). Recent findings by Yan and coworkers show that MANF binds to the nucleotide-binding domain of GRP78 and inhibits nucleotide exchange on GRP78 (Yan et al. 2019). Thus, MANF seems to promote protein folding in the ER by stabilizing GRP78–substrate interactions.



**Figure 2.4. Proposed mechanism for unfolded protein response (UPR) regulation by MANF.** (A) In the absence of endoplasmic reticulum (ER) stress, binding of glucose-regulated protein 78 kDa (GRP78) to the luminal domain of three ER transmembrane sensor proteins, activating transcription factor 6 (ATF6), inositol-requiring enzyme 1 (IRE1) and protein kinase RNA-like ER kinase (PERK), keeps them inactive. MANF is suggested to bind GRP78 in a Ca<sup>2+</sup> dependent manner helping to keep the UPR sensor proteins inactive. KDEL receptor (KDEL) retains MANF in the ER via the C-terminal RTDL sequence. MANF may also be retrieved from the Golgi back to the ER by KDEL. (B) Upon ER stress, depletion of Ca<sup>2+</sup> from the ER leads to dissociation of MANF from GRP78 and its secretion to the extracellular space, possibly due to increased competition in KDEL-mediated retention. GRP78 dissociates from ATF6, IRE1 and PERK and binds to accumulating misfolded proteins in the ER. This activates UPR related cytosolic signaling cascades. ATF6 is translocated to the Golgi, where it gets cleaved and activated. Active ATF6 moves into the nucleus, where it functions as a transcription factor for genes involved in protein folding, ER-associated protein degradation (ERAD) and lipid synthesis. It also upregulates the expression of GRP78, MANF, pro-apoptotic transcription factor C/EBP homologous protein (CHOP) and Xbp1 mRNA. Dissociation of GRP78 allows IRE1 to be dimerized and autophosphorylated. Endoribonuclease activity of the cytoplasmic domain of IRE1 removes an intron from Xbp1 mRNA. Spliced Xbp1s mRNA is then translated to XBP1s transcription factor, which upregulates genes for chaperons, ERAD, lipid synthesis and MANF. In addition, IRE1 can reduce translational workload by degrading ER associated mRNAs. Chronic activation of IRE1 can recruit TRAF2 and ASK1, which phosphorylates c-Jun N-terminal kinase (JNK) and leads to the nuclear translocation of NF-kB. NF-kB upregulates genes of inflammatory response, autophagy regulation and apoptosis. Dimerization and autophosphorylation of PERK followed by phosphorylation of translation initiation factor eIF2 $\alpha$  leads to general inhibition of protein synthesis and increase in the translation of transcription factor ATF4. ATF4 induces genes related to protein folding, amino acid metabolism and restoring oxidative balance, as well as CHOP. PM – plasma membrane; SERCA2 – sarco/endoplasmic reticulum Ca<sup>2+</sup>-ATPase2. Figure drawn by the author, inspired by Lindahl et al. (2017), Kim et al. (2017) and Danilova and Lindahl (2018).

## Regulation of inflammation by MANF and CDFN

Inflammation and ER stress are closely linked with each other. In many pathologies, such as neurodegenerative diseases, all three ER stress-induced UPR sensor proteins (i.e. IRE1, PERK and ATF6) participate in mediating inflammatory processes as reviewed by (Zhang and Kaufman 2008; Cao et al. 2016; Sprenkle et al. 2017). ER stress and immune cell activation share many common regulators, e.g. NF- $\kappa$ B (nuclear factor kappa-light-chain-enhancer of activated B cells), JNK and p38, forming a vicious cycle of the two signaling pathways that may exacerbate their pro-apoptotic effects. As a consequence, the neuroprotective and regenerative properties of CDFN and MANF may ultimately be due to regulation of inflammatory responses in the CNS (Zhao et al. 2014; Chen et al. 2015; Neves et al. 2016). Immune cell activation occurs at early stages of tissue reparation and its regulation is crucial to regenerative success (Sousa-Victor et al. 2018). The precise mechanisms through which CDFN and MANF can affect neuroinflammation still remain undetermined, but the studies published to date support the notion that they can reduce the secretion of pro-inflammatory cytokines and promote the pro-regenerative activation of immune cells.

MANF has been shown to exert immune modulatory functions that bias immune cells toward an anti-inflammatory phenotype. Results from Chen and colleagues suggest that MANF acts as a negative regulator of inflammation by suppressing NF- $\kappa$ B pathway activation (Chen et al. 2015). They showed that under the condition of inflammation or ER stress MANF was upregulated and translocated into the nucleus where its C-terminal domain interfered with the binding of p65, a subunit of NF- $\kappa$ B, to the target gene promoters. Consequently, MANF suppressed the expression of NF- $\kappa$ B-dependent cytokines and inflammatory cell proliferation. In an *in vitro* model of ischemia, pretreatment of rat primary astrocytes with MANF alleviated oxygen-glucose deprivation induced cell damage, suppressed ER stress and decreased the secretion of pro-inflammatory cytokines IL-1 $\beta$ , IL-6, and TNF- $\alpha$  (Zhao et al. 2013). Anti-inflammatory effects of MANF were also reported in lipopolysaccharide (LPS) -exposed neural stem cells where MANF inhibited inflammatory signaling (IL-1 $\beta$ , TNF- $\alpha$  and IFN- $\gamma$ ) by regulating NF- $\kappa$ B and p38-MAPK pathways (Zhu et al. 2016).

CDFN was shown to be upregulated in rat primary microglia in response to LPS-induced inflammation (Zhao et al. 2014). Cultured microglial cells were exposed to LPS which induced distinct cytotoxicity and secretion of pro-inflammatory cytokines PGE<sub>2</sub> and IL-1 $\beta$ . Pretreatment with CDFN was able to attenuate these reactions, possibly due to suppressed activation of JNK pathway. In agreement with this, overexpression of CDFN in primary astrocytes was sufficient to alleviate ER stress induced cell damage and production of pro-inflammatory cytokines IL-1 $\beta$ , IL-6 and TNF- $\alpha$  (Cheng et al. 2013). The ability of CDFN to limit neuroinflammation has also been observed *in vivo*. In a rat model of spinal cord injury, the transplantation of mesenchymal stem cells overexpressing CDFN was able to decrease the production of PGE<sub>2</sub>, IL-1 $\beta$  and TNF- $\alpha$ , promote nerve regeneration and improve the overall motor function of the injured rats (Zhao et al. 2016). Importantly, in a 6-hydroxydopamine (6-OHDA) model of PD in rats, CDFN transfection into the SN reduced glial activation and IL-6 levels in the midbrain (Nadella et al. 2014). Finally, it should be noted that CDFN and MANF may also exert their actions through different mechanisms leading to

synergistic effects as proposed by Cordero-Llana and colleagues based on their observation that combined overexpression of both CDNF and MANF in the SNpc led to more efficacious dopaminergic neuroprotection in 6-OHDA-lesioned rats as compared to the overexpression of either one of the NTFs alone (Cordero-Llana et al. 2015).

### 2.1.5 Retrograde signaling of neurotrophic factors

In order to regulate neuronal development, plasticity and survival, target-derived NTFs have to be transported retrogradely from the synapses to the cell bodies where they can modulate gene transcription. Because the morphology of neurons is highly polarized, NTF-receptor complexes internalized at the nerve terminals may need to travel long distances along the axons to reach downstream signaling effectors localized in the cell body.

An early study from 1974 showed distal uptake and axonal transport of exogenous radiolabeled NGF and suggested that the retrograde transport of NTFs to neuronal cell bodies plays an important role in their survival promoting effects (Hendry et al. 1974). Later, endogenous unlabeled NGF was also confirmed to be retrogradely transported *in vivo* (Palmatier et al. 1984; Mufson et al. 1999). Upon NGF binding to TrkA receptor, the activated receptor-ligand complex is internalized by clathrin-dependent endocytosis (Ehlers et al. 1995; Grimes et al. 1996; Riccio et al. 1997; Howe et al. 2001; Ye et al. 2003). The endocytic vesicle containing the activated components of MAPK, PLC $\gamma$  and PI3K/Akt pathways is then retrogradely transported along the axon to the soma by a microtubule-associated protein complex called dynein (Vallee et al. 1989; Bhattacharyya et al. 2002; Vallee et al. 2004). This phenomenon is collectively called “signaling endosome” hypothesis.

Like neurotrophins, GFLs can also provide long-distance trophic support to neurons. For example, in the nigrostriatal pathway GFLs are taken up by axon terminals of dopamine neurons and transported to the cell bodies: receptor-mediated uptake and retrograde transport of GDNF have been demonstrated by injecting biologically active  $^{125}\text{I}$ -labeled GDNF into the adult rat striatum and using autoradiography to detect accumulated radioactivity in the ipsilateral SNpc 24h later (Tomac et al. 1995b; Voutilainen et al. 2009, 2011). This suggests that in the adult nigrostriatal system GDNF acts as an endogenous target-derived NTF for dopamine neurons. Similarly, GDNF was shown to be internalized and retrogradely transported by spinal motor neurons in neonatal rats (Yan et al. 1995). In compartmentalized cultures of sympathetic neurons, addition of GDNF to distal axons induced neuronal survival and neurite outgrowth in the cell bodies (Coulpier and Ibáñez 2004). The trophic effects were associated with retrograde transport of GDNF and GFR $\alpha$ 1, as well as activation of RET and Akt in the cell bodies suggesting a saturable, receptor-mediated mechanism for the retrograde signaling. A competition study investigating the retrograde transport of  $^{125}\text{I}$ -labelled GFLs demonstrated that the transport is mediated selectively by corresponding GFR $\alpha$  co-receptors and only limited cross-binding occurs (Leitner et al. 1999). Proteasomal degradation of RET in distal axons has been suggested to regulate the trophic signaling by modulating the amount of active GFL-GFR $\alpha$ -RET complexes that can be transported to cell bodies (Tsui and Pierchala 2010).

The molecular mechanisms of the retrograde GFL-GFR $\alpha$ -RET signaling are more poorly understood than the mechanism of retrograde NGF-TrkA signaling (Ito and Enomoto 2016). In response to receptor activation, GFL-GFR $\alpha$ -RET complex can be internalized from the plasma membrane via clathrin-mediated endocytosis (Richardson et al. 2006; Richardson et al. 2012; Crupi et al. 2015). The resulting early endosome contains phosphorylated RET and signaling effectors that can activate downstream trophic pathways. It might also anchor to the microtubule network and be transported to the soma by dynein motors in accordance with the “signaling endosome” hypothesis.

Intrastrially injected CDNF was shown to be internalized by different cortical and striatal neurons, possibly via the endocytic pathway, and undergo retrograde transport to the SNpc along nigrostriatal dopamine neurons (Voutilainen et al. 2011; Mätlik et al. 2017). <sup>125</sup>I-labeled CDNF also seemed to spread to the frontal cortex and hippocampus but active transportation to these areas could not be confirmed because the spread was not markedly affected by an excess of unlabeled CDNF (Voutilainen et al. 2011). Intrastrially injected <sup>125</sup>I-MANF, instead, was retrogradely transported, possibly via glutamatergic projection neurons, to the frontal cortex but not to the SN (Voutilainen et al. 2009).

In addition to retrograde transport, there is some evidence of anterograde transport of NTF signaling. In retinal ganglion cells, neurotrophins are anterogradely transported in vesicles along axonal microtubules to the midbrain superior colliculus using conventional kinesin-I motors (Butowt and von Bartheld 2007, 2001). However, the detailed molecular mechanisms and physiological significance of anterograde transport of NTFs remain largely elusive. Interestingly, in many neurodegenerative diseases there are alterations in axonal transport suggesting that defective retro- or anterograde NTF signaling may contribute to neurodegenerative processes (Bronfman et al. 2007; Ito and Enomoto 2016).


### **2.1.6 Neurotrophic factor mimetics, inducers and variants**

Peripherally administered NTFs do not cross the BBB and have short half-lives. An alternative strategy to activate NTF signaling is to use small molecules that bind to and activate NTF receptors or enhance NTF expression. The elucidation of the 3D structures of NTFs and their receptors makes it possible to use rational drug discovery methods in the design and development of such compounds. NTF mimetics potentially afford an opportunity to attain better drug-like properties as compared to recombinant NTFs: Ability to penetrate the BBB would make them suitable for peripheral administration, thus overcoming the high risks associated with the intracranial delivery of NTF proteins or viral vectors. Better diffusion in the brain parenchyma would ensure sufficient dose in the target area, selective binding profile and possibility for intermittent delivery would contribute to a better safety profile. In addition, small molecules would evade the potential risk of immunogenicity related to recombinant protein therapies and would be easier to manufacture in large quantities.



### 2.1.6.1 GFL mimetics

Considering the potent neurotrophic effects GFLs on midbrain dopamine neurons, targeting GFL-GFR $\alpha$ -RET signaling with GFL mimetics is of therapeutic interest in relation to PD (Lin et al. 1993). The proposed mechanisms of action for a GFL mimetic activating GFR $\alpha$ -RET receptor complex are shown in Figure 2.5. Tokugawa and colleagues described the first small molecule compound, XIB4035, mimicking the biological functions of GDNF (Tokugawa et al. 2003). XIB4035 binds to GFR $\alpha$ 1 and activates the GFR $\alpha$ 1-RET complex inducing RET phosphorylation and neurite outgrowth in a mouse neuroblastoma cell line. Later, this GDNF mimetic was shown to improve sensory function in animals with advanced neuropathy and to act as a positive modulator of GFR $\alpha$ -mediated signaling instead of being a direct GFR $\alpha$ 1 agonist (Hedstrom et al. 2014).



**Figure 2.5. Putative mechanisms of action of GDNF family ligand (GFL) mimetics.** There are three proposed mechanisms for the activation of GFR $\alpha$ -RET receptor complex by an agonist (i.e. endogenous GFL dimer or GFL mimetic). **(A)** The agonist first binds to one GDNF family receptor  $\alpha$  (GFR $\alpha$ ) co-receptor, which then engages a second GFR $\alpha$ . Subsequently, the agonist-(GFR $\alpha$ )<sub>2</sub> complex recruits two receptor tyrosine kinase RET molecules leading to autophosphorylation of intracellular tyrosine residues of RET, and activation of intracellular signaling pathways like PI3K/Akt and Ras/MAPK. **(B)** The agonist binds to a preformed GFR $\alpha$ -RET heterodimer, and thereby recruit another GFR $\alpha$ -RET pair and activate downstream signaling cascades. **(C)** The receptor complex exists as a preformed (GFR $\alpha$ )<sub>2</sub>-(RET)<sub>2</sub> heterotetramer which acquires a conformational change in response to the binding of the agonist, enabling the phosphorylation of the tyrosine residues of RET. PM - plasma membrane; MAPK - mitogen activated protein kinase; PI3K - phosphatidylinositol 3-kinase. Figure drawn by the author.

A novel family of small molecule GFL mimetics, termed BT13, BT18 and BT44, can selectively activate RET and its downstream signaling pathways (Bespalov et al. 2016; Mahato et al. 2020; Sidorova et al. 2017; Sidorova and Saarma 2020; Viisanen et al. 2020). These compounds have been shown to promote neurite outgrowth from cultured dorsal root ganglia sensory neurons and attenuate hyperalgesia in rat models of neuropathic pain. BT13 also supports the survival of cultured dopamine neurons, protects them from 6-OHDA and 1-methyl-4-phenylpyridinium (MPP<sup>+</sup>) induced cell death in a RET-dependent manner, and is able to stimulate dopamine release

in the mouse striatum (Mahato et al. 2020). The preferred binding sites of BT13 in the GFR $\alpha$ 1-RET complex were investigated using molecular docking simulations and identified to be the allosteric modulation site of GFR $\alpha$ 1 and the GFR $\alpha$ 1-interacting site of RET (Ivanova et al. 2018). The latter, more likely binding site, enables BT13 to act as a direct RET agonist.

#### 2.1.6.2 NTF inducers

A large number of compounds, both pharmacological and natural agents, has been suggested to stimulate endogenous NTF synthesis, which potentially has disease-modifying relevance in PD management. These compounds are extensively reviewed by Saavedra et al. (2008) and include, but are not limited to, dopamine agonists, monoamine oxidase (MAO) inhibitors, catecholamine reuptake inhibitors, anti-depressants, anti-psychotics, vitamin D<sub>3</sub>, estrogen and various medicinal herb extracts. Rasagiline, a clinically used MAO-B inhibitor, is one of such compounds that has attracted attention. It seems to upregulate BDNF and GDNF levels *in vitro* and *in vivo* (Bar-Am et al. 2005; Maruyama et al. 2004; Weinreb et al. 2009). In a rat 6-OHDA model of PD, rasagiline protected nigrostriatal dopamine neurons and alleviated motor decline (Blandini et al. 2004). Likewise, in a mouse lactacystin model of PD, rasagiline showed neuroprotective and neurorestorative effects and induced motor recovery (Zhu et al. 2008). Rasagiline has also been tested in early-stage PD patients in two phase II trials with delayed-start design (Parkinson Study Group 2004; Olanow et al. 2009; Hauser et al. 2009). It showed some disease-modifying effects, but further evidence is needed for decisive conclusions.

Dopaminergic neurotransmission itself is proposed to regulate the expression of GDNF and BDNF in the nigrostriatal system (Saavedra et al. 2008). For example, D1 and D2 receptor agonists increase the synthesis of GDNF in midbrain and striatal primary cultures (McNaught and Jenner 2000; Ohta et al. 2000; Guo et al. 2002; Ohta et al. 2004). Consequently, striatal dopamine depletion may lead to downregulation of GDNF which can contribute to the disease progression in PD patients. Clinically used D2-like dopamine receptor agonists pramipexole and ropinirole have been speculated to have neurotrophic properties. In rodent models of PD, they protected nigrostriatal dopamine neurons against 6-OHDA- and 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) -induced toxicity (Iida et al. 1999; Zou et al. 2000; Anderson et al. 2001; Ramirez et al. 2003). It should be noted, however, that the neuronal uptake of 6-OHDA and MPTP via DAT is a prerequisite for their toxic effects and that the neuroprotection seen in these models could be attributed to the regulation of DAT function by presynaptic D2 autoreceptor activation (Joyce et al. 2004). Neuroimaging evidence from clinical studies also suggests that pramipexole and ropinirole can affect the disease progression (Clarke and Guttman 2002; Parkinson Study Group 2002; Whone et al. 2003). The clinical relevance of these observations, however, has remained insignificant.

#### 2.1.6.3 NTF variants

Structurally modified or truncated NTF variants have been engineered in search of improved pharmacokinetic properties of NTFs (Piltonen et al. 2009; Runeberg-Roos et al. 2016). Insufficient distribution in the brain parenchyma has been proposed as one major factor contributing to the

lack of efficacy in the phase II clinical trials with GDNF and NRTN. As mentioned earlier, the diffusion of GDNF and NRTN in the brain tissue is hindered by their high affinity binding to heparan sulfates in the extracellular matrix. Thus, the promising neuroprotective effects of these GFLs in relatively small rat brain may not necessarily translate to much larger human brain where the GFLs have to diffuse long distances from the site of application.

To improve the diffusion properties of GFLs, biologically active variants with reduced heparin binding affinity have been investigated (Piltonen et al. 2009; Smith et al. 2015; Runeberg-Roos et al. 2016; Grondin et al. 2019). N-terminally truncated, non-heparin-binding  $\Delta 38\text{N}$ -GDNF was more widely distributed in the rat striatum than wild-type GDNF, stable in a brain extract stability assay and protective against amphetamine-induced turning behavior in a 6-OHDA model of PD (Piltonen et al. 2009). Nevertheless, the  $\Delta 38\text{N}$ -GDNF was unable to significantly promote the survival of dopaminergic cell bodies in the SNpc. Overall, widespread striatal diffusion of  $\Delta 38\text{N}$ -GDNF did not improve its efficacy as compared to wild-type GDNF. These data suggest that although heparan sulfate binding of GDNF restricts its distribution in the brain, it may be needed for the optimal neuroprotective effects.

Another GDNF variant (GDNFv) was engineered from wild-type GDNF by removing 31 amino acids from the N-terminal domain to reduce heparin binding and substituting two conservative amino acids (N38Q and D95E) to improve chemical stability (Smith et al. 2015). These modifications resulted in significantly improved distribution in the rat brain as well as in the larger rhesus macaque brain after intrastriatal administration, better chemical stability and lower predicted immunogenicity as compared to wild-type GDNF. *In vitro*, GDNFv was equivalent to wild-type GDNF in GFR $\alpha$ 1 receptor binding, RET phosphorylation and neurite outgrowth assays. In a 6-OHDA model of PD in rats, its ability to protect dopamine neurons was comparable to wild-type GDNF, and in naïve rhesus macaques GDNFv enhanced dopamine turnover (Grondin et al. 2019).

Structural modifications of NRTN were shown to increase its diffusion properties in the rat and monkey brain as compared to wild-type NRTN (Runeberg-Roos et al. 2016). This NRTN variant was also able to restore dopamine fibers in the striatum and alleviate motor impairment in 6-OHDA-lesioned rats more potently than GDNF showing promise for further development of NTF variants into disease-modifying therapies for PD.

## 2.2 Basal ganglia circuitry and dopaminergic system

Basal ganglia are comprised of a highly conserved chain of subcortical nuclei in the forebrain that orchestrate action selection, motor coordination, habit formation, learning and motivation (Gerfen and Surmeier 2011; Macpherson and Hikida 2019). They are in a central position to facilitate voluntary movements and to inhibit interfering or competing movements. Hence, the functions mediated by the basal ganglia cover behaviors that are vital for animal survival. The largest nucleus of the basal ganglia is the striatum. Based on the distinct functions and connections, the striatum is subdivided into the dorsal striatum consisting of the caudate nucleus and putamen, and the ventral striatum including the nucleus accumbens (NAcc) and olfactory tubercle. The other basal ganglia nuclei include the pallidal nuclei, that are comprised of the

external and internal segment of the globus pallidus (GPe and GPi, respectively) and the ventral pallidum, subthalamic nucleus (STN), substantia nigra pars compacta and pars reticulata (SNpc and SNr, respectively) and ventral tegmental area (VTA). Perturbations in basal ganglia functions underlie debilitating movement disorders that are characterized by impaired voluntary movements, the presence of involuntary movements or both, such as in PD and Huntington's disease (Macpherson and Hikida 2019). Dysfunctions of the basal ganglia also give rise to a spectrum of psychiatric conditions including schizophrenia, depression, anxiety, obsessive-compulsive disorder and addiction.


### **2.2.1 Organization of the basal ganglia circuits**

Basal ganglia circuitry receives its information from the cortex, modulates it under the control of midbrain dopaminergic innervation, and sends it back to the cortex through the thalamus (Gerfen and Surmeier 2011; Macpherson and Hikida 2019). The primary input nucleus of the basal ganglia is the striatum. It is anatomically connected to the cerebral cortex, thalamus and limbic system, and receives direct glutamatergic inputs from all main cortical areas (motor, sensory and associational areas) as well as thalamic nuclei. Midbrain dopamine neurons located in the SNpc and VTA send wide and dense dopaminergic innervation to the striatum where it merges with and modulates the cortical and thalamic glutamatergic input (Hooks et al. 2018; Klaus et al. 2019). The axonal arborization of a single dopamine neuron is shown to cover up to 6% of total striatal volume in rats (Matsuda et al. 2009). The basic organization of the basal ganglia circuitry is illustrated in Figure 2.6.

The striatum hosts two groups of GABAergic medium spiny neurons (MSNs; also called spiny projection neurons, SPNs) which give rise to two parallel projection pathways that convey information from the striatum to the output nuclei of the basal ganglia, i.e. the GPi and SNr (Gerfen and Surmeier 2011; Macpherson and Hikida 2019). The GPi and SNr are the final information-processing nuclei before output signals are sent to the ventral lateral nucleus and ventral anterior nucleus of the thalamus, superior colliculus and pedunculopontine nucleus (PPN) via GABAergic projections. The GPi and SNr are often considered as one somatotopic structure (the GPi/SNr), where the GPi is responsible for axial and limb movements and the SNr for head and eye movements.

The excitatory corticostriatal glutamatergic input arises from the cortex layer 5 pyramidal neurons and is directed to the dendritic spines of the MSNs. Similarly, thalamostriatal glutamatergic projections arise from the intralaminar thalamic nuclei and form synapses on the dendritic shafts and spine heads of the MSNs. These inputs are modulated by dopaminergic input from the nigrostriatal pathway which is directed to the spine necks of MSNs (Figure 2.8). Approximately half of the striatal MSNs express excitatory dopamine D1 receptors, substance P and dynorphin (D1-MSNs), and form the "direct" striatonigral pathway that projects monosynaptically to the GPi/SNr (Gerfen and Surmeier 2011; Macpherson and Hikida 2019). The other half expresses inhibitory dopamine D2 receptors and enkephalin (D2-MSNs) and project polysynaptically to the GPi/SNr forming the "indirect" striatopallidal pathway (Figure 2.6). D2-MSNs first extend axonal

projections to the GABAergic neurons of the GPe, which in turn provide inhibitory innervation to the STN. Thereafter, glutamatergic neurons of the STN send excitatory projections to the GPi/SNr.



**Figure 2.6. Simplified diagram of the basal ganglia-thalamocortical circuitry.** The striatum functions as an integrator of the basal ganglia input signals. It receives excitatory glutamatergic input from the cortex and thalamus (wide green arrows). In addition, dopaminergic input from the substantia nigra pars compacta (SNpc) arrives to the striatum (not shown) and modulates the glutamatergic input. The inhibitory GABAergic outputs of the basal ganglia project from the internal segment of the globus pallidus (GPi) and substantia nigra pars reticulata (SNr) to thalamus, superior colliculus and pedunculopontine nucleus (PPN) (wide red arrows). The direct pathway arises from GABAergic D1 receptor expressing medium spiny neurons (MSNs) and projects directly to the output nuclei GPi and SNr (thin red arrows). The indirect pathway originates from GABAergic D2 receptor expressing MSNs that project only to the external segment of the globus pallidus (GPe) (thin blue arrows). The GPe further sends GABAergic afferents to the subthalamic nucleus (STN), which connects the signal to the output nuclei via glutamatergic projections (thin green arrows). Also, hyperdirect pathway providing glutamatergic input from the cortex to the STN (wide green arrow), direct pathway axon collaterals from the striatum to the GPe (thin red arrow), and collaterals from the GPe to the output nuclei (thin blue arrow) are shown in the diagram. Reproduced with the permission of Annual Reviews, from Gerfen and Surmeier (2011), Annual Review of Neuroscience, © Annual Reviews 2011.

Evidence from optogenetical and electrophysiological measurements has shown, that direct pathway D1-MSNs also extend axon collaterals from the dorsal striatum and NAcc to the GPe and ventral pallidum, respectively, indicating that the canonical “direct/indirect” architecture of the striatal output pathways is more complex than originally thought (Gerfen and Surmeier 2011; Kupchik et al. 2015). Moreover, the GPe sends some GABAergic projections directly to the GPi/SNr, as well as back to the striatum, where they are able to inhibit striatal movement representations (see below chapter 2.2.4) (Bevan et al. 1998; Gerfen and Surmeier 2011; Mallet et al. 2016, 2012). The STN also receives glutamatergic input in addition to GABAergic input from the GPe. These glutamatergic projections originate from the cortex forming the “hyperdirect” pathway that bypasses the striatum (Nambu et al. 2002). The hyperdirect pathway has been proposed to be critical for suppressing thalamocortical output, and thereby inhibiting erroneous or competing motor programs (Nambu et al. 2002; Aron and Poldrack 2006). Dopamine neurons of the ventral tier of the SNpc also send dendrites through the SNr where they are able to release dopamine (Zhou et al. 2009). This dopaminergic connection between the SNpc and SNr is termed the “ultra-short pathway”. The dendritic dopamine release can activate presynaptic D1 receptors on the striatonigral MSN terminals which enhances GABA release in the SNr, and thus further suppresses the activity of nigral output projections to the thalamus (Robertson 1992).

### **2.2.2 Functions of the basal ganglia circuits**

The dorsal striatum is mainly responsible for cognition involving motor and executive functions, while the ventral striatum (especially the NAcc) primarily controls motivation, reward, positive reinforcement and Pavlovian-instrumental conditioning (Macpherson and Hikida 2019). The dorsal striatum is traditionally further divided into dorsomedial region (DMS) and dorsolateral region (DLS) based on anatomical and functional characteristics (McGeorge and Faull 1989). DMS receives input from the frontal- and parietal-associative cortices and plays an important role in skill acquisition and goal-oriented behavior (Gerfen and Surmeier 2011). DLS, on the other hand, receives the main input from the sensorimotor cortex, and this loop is responsible for automated action programs, habit formation and locomotion. Additionally, the basolateral amygdala sends excitatory afferents to the dorsal striatum which, according to experiments in non-human primates (NHPs), seem to modulate striatal activity based on the emotional context and internally generated goals (Kelley et al. 1982; Pan et al. 2010; Hernádi et al. 2015; Maeda et al. 2018). The NAcc receives projections from the limbic structures such as prefrontal and anterior cingulate cortices, amygdala and hippocampus. Recent developments in chemogenetic and optogenetic tools, viral tracers and electrophysiological techniques have enabled further division of the striatum into smaller functional subregions beyond the simplified ventral-dorsal and medial-lateral divisions (Chuhma et al. 2019).

Besides the anatomical and functional organization, the dorsal and ventral striatum can be neurochemically divided into two main compartments known as striosomes and matrix (Brimblecombe and Cragg 2017; Crittenden and Graybiel 2016; Graybiel and Ragsdale 1978). Striosomes occupy on average 10–15% of the striatal volume and form small labyrinth-like zones that are surrounded by larger matrix compartment. Striosomes are defined histochemically based on high expression of e.g.  $\mu$ -opioid receptors, substance P, D1 receptors and calretinin. Matrix, by contrast, is enriched with acetylcholinesterase, choline acetyltransferase, calbindin, somatostatin and D2 receptors. Both compartments contain direct and indirect pathway MSNs. The most prominent difference in the neuronal connectivity of the two compartments is the direct D1-MSN projection from striosomes to dopamine neurons of the ventral tier of the SNpc forming striato-nigro-striatal loop. Striosomes also receive more innervation from the ventral tier, and matrix from the dorsal tier of the SNpc. Interestingly, studies in rats suggest that dopamine levels and TH-immunoreactivity are higher in matrix but the density of dopaminergic axons is greater in striosomes. In addition to the dopaminergic innervation, striosomes receive abundant input from the limbic areas of the prefrontal cortex, whereas matrix receives cortical inputs primarily from the sensorimotor and associative cortices. The inputs from the thalamus and amygdala are also dissimilar between the two compartments. Studies of functional differences between striosomes and matrix are still at an early stage. Striosome-related circuits are proposed to regulate motivation, behavioral flexibility and choice under stressed conditions (Crittenden and Graybiel 2016). The two compartments are also suggested to be differentially involved in pathological processes in a range of psychomotor disturbances including PD, Huntington's disease, attention deficit and hyperactivity disorder (ADHD).

As described above, the striatum receives dense and convergent excitatory inputs from multiple cortical areas and thalamic nuclei. Thus, it serves as an integrator of this sensorimotor, cognitive and motivational information, selects the most appropriate behaviors and translates them into specific changes in the basal ganglia outputs. Cortico-basal ganglia neurocircuits can be divided into three functional loops, the sensorimotor, associative/cognitive and limbic loops, which play different roles in context-dependent action planning, selection and initiation (Gerfen and Surmeier 2011; Macpherson and Hikida 2019).

According to the classical model of action selection the basal ganglia tonically inhibit downstream motor centers (i.e. the motor cortex, superior colliculus and PPN) via GABAergic output projections, thus preventing the occurrence of unwanted or wrongly timed movements (Albin et al. 1989; Gerfen and Surmeier 2011; Klaus et al. 2019). Cortical and thalamic inputs give rise to the activation of the direct and indirect pathways which exert opposing effects on output nuclei. The activation of the direct pathway causes pauses in the tonic inhibition of the output projections enabling the desired movements, whereas the indirect pathway further suppresses the competing actions. Nigrostriatal dopamine release dichotomously modulates the direct and indirect projections facilitating the movement initiation.

Traditionally, rapid phasic activity of the midbrain dopamine neurons is suggested to drive reward, learning and transitions between movement states (Jin and Costa 2010; Matsumoto and Hikosaka 2009; Schultz 2007). Brief increase in the firing rate of the SNpc neurons and subsequent transient fluctuation in the striatal dopamine level occur immediately before or around self-paced movement initiation (Howe and Dombeck 2016; da Silva et al. 2018). It was recently shown that the transient increase in dopaminergic activity immediately preceding movement initiation correlates with the vigor of the initiated movement (da Silva et al. 2018). Brief optogenetic activation of the SNpc dopamine neurons before movement initiation, but not during an ongoing movement, increased the probability and vigor of future movements, while optogenetic inhibition resulted in less probable and less vigorous movements.

The classical model is challenged by accumulating evidence suggesting that both direct and indirect projection pathways are coactivated during movement and inhibited during immobility (Klaus et al. 2019). The new model proposes that both direct and indirect pathways might work in concert to facilitate the desired motor actions and suppress the unwanted movements by concurrently activating different basal ganglia output neurons. The cortex represents alternative motor plans to specific striatal MSN ensembles. Action specificity arises from the pattern of cortical inputs, while dopaminergic inputs nonspecifically modulate the excitability of the MSNs. Only with adequate phasic dopamine activity specific MSN ensembles become active depending on the strength of their cortical inputs. Too low dopaminergic input results in subthreshold striatal output despite the presence of strong cortical signals. In this model, the relative timing and activity levels of specific MSN ensembles are thought to have a critical impact on the final basal ganglia output signal. However, new cell type -specific recordings with single-cell resolution have demonstrated that the spatiotemporal dynamics of MSN subtypes during natural behavior is action-specific and rather complex (Klaus et al. 2017; Markowitz et al. 2018; Parker et al. 2018).



The abovementioned models of the basal ganglia functions are based on the current understanding of the circuitry and provide an oversimplified picture of their true physiology. The descriptions provided here, for example, completely ignore the fact that the GPe, GPi, STN and SNr neurons can also serve as autonomous pacemakers generating action potentials without synaptic input, consequently adding another dimension to the modulation of the basal ganglia circuitry (Gerfen and Surmeier 2011). There are still a great many questions to be answered before we can fully understand the role of the basal ganglia in action selection and motor control.

### **2.2.3 Striatal interneurons**

The distribution of the neuron populations is fairly uniform throughout the striatum: MSNs comprise 90-95% of the striatal neurons, and the remaining 5-10% are interneurons (Gerfen and Surmeier 2011; Chuhma et al. 2019). Despite the relatively small population, striatal interneurons have an important role in regulating the striatal output. Their behavioral relevance, however, remains poorly understood thus far.

Striatal interneurons can be divided into large cholinergic interneurons and three types of smaller GABAergic interneurons: (a) fast spiking (FS) PV-positive, (b) low-threshold spiking (LTS) somatostatin-, neuropeptide Y- and nitric oxide synthase-positive, and (c) calretinin-positive interneurons (Kawaguchi et al. 1995; Tepper et al. 2008; Kreitzer 2009). FS interneurons are most abundant in the rat striatum and their electrophysiological properties are dominated by short-duration action potentials, abrupt high-frequency firing and hyperpolarized resting potential. FS interneurons receive excitatory synapses from the cortex and thalamus and inhibitory inputs from other interneurons and a subpopulation of GPe neurons. In addition, dopamine and acetylcholine can increase their activity via D5 and nicotinic acetylcholine receptors (nAChRs), respectively. Gap junctions between the dendrites contribute to electrotonic coupling that can lead to synchronous firing among local FS interneuron populations.

LTS interneurons are physiologically distinguished from the FS cells by less negative resting potential, long-duration action potentials and higher input resistance (Kawaguchi et al. 1995; Kreitzer 2009). Like FS interneurons, LTS interneurons are also innervated by glutamatergic afferents from the cortex and thalamus and form synapses with MSNs. Dopamine can modulate the activation of LTS neurons via D5 receptors, and acetylcholine can exert either activating or inhibiting modulation via muscarinic M1 or M2 receptors, respectively. Due to the wide variety of transmitters, LTS interneurons can affect the activity of MSNs not only through the release of GABA but also nitric oxide, somatostatin and neuropeptide Y.

Cholinergic interneurons constitute only approximately 1–2% of striatal cells, but their effects are much more significant (Kreitzer 2009). They show tonic spontaneous firing on average at 2-10 Hz, long-lasting afterhyperpolarizations, depolarized resting potential and relatively high input resistance. Cholinergic interneurons receive excitatory innervation primarily from the intralaminar thalamic nuclei and, to a lesser extent, from the cortex as well as inhibitory synapses from MSNs. Their widespread axonal arborization connects mainly with D1- and D2-MSNs, but also with FS interneurons, through perisomatic synapses forming the feedforward thalamostriatal circuit



(Gerfen and Surmeier 2011). Nigrostriatal dopamine release modulates the activity of cholinergic interneurons through D2 and D5 receptors. D2, together with M2 and M4 receptor signaling, reduces their excitability (Calabresi et al. 1998; Chuhma et al. 2014; Maurice et al. 2004), whereas D5 receptor activation stimulates them (Aosaki et al. 1998). M2 also functions as a presynaptic autoreceptor regulating acetylcholine release (Hersch et al. 1994). Cholinergic interneurons can modulate direct-pathway MSNs through their M1 and M4 receptors and indirect-pathway MSNs through M1 receptors (Bernard et al. 1992; Yan and Surmeier 1996), as well as dopaminergic transmission through nAChRs on dopaminergic axon terminals (Wonnacott et al. 2000; Jones et al. 2001; Zhou et al. 2001; Cachope et al. 2012; Threlfell et al. 2012; Mamaligas et al. 2016). Thus, dopamine and acetylcholine together exert complex modulation of neuronal activity in the striatum that strongly influences basal ganglia output. On top of that, other neuromodulators such as adenosine, endocannabinoids, nitric oxide and various neuropeptides play important roles in shaping the final output signals.


#### **2.2.4 Striatal microcircuits regulating the direct and indirect pathways**

There are three distinct inhibitory microcircuits in the striatum that counterbalance the glutamatergic excitation of the MSNs (Gerfen and Surmeier 2011). They are thought to contribute to action selection by suppressing unwanted activities. Firstly, the feedforward inhibitory circuit originates from corticostriatal inputs that enhance the activity of FS interneurons (Mallet et al. 2005). The interneurons then form perisomatic GABAergic synapses on both D1- and D2-MSNs and effectively reduce their spiking. Secondly, FS interneurons receive GABAergic innervation from the GPe neurons forming the inhibitory pallidostriatal feedback circuit (Gerfen and Surmeier 2011). On this feedback circuit, dopamine exerts dual excitatory actions by directly depolarizing the interneurons through D5 receptors and reducing their synaptic inhibition through presynaptic D2 receptors on pallidostriatal nerve terminals (Bracci et al. 2002; Centonze et al. 2002, 2003; Wiltchko et al. 2010). Thirdly, the striatal feedback circuit involves local axon collaterals of D1- and D2-MSNs that make synaptic connections with the dendrites of other MSNs providing lateral inhibition (Gerfen and Surmeier 2011). The release of GABA on these collateral synapses is regulated by dopamine; D2 receptor activation decreases and D1 activation increases the release of GABA. Taken together, we can appreciate the complex role of striatal interneurons in regulating the activity of direct- and indirect-pathway MSNs. Their exact effects on the basal ganglia circuitry are yet to be resolved.

#### **2.2.5 Dopamine cell groups and projection pathways**

Arvid Carlsson, Eric Kandel and Paul Greengard were awarded the Nobel Prize in medicine and physiology in 2000 for their seminal work in the field of catecholaminergic neurotransmission in the late 1950s that lead to recognition of dopamine as an independent neurotransmitter in the brain (Carlsson 2001, 1959; Carlsson et al. 1958; Sano et al. 1959). Soon after that, in 1964, Dahlström and Fuxe mapped for the first time the locations of monoamine-containing cell bodies and nerve terminals in the rat brain (Dahlström and Fuxe 1964). Later, the development of sensitive immunohistochemical and tract-tracing techniques has provided a more detailed picture of the dopamine system making it one of the most completely mapped neurotransmitter systems

in the brain. The current understanding of the nine different dopamine-containing neuronal groups (groups A8–A16) in the mammalian brain is reviewed in (Björklund and Dunnett 2007) and depicted in Figure 2.7.



**Figure 2.7. Dopamine neuronal groups and dopaminergic pathways in the rat brain.** Dopamine neuron cell bodies are localized in nine distinctive groups (A8–A16), distributed from the midbrain to olfactory bulb (OB). The nigrostriatal pathway connecting dopamine neurons located in the substantia nigra pars compacta (A9) to the dorsal striatum is highlighted with wide red arrows and a circle. The mesolimbic and mesocortical pathways from the ventral midbrain dopamine neuron groups are illustrated by wide green arrows. The tuberoinfundibular pathway from the A12 and A14 groups and the diencephalospinal pathway from the A11 group are shown by thin green arrows. The cell groups are defined based on tyrosine hydroxylase immunoreactivity. HPC – Hippocampus. Adapted with permission from Elsevier: *TRENDS in Neurosciences*, Björklund and Dunnett (2007), © Elsevier 2007.

Ventral midbrain dopamine neurons are arranged in three different groups: the retrorubral field (A8 group), the SNpc (A9 group) and the VTA (A10 group) (Björklund and Dunnett 2007). These neuron populations project to defined areas of the forebrain and modulate specific functions according to their target areas. The dopaminergic projections are traditionally divided into three major pathways: Dopamine neurons located in the SNpc mainly innervate the dorsal striatum, forming the nigrostriatal pathway (Fallon and Moore 1978; Björklund and Dunnett 2007; Howe and Dombeck 2016). The nigrostriatal pathway is predominantly responsible for the regulation of motor functions and is implicated in the pathophysiology of PD (see chapter 2.3.2). Dopamine neurons located in the VTA primarily project to the ventral striatum (NAcc and olfactory tubercle) and cortical areas (prefrontal cortex and cingulate cortex) forming the mesolimbic and mesocortical pathways, respectively. These pathways regulate emotional behavior, motivation, reward, learning and cognitive functions, and are afflicted in an array of psychiatric disorders (Hornykiewicz 1978; Parker et al. 2016; Macpherson and Hikida 2019). The retrorubral field is a dorso-caudal extension of the SNpc and projects to both striatal, limbic and cortical areas (Björklund and Dunnett 2007).

The reality of the anatomical organization and projection patterns of the dopamine neurons in the midbrain cell groups is more complex than the original division described above (Loughlin and Fallon 1984; Swanson 1982). The cell groups can be further divided into different subcompartments (such as the dorsal and ventral tier of the SNpc and VTA) that differ in their cellular morphology, some molecular markers and projection patterns. For example, the dorsal

tier of the SNpc also contains neurons that send afferents to limbic and cortical areas (François et al. 1999). Using sophisticated tissue clarification, optogenetic and fiber photometry tools Lerner and colleagues identified two parallel dopamine neuron subpopulations within the SNpc with divergent biophysical properties and distinct output projections to the DMS and DLS (Lerner et al. 2015). The intermixing of the cell bodies of the nigrostriatal, mesolimbic and mesocortical pathways is more prominent in NHPs than in rodents (Williams and Goldman-Rakic 1998; François et al. 1999). Midbrain dopamine neurons also send sparse collateral innervation to several other basal ganglia structures including the GPe, GPi and STN enabling dopaminergic modulation of the basal ganglia circuitry at pallidal and subthalamic levels (Lindvall and Björklund 1979; Lavoie et al. 1989; Smith et al. 1989; Hassani et al. 1997; Gauthier et al. 1999; Prensa and Parent 2001).

The total number of TH-ir cells in the ventral midbrain (in A8, A9 and A10 groups together, bilaterally) is approximately 20 000–30 000 in mice and 40 000–45 000 in rats (Björklund and Dunnett 2007). About half of those cells are located in the SNpc. In humans, the same numbers are approximately 400 000–600 000, with >70% of the neurons located in the SNpc. The larger number of the midbrain dopamine neurons in humans is accompanied, for example, with much more extensive mesocortical innervation as compared to rodents.

Dopamine neurons in the diencephalon are divided into four groups: A14 (the rostral hypothalamic periventricular group), A13 (the dorsal hypothalamic group of the medial zona incerta), A12 (the infundibular group of the arcuate nucleus), and A11 (the posterior and dorsal hypothalamic group and the periventricular grey matter of the caudal thalamus) (Björklund and Nobin 1973). The best known projection pathway arising from these cell groups is the tuberoinfundibular pathway which originates from the A12 and A14 dopamine neurons and innervates the median eminence and the neural lobe of the pituitary gland (Goldsmith et al. 1990; Grattan 2015). Dopamine released from the tuberoinfundibular projections suppresses the secretion of prolactin from anterior pituitary gland by binding to D2 receptors.

The A11 group gives rise to diencephalospinal dopamine system. The A11 neurons project to all levels of the spinal cord providing the only source of inhibitory dopaminergic innervation for the spinal cord (Barraud et al. 2010; Puopolo 2019; Qu et al. 2006; Skagerberg and Lindvall 1985). Descending fibers innervate both the dorsal and ventral horn of the spinal cord and have an important role in sensorimotor integration and pain control. Dysfunction of the A11 dopamine system is suggested to have implications in the pathophysiology of pain (Fleetwood-Walker et al. 1988; Puopolo 2019), migraine (Charbit et al. 2009; Charbit et al. 2010) and restless legs syndrome (Clemens et al. 2006; Lanza et al. 2017).

The functions of the A13 group dopamine neurons and their so called incertohypothalamic projections are less well established (Fougère et al. 2019). They have been suggested to provide local hypothalamic innervation and send projections to locomotor centers of the brainstem. They may be associated with visceral and sensorimotor activities. The A15 group in the rostral hypothalamus contain TH-ir neurons but not detectable levels of dopamine or noradrenaline (Björklund and Dunnett 2007). The A15 neurons give rise to local diencephalic innervation but

their functional role has remained rather obscure. There is some evidence of their role in hormonal regulation (Clarkson and Herbison 2011; Liu and Herbison 2013; Brown et al. 2015).

Finally, the A16 group located in the olfactory bulb hosts the most numerous population of dopamine neurons in the brain (Pignatelli and Belluzzi 2017). Dopamine neurons form the most external (glomerular) layer of the olfactory bulb and function as interneurons co-releasing dopamine and GABA from separate vesicle pools. They modulate the activity of the olfactory nerve terminals and play a key role in odor processing. Interestingly, the A16 cells are regenerated throughout the lifetime; the RMS originating from the SVZ provides a constant source of new dopamine neurons to be integrated to the neuronal network of the olfactory bulb.

## **2.2.6 Dopamine receptors and modulatory effects**

### **2.2.6.1 Five dopamine receptors**

Reflecting the dense dopaminergic innervation from the ventral midbrain, the striatum has a prominent expression of dopamine receptors. The dopamine receptors are metabotropic G protein-coupled receptors and can be divided into two classes on the basis of the G-protein to which they couple: D1-like receptors (D1 and D5) are stimulatory and couple to  $G_s$  and  $G_{olf}$  proteins, whereas D2-like receptors (D2, D3 and D4) are inhibitory and couple to  $G_i$  and  $G_o$  proteins (Neve et al. 2004).  $G_s$  and  $G_{olf}$  proteins stimulate adenylate cyclase enzyme which elevates intracellular levels of cyclic adenosine monophosphate (cAMP). cAMP further activates protein kinase A (PKA) which has a broad range of cellular targets, including receptors, enzymes, ion channels and transcription factors. D1 receptor may also activate phospholipase C (PLC) in a cAMP-independent manner by interacting with  $G_q$ . PLC catalyzes the hydrolysis of a membrane lipid phosphatidylinositol 4,5-bisphosphate ( $PIP_2$ ) to two secondary messengers, diacyl glycerol (DAG) and inositol 1,4,5-trisphosphate ( $IP_3$ ).

$G_i$  and  $G_o$  target many effectors such as ion channels, phospholipases and receptor tyrosine kinases through a membrane-delimited mechanism involving the liberation of  $G_{\beta\gamma}$  subunits (Neve et al. 2004). For example, D2 receptor stimulation in striatal MSNs activates PLC via  $G_{\beta\gamma}$  subunits leading to mobilization of intracellular  $Ca^{2+}$  storages. As a result, transmembrane  $Ca^{2+}$  currents through L-type  $Ca^{2+}$ -channels are suppressed and neuronal excitability reduced (Hernández-López et al. 2000).  $G_{i/o}$  is also negatively coupled to adenylate cyclase leading to decreased levels of cAMP and activate protein phosphatases that directly counter the effects of PKA (Neve et al. 2004). Thus, the activation of D1-like and D2-like receptors leads to opposing cellular responses.

All five dopamine receptors (D1-D5) are expressed in the striatum, but D1 and D2 are by far the most abundant (Gerfen and Surmeier 2011). Their expression in direct- and indirect-pathway MSNs is dichotomous: direct-pathway MSNs express excitatory D1 receptors and indirect-pathway MSNs inhibitory D2 receptors. D1 and D2 receptors have different binding affinities (Macpherson and Hikida 2019). The high-affinity D2 receptors are occupied at lower dopamine concentrations (~10 nM) and respond more readily to tonic basal dopamine levels (~10-30 nM) inhibiting D2-MSNs. The low-affinity D1 receptors, instead, are primarily activated by increased phasic

dopamine release (at  $\sim 1 \mu\text{M}$ ) transiently enhancing the responsiveness of D1-MSNs (Surmeier et al. 2007). Hence, phasic dopamine release has an opposite effect on the activity of MSNs. D2 receptor signaling decreases the responsiveness of indirect-pathway MSNs to uncoordinated cortical activity, whereas transient D1 receptor signaling increases the excitability of the direct-pathway MSNs contributing to properly coordinated action initiation.

D2 receptors expressed at presynaptic axon terminals function as autoreceptors regulating synaptic neurotransmission. D2 autoreceptor activation inhibits the opening of voltage-gated  $\text{Ca}^{2+}$  channels resulting in decreased axonal dopamine release (Cardozo and Bean 1995). Additionally, D2 autoreceptor activation decreases dopamine release by other mechanisms that include inhibiting dopamine synthesis, enhancing dopamine reuptake via DAT and regulating vesicular monoamine transporter (VMAT) expression (Schmitz et al. 2003).

#### 2.2.6.2 Dopaminergic modulation of synaptic plasticity

In addition to modulating the short-term activity of the corticostriatal circuits, dopamine has an important part to play in regulating the long-term weakening and strengthening of synaptic transmission, i.e. LTD and LTP, respectively (Gerfen and Surmeier 2011). These synaptic plasticity events are believed to be the underlying mechanisms leading to learning and habit formation. Transient elevation in striatal dopamine release decreases the excitability of D2-MSNs and promotes the induction of LTD at corticostriatal synapses. LTD is also regulated by cholinergic interneurons and postsynaptic release of endocannabinoids which activate presynaptic CB1 cannabinoid receptors causing a decrease in glutamate release (Gerdeman et al. 2002; Tozzi et al. 2011).


In contrast, at the glutamatergic synapses of D1-MSNs, activation of D1 receptors increases the excitability and promotes the induction of LTP (Gerfen and Surmeier 2011). LTP seems to depend on the coactivation of D1 and NMDA receptors, as well as BDNF-TrkB signaling (Kerr and Wickens 2001; Calabresi et al. 2007; Jia et al. 2010). D1 receptor activation reinforces the function of AMPA and NMDA receptors and enhances their surface trafficking which is likely to contribute to the induction of LTP (Snyder et al. 2000; Sun et al. 2005; Hallett et al. 2006). Hence, the plastic changes caused by transient elevation in striatal dopamine release promotes the ability of cortical input to activate direct-pathway MSNs (LTP) and reduces its ability to turn on indirect-pathway MSNs (LTD). Ultimately, both changes facilitate movement initiation.

### 2.2.7 Dopamine lifecycle

#### 2.2.7.1 Dopamine biosynthesis and its regulation

Although dopamine is an important neurotransmitter in the brain, almost half of dopamine in the body is synthesized in the gastrointestinal tract (Eisenhofer et al. 1997). Here, the focus is on dopamine lifecycle in the CNS which is summarized in Figure 2.8. The biosynthesis of dopamine takes place in the cytosol of catecholaminergic neurons in a two-step reaction (Meiser et al. 2013). In the first step, TH catalyzes the hydroxylation of L-tyrosine at the phenol ring to L-3,4-

dihydroxyphenylalanine (L-DOPA). For this, TH requires a cofactor tetrahydrobiopterin (BH<sub>4</sub>), O<sub>2</sub> and Fe<sup>2+</sup>. In the second step, L-DOPA is decarboxylated to dopamine by aromatic amino acid decarboxylase (AADC, also known as DOPA decarboxylase) in a reaction that uses pyridoxal phosphate (vitamin B6) as a cofactor. The biosynthesis of dopamine and other catecholamines (noradrenaline and adrenaline) is under strong regulatory control. The activity of AADC, for example, is under feedback regulation by the reaction product.



**Figure 2.8. Nigrostriatal dopaminergic axon terminal and dopamine lifecycle.** Dopamine (DA) synthesis takes place in the cytosol. Tyrosine hydroxylase (TH) hydroxylates L-tyrosine to L-3,4-dihydroxyphenylalanine (L-DOPA) using tetrahydrobiopterin (BH<sub>4</sub>) as a cofactor. L-DOPA is decarboxylated to DA by aromatic amino acid decarboxylase (AADC). DA is sequestered into synaptic vesicles via vesicular monoamine transporter 2 (VMAT2). Upon depolarization of the axon terminal, synaptic vesicles are fused into the presynaptic membrane in a Ca<sup>2+</sup>-dependent manner resulting in quantal release of DA into the extracellular space. Dopaminergic axons form synaptic contacts on the dendritic spines necks of medium spiny neurons (MSN). Released DA diffuses into perisynaptic sites and activates postsynaptic D1- and D2-like receptors located along the dendritic membrane of MSNs. Presynaptic D2 receptors function as autoreceptors providing feedback inhibition of DA synthesis and release. Dopamine transporter (DAT) is responsible for the neuronal high-affinity reuptake of DA and termination of the signaling, after which DA is recycled back into the synaptic vesicles via VMAT2. DAT is distributed outside of the synaptic membrane. Polyspecific organic cation transporter 3 (OCT3) mediates non-neuronal uptake of DA into surrounding glial cells. In the glia, DA undergoes degradation, where DA is first metabolized to 3,4-dihydroxyphenylacetaldehyde (DOPAL) via monoamine oxidase (MAO). DOPAL is rapidly further converted to 3,4-dihydroxyphenylacetic acid (DOPAC) via aldehyde dehydrogenase (ALDH). Catechol-O-methyl transferase (COMT) methylates DOPAC to the main metabolite homovanillic acid (HVA). Alternatively, DA is first metabolized to 3-methoxytyramine (3-MT) via COMT, which is then converted to HVA by MAO and ALDH. Figure drawn by the author.

TH is the rate-limiting enzyme in the biosynthesis of catecholamines (Kumer and Vrana 1996). The regulation of TH expression and activity is strong and versatile. TH mRNA levels are under transcriptional and RNA stability regulation. TH protein undergoes translational control and allosteric modulations (Daubner et al. 2011). Dopamine competes with the cofactor BH4 to bind to the catalytic site of TH (Kumer and Vrana 1996). Thus, high dopamine levels exert feedback inhibition on the enzyme activity. In addition, various kinases can phosphorylate four different serine residues in the N-terminal regulatory domain of TH which results in substantially increased enzyme activity (Dunkley et al. 2004). Reversely, phosphatases can dephosphorylate TH, and therefore function as TH deactivators. Interaction with other proteins, such as DJ-1, alpha-synuclein ( $\alpha$ -syn), VMAT2, AADC and BH4 synthesizing guanosine triphosphate cyclohydrolase (GTPCH), controls the stability and activity of TH (Daubner et al. 2011). TH can also undergo ubiquitylation and subsequent proteasomal degradation (Døskeland and Flatmark 2002).

TH-immunolabelling is commonly used to identify dopamine neurons in the ventral midbrain. However, due to the strong regulation of TH expression in response to aging, neuronal insults or altered functional demands, TH-immunoreactivity is not always a reliable marker for identifying viable dopamine neurons (Björklund and Dunnett 2007). Thus, in the absence of TH-immunoreactivity, additional markers such as neuromelanin, DAT, AADC or VMAT2 may be needed to identify cells as functional dopamine neurons.

Interestingly, Nurr1, a transcription factor belonging to the orphan nuclear receptor superfamily, is predominantly expressed in dopamine neurons and essential in the development and maintenance of the midbrain dopaminergic system (Jankovic et al. 2005). Nurr1 regulates the expression of TH, DAT, VMAT2, AADC, GDNF and RET, in other words, all the proteins that are crucial for the normal function of dopamine neurons. Studies in Nurr1 knockout mice have shown that Nurr1 deficiency results in impaired function and increased vulnerability of midbrain dopamine neurons. Nurr1 expression is also decreased in the midbrain samples of PD patients. Thus, it may have a role in the pathogenesis of PD.

Besides the classical TH-dependent pathway, the biosynthesis of dopamine can also follow a cytochrome P450-mediated pathway where decarboxylation of tyrosine to tyramine by AADC precedes the hydroxylation step, in which CYP2D6 converts tyramine to dopamine (Hiroi et al. 1998; Bromek et al. 2011). However, the contribution of this pathway to the total dopamine synthesis seems to be minimal.

After synthesis, dopamine is readily sequestered into synaptic vesicles by VMAT2 (Chaudhry et al. 2008). Vesicular dopamine transport depends on the electrochemical  $H^+$  gradient generated by the vacuolar  $H^+$ -ATPase. Inside the vesicles, oxidation-prone dopamine is stabilized by acidic pH which prevents the generation of toxic derivatives in the cytosol (Vergo et al. 2007). In noradrenergic neurons, dopamine is further hydroxylated to noradrenaline by dopamine- $\beta$ -hydroxylase inside the vesicles (Daubner et al. 2011). Amphetamine and similar psychostimulants deplete dopamine from the synaptic vesicles by collapsing the electrochemical  $H^+$  gradient that drives the VMAT2-mediated dopamine transport (Sulzer and Rayport 1990; Chaudhry et al. 2008). Amphetamines can also inhibit VMAT2 directly.

### 2.2.7.2 Dopamine release and reuptake

Nigrostriatal dopamine neurons are autonomous pacemakers. They typically exhibit two main activity patterns: Under basal conditions, most dopamine neurons show tonic activity with regularly spaced spikes at 2-5 Hz which results in tonic dopamine release in the striatum (Grace and Bunney 1984; Hyland et al. 2002). Rewarding events or other stimuli can trigger brief higher-frequency bursts of approximately two to six action potentials at 15-30 Hz which evoke transient dopamine overflow from the synaptic vesicles. Aversive events, on the other hand, transiently decrease nigrostriatal dopaminergic activity. These transient changes in striatal dopamine release regulate motivation and movement initiation (Klaus et al. 2019).

In response to action potentials arriving to an axon terminal, depolarization opens voltage-gated  $\text{Ca}^{2+}$  channels allowing rapid influx of  $\text{Ca}^{2+}$ . Inside the axon terminal,  $\text{Ca}^{2+}$  induces conformational changes in synaptic soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE)-complex proteins which eventually lead to exocytotic release of synaptic vesicles filled with dopamine (Sulzer et al. 2016). Using electrochemical cytometry, it has been estimated that the average number of dopamine molecules per vesicle is 33 000 in striatal axon terminals of healthy mice, i.e. the quantal size of dopamine release is 33 000 molecules (Omiatek et al. 2013). Under basal conditions, the extracellular dopamine concentration resulting from tonic neuronal activity is in the range of 10-30 nM (Gonon and Buda 1985; Venton et al. 2003). It is high enough to tonically activate the high-affinity pre- and postsynaptic D2 receptors.

Some of the midbrain dopamine neurons, that express the vesicular glutamate transporter 2 (VGLUT2), can co-release glutamate in addition to dopamine (Chuhma et al. 2014; Hnasko et al. 2010; Stuber et al. 2010; Tecuapetla et al. 2010). Dopamine neuron -evoked AMPA and NMDA receptor activation have been reported in MSNs and cholinergic interneurons of the NAcc. Recently, SNpc dopamine neuron inputs were shown to drive pauses in the firing of DMS and bursts in the firing DLS cholinergic interneurons (Cai and Ford 2018). The pauses were mediated by D2 receptors while the bursts were driven by glutamate co-release and activation of excitatory metabotropic glutamate receptors (mGluRs). Also, rapidly after dopaminergic activation, firing in both direct- and indirect MSNs is inhibited by co-release of GABA from a subset of dopaminergic axon terminals (Tritsch et al. 2012; 2014; Chuhma et al. 2014). The co-release of GABA seems to require its cellular uptake by membrane GABA transporters (mGAT1 and mGAT4) and loading into dopaminergic vesicles by VMAT2.

$\alpha$ -Syn is a 140-amino acid long presynaptic protein that interacts with negatively charged phospholipid membranes through its N-terminus, is ubiquitously expressed throughout the nervous system and is prone for aggregation in PD (Burré et al. 2013). Although the precise function of  $\alpha$ -syn is not known, it seems to be an essential negative regulator of synaptic neurotransmission and activity-dependent dopamine release (Abeliovich et al. 2000; Larsen et al. 2006; Nemani et al. 2010). A growing body of evidence suggests a role for  $\alpha$ -syn in the maintenance of synaptic vesicle pools and/or as an auxiliary chaperone required for normal SNARE-complex assembly (Burré et al. 2010; Cabin et al. 2002; Chandra et al. 2005; Murphy et al. 2000; Snead and Eliezer 2014). Transgenic mice overexpressing human wild-type  $\alpha$ -syn show



selective alteration in the distribution of dopaminergic synaptic vesicles and deficits in dopamine release from the nigrostriatal axon terminals (Janezic et al. 2013). This early-onset phenotype is followed by age-dependent loss of dopamine neurons, reduced neuronal firing rate and motor impairments.

Other neurotransmitters can also regulate axonal dopamine release. In the dorsal striatum, glutamate spillover from its synapses can reduce dopamine release by acting on presynaptic mGluR1 on dopaminergic axon terminals (Zhang and Sulzer 2003). GABA has been shown to decrease dopamine release via presynaptic GABA<sub>B</sub> receptors (Charara et al. 1999; Smolders et al. 1995). In the NAcc, kappa and delta opioid receptors seem to be able to decrease dopamine release via a presynaptic mechanism (Schlösser et al. 1995; Svingos et al. 1999; 2001).

Dopamine released into the extracellular space can interact with postsynaptic dopamine receptors and presynaptic autoreceptors (Figure 2.8). The postsynaptic receptors are mostly located at distant extrasynaptic sites, often with a higher density in the perisynaptic zone of the dendritic spines of striatal MSNs (Hersch et al. 1995; Yung et al. 1995; Caillé et al. 1996). Therefore, dopamine has to spillover from the release sites and diffuse rather long distances to reach its receptors (Pickel et al. 1996). This kind of “volume transmission” is characteristic for monoamine transmitters and can modulate not only neuronal activity in larger brain areas, but also neuron-glia interactions (Fuxe et al. 2015; Taber and Hurley 2014). Diffusion-based dilution is also the main factor eventually terminating the dopamine signaling in most brain areas (Cragg and Rice 2004; Rice and Cragg 2008).

Another important factor contributing to the termination of the dopamine signaling and maintaining dopamine homeostasis is its effective reuptake from the extracellular space primarily via DAT (Meiser et al. 2013). DAT is rarely expressed on the active zone of a synapse. Instead, the dopamine reuptake sites are distributed along the membrane of dopaminergic fibers where they effectively restrict the extracellular diffusion of larger dopamine transients (Nirenberg et al. 1996; Pickel et al. 1996; Hersch et al. 1997). After release, dopamine can either be recycled back into dopaminergic axon terminals by DAT, or it can be degraded after non-neuronal uptake into surrounding glial cells by organic cation transporter 3 (OCT3) (so called uptake 2). OCT3 is a low-affinity and high-capacity transporter for monoamines widely expressed in neurons, astrocytes, microglia and oligodendrocytes (Eisenhofer 2001; Gasser 2019). It contributes to the clearance of monoamines after synaptic release and restricts their spread. In the striatum, DAT-mediated neuronal uptake is primarily responsible for the clearance of extracellular dopamine. Substrate-induced and PKC-dependent trafficking of DAT away from the cell surface is the predominant mechanism to regulate DAT function (Gulley and Zahniser 2003). For example, amphetamine typically induces DAT internalization. In addition, presynaptic D2 receptors mediate the upregulation of DAT activity. Neuronal uptake of dopamine is followed by its sequestration back into the synaptic vesicles via VMAT2.

### 2.2.7.3 Dopamine degradation

Following the uptake 2 by glial cells, dopamine is readily degraded by MAO and catechol-*O*-methyl transferase (COMT) (Meiser et al. 2013). There are two separate genes coding two isoforms of MAO, MAO-A and MAO-B, which are both found in the CNS. MAO is localized in the outer membrane of mitochondria and expressed in neurons, astrocytes and microglia. MAO catalyzes oxidative deamination of dopamine to a reactive intermediate product 3,4-dihydroxyphenylacetaldehyde (DOPAL) generating hydrogen peroxide as a side product (Eisenhofer et al. 2004). DOPAL is further metabolized predominately by aldehyde dehydrogenase to form 3,4-dihydroxyphenylacetic acid (DOPAC), or by aldehyde reductase to form 3,4-dihydroxyphenylethanol (DOPET), both of which are more stable metabolites. In neurons, dopamine that is leaking from the synaptic vesicles into the cytosol, is degraded by MAO and aldehyde dehydrogenase (Meiser et al. 2013). Notably, there are species specific differences in the affinities of MAO-A and MAO-B: in human, dopamine is mostly oxidized by MAO-B, but in rats by MAO-A.

COMT has two splicing isoforms encoded by one gene, soluble S-COMT and membrane-bound MB-COMT. Their role is to transfer a methyl group from S-adenosylmethionine to one of the two catechol hydroxyl groups (Gulberg and Marsden 1975; Männistö and Kaakkola 1999). This 3-*O*-methylation of DOPAC is Mg<sup>2+</sup>-dependent and leads to the formation of homovanilic acid (HVA) which is the main degradation product of dopamine and ultimately secreted into urine. In an alternative degradation pathway, dopamine is first methylated to 3-methoxytyramine (3-MT) by COMT, which is then oxidized to HVA by MAO and aldehyde dehydrogenase (Meiser et al. 2013). There is no significant COMT activity in presynaptic axon terminals but some activity in postsynaptic dendrites. The main COMT-activity seems to reside in astrocytes and microglial cells (Myöhänen et al. 2010).

Dopamine and its metabolites can also undergo phase II conjugation reactions before excretion, although the conjugation reactions seem to play only a minor role in dopamine metabolism in the CNS (Uutela et al. 2009). *O*-Sulfatation of dopamine or its metabolites is catalyzed by phenol sulfotransferases (Buu et al. 1981; Swahn and Wiesel 1976), and *O*-glucuronidation by uridine diphosphoglucuronosyl-transferases (Wang et al. 1983; Uutela et al. 2009).

The metabolism of dopamine inflicts constant oxidative stress on dopaminergic neurons that might be one predisposing factor for the specific neuronal vulnerability in PD (Meiser et al. 2013). High dopamine content exposes to vesicular leakage of dopamine into the cytosol. The degradation of cytosolic dopamine generates reactive oxygen species (ROS). As described above, oxidative deamination by MAO generates hydrogen peroxide that is a major source of oxidative stress in dopaminergic neurons. Besides this, the electron-rich catechol moiety of dopamine is also prone to oxidation if exposed to the neutral pH of the cytosol. Dopamine and L-DOPA can also be enzymatically oxidized to form highly reactive ortho-quinones dopamine-*o*-quinone and dopaquinone, respectively (Sulzer and Zecca 1999). Both, quinones and ROS, are apt to react unspecifically with cellular components and perturb their functionality. Oxidative stress and

changes in cellular functions can trigger neuroinflammation which is a characteristic feature in PD and can contribute to the degeneration of dopamine neurons.

## 2.3 Parkinson's disease

The first coherent descriptions of a clinical syndrome resembling PD can be found from ancient Indian medical literature dating back to 300 BC (later known as *Kampavata*) (Ovallath and Deepa 2013). In Western medicine, a British physician James Parkinson provided the first sound picture of PD with all the characteristic motor symptoms in his famous monograph *An Essay on the Shaking Palsy* published in 1817 (Parkinson 2002). Almost 150 years later, striatal dopamine depletion was associated with PD (Ehringer and Hornykiewicz 1960), and the effectiveness of L-DOPA in alleviating PD symptoms was demonstrated and brought into clinical practice (Birkmayer and Hornykiewicz 1961; Cotzias et al. 1969).

Although the emerge of motor symptoms is typically diagnostic, PD is not solely a movement disorder but associated with numerous debilitating non-motor symptoms (Kalia and Lang 2015). Until today, the most urgent unmet medical need in PD is a disease-modifying therapy that could halt or slow the progression of the disease and restore the neuronal circuits that have already been degenerated by the time of the diagnosis. Accumulating preclinical evidence for the neurorestorative properties of several NTFs seems convincing (Paul and Sullivan 2018). However, to date, none of them has translated into clinical use, although several clinical trials have been conducted and some are currently underway. This calls for new approaches to harness the regenerative potential of neuroplasticity in the brain.

Apart from the lack of a cure and incomplete understanding of the pathogenesis, a particular clinical challenge of PD management is that there are no biomarkers which would allow definitive diagnosis at early stages of the disease. Better diagnostic tools would also facilitate better distinction between different subtypes of PD enabling more effective personalized therapies.

### 2.3.1 Epidemiology and risk factors

PD is the second most common neurodegenerative disease after AD, and the most common movement disorder (Hebert et al. 2013; Marras et al. 2018). It is estimated to affect more than 6 million people worldwide causing over 200 000 deaths per year (Dorsey et al. 2018). The incidence of PD sharply increases with age which is the biggest risk factor for the disease (Driver et al. 2009; Pringsheim et al. 2014). On average 1% of people aged over 60 years lives with the disease, and the prevalence increases to about 2-4% in individuals over the age of 80 (de Lau and Breteler 2006; Pringsheim et al. 2014). With the overall aging of the population, PD is expected to impose an increasing socio-economic burden on societies. Men are approximately 1.5 times more likely to be afflicted than women (de Lau and Breteler 2006). The risk of developing sporadic PD is multifactorial involving a complex interplay between normal process of aging, genetic susceptibility, epigenetic mechanisms and environmental risk factors (Pang et al. 2019). The fundamental cause of the disease remains unknown in the large majority of the cases.

### 2.3.1.1 Environmental risk factors

Several comprehensive meta-analyses have identified pesticide, herbicide and solvent exposure, traumatic head injury, consumption of dairy products, rural living, beta-blocker use, agricultural occupation and well-water drinking as environmental risk factors that significantly increase the risk of PD (Noyce et al. 2012; Pezzoli and Cereda 2013; Ascherio and Schwarzschild 2016; Yang et al. 2017b; Pang et al. 2019). Environmental factors found to have a significant negative association with PD are tobacco use (cigarette smoking or smokeless tobacco) and caffeine consumption. Importantly, physical activity has been reported to reduce the risk of PD, and this potential protective effect has been related to the ability of exercise to stimulate antioxidant defense, angiogenesis and NTF expression (Cohen et al. 2003; Zigmond and Smeyne 2014; Bellou et al. 2016).

### 2.3.1.2 Genetic susceptibility

Only 5-10% of PD patients suffer from a familial form of the disease (Kalinderi et al. 2016). To date, more than 20 causal genes and 90 independent genetic risk variants for PD have been identified (Blauwendraat et al. 2020). The gene encoding  $\alpha$ -syn, *SNCA*, was the first one found to cause monogenic PD (Polymeropoulos et al. 1997). *SNCA* missense mutations and multiplications of the gene locus cause autosomal dominant form of the disease by increasing the tendency of  $\alpha$ -syn to misfold and form aggregates (Deng et al. 2018). The most frequent cause for late-onset and autosomal dominant PD are missense gain-of-function mutations in *LRRK2*-gene (Paisán-Ruiz et al. 2004; Zimprich et al. 2004; West et al. 2005; Deng et al. 2018; Pang et al. 2019). *LRRK2* mutations and elevated kinase activity can also be found in some sporadic cases (Di Maio et al. 2018; Simón-Sánchez et al. 2009). *LRRK2* encodes a large multidomain protein called leucine-rich repeat kinase 2 which is involved in multiple cellular processes, e.g. neurite outgrowth, synaptogenesis, vesicular trafficking, autophagy and protein synthesis (Dzambo and Halliday 2012; Martin et al. 2014).

Loss-of-function mutations in *PRKN*, *PINK1* and *DJ-1* are the most common examples of genes accounting for autosomal recessive forms of PD (Pang et al. 2019). Recessively inherited mutations in these genes are often associated with early-onset PD (Schrag and Schott 2006). Parkin (encoded by *PRKN*), *PINK1* and *DJ-1* are implicated in mitochondrial function, dynamics and quality control (McCoy and Cookson 2012). Mutations in *GBA*, which encodes a lysosomal enzyme  $\beta$ -glucocerebrosidase, form another major risk factor for early-onset PD and cause a more severe phenotype of PD with increased risk for dementia and reduced survival (Cilia et al. 2016; Lwin et al. 2004; Sidransky and Lopez 2012). Apart from the abovementioned typical examples, advances in human genetics and bioinformatics have identified various other PD-causing genes and risk loci as reviewed in (Deng et al. 2018) and (Blauwendraat et al. 2020).


## 2.3.2 Pathology of Parkinson's disease

### 2.3.2.1 Neurodegeneration

PD is a complex neurodegenerative disorder. The main pathological feature is progressive loss of dopamine neurons within the SNpc (Kalia and Lang 2015). The neuronal loss and the accompanying denervation of the nigrostriatal pathway leads to decreased dopamine release in the dorsal striatum (Figure 2.9.A and C). Striatal dopamine depletion disrupts the equilibrium between striatonigral and striatopallidal pathway MSNs resulting in increased activation of the GABAergic output nuclei of the basal ganglia (Mallet et al. 2006; McGregor and Nelson 2019). The outcome of these pathological changes is reduced activation of the motor cortex, and ultimately, the appearance of the cardinal parkinsonian motor symptoms (see below chapter 2.3.3).

When PD is diagnosed, a substantial proportion of dopamine neurons in the SNpc has already died. It has been estimated that at the onset of the motor symptom there is approximately 30% loss of dopaminergic cell bodies in the SNpc and 50-70% loss of striatal dopaminergic terminals in comparison to age-matched controls (Burke and O'Malley 2013). Thus, the molecular and cellular neuropathology is likely to start years or decades before the diagnosis. Over the course of the disease, the neurodegeneration continues progressing and results in 60-80% loss of nigral dopamine neurons and almost complete loss of striatal fibers at the time of death (Burke and O'Malley 2013; Kordower et al. 2013). The greater loss of striatal fibers as compared to the loss of neuronal bodies in the SNpc suggests that in PD the neurodegeneration occurs in a "dying-back" fashion with the distal axons deteriorating first and the cell bodies dying only months or years later. In addition, the substantial neurodegeneration observed at the time of the diagnosis suggests a prolonged preclinical stage. These two characteristics offer a clear therapeutic window for disease-modifying interventions.

Axonal degeneration of the nigrostriatal dopamine neurons is proposed to be an early pathological process and mediated by different mechanisms than those responsible for the degeneration of the cell bodies (Cheng et al. 2010; Tagliaferro and Burke 2016; O'Keefe and Sullivan 2018). Axonopathy is accompanied with disturbances in axonal transport which is another important feature in the early-stage PD pathology (Chu et al. 2012; De Vos et al. 2008). The lack of proper axonal transport can interfere with many neuronal functions crucial, for example, for synaptic communication. Accumulation of  $\alpha$ -syn inclusions in the axons in the early stages of the disease has been shown to hamper the retrograde transport of signaling endosomes from axon terminals to the cell bodies (Volpicelli-Daley et al. 2014). Importantly, this may have implications for neuroprotective actions of striatally secreted NTFs and reduce the effectiveness of interventional neurotrophic therapeutics administered into the striatum (Bartus et al. 2011).



**Figure 2.9. Neuropathology of Parkinson's disease (PD).** (A) Left panel: The cell bodies of dopamine neurons are located in the substantia nigra pars compacta (SNpc). In intact nigrostriatal pathway, neuromelanin in dopamine neurons produces the dark pigmentation of the SNpc as shown in the photograph of the brainstem transverse section (black arrows). Dopamine neurons project to the dorsal striatum (i.e. putamen and caudate nucleus) as illustrated with thick solid red lines. Right panel: Degeneration of the nigrostriatal pathway is the pathological hallmark of PD. The loss of neuromelanin-containing dopamine neurons causes depigmentation of the SNpc (black arrows). The loss of dopaminergic projections to the putamen (dashed red line; prominent loss) and the caudate nucleus (thin solid red line; less prominent loss) results in dopamine depletion in the dorsal striatum. Reproduced with permission from Elsevier: Neuron, Dauer and Przedborski (2003), © Cell Press 2003 (B) Intracellular Lewy pathology in SNpc specimens from a PD patient. Upper panel: Nerve cell body with three Lewy body inclusions that are double-stained for alpha-synuclein and ubiquitin. Lower panel: Lewy neurites in neuronal processes double-stained for alpha-synuclein and ubiquitin. Images adopted from Spillantini et al. (1998), © 1998 by The National Academy of Sciences (C) Classical (simplified) "Box and Arrow model" of cortico-basal ganglia-thalamocortical circuitry depicting the essential neuronal pathways in the healthy brain (left panel) and PD affected brain (right panel). In PD, the excitability of the direct and indirect pathways shifts in opposite directions following the dopamine depletion in the dorsal striatum; the excitability of D1 receptor expressing medium spiny neurons (D1-MSN) reduces whereas the excitability of D2-MSN increases. These changes contribute to the increased firing of the output nuclei internal globus pallidus (GPi) and substantia nigra pars reticulata (SNr). The resultant suppression of the motor thalamus leads to the decreased activation of the motor cortex. VL - ventral lateral nucleus of the thalamus; VA - ventral anterior nucleus of the thalamus; IL - intralaminar thalamic nuclei; STN - subthalamic nucleus; GPe - external globus pallidus. Figure drawn by the author, inspired by Albin et al. (1989).

The neuropathological processes are not confined exclusively to dopaminergic neurons but affect several non-dopaminergic neurotransmitter systems too. Histopathological, biochemical and imaging findings in PD patients have demonstrated the concomitant degeneration of serotonin, noradrenaline, acetylcholine and various neuropeptides containing neurons (Halliday et al. 1990; Gesi et al. 2000; Hirsch et al. 2003; Kish et al. 2008). Widespread neuronal loss can be found in several non-dopaminergic nuclei including the locus coeruleus, the nucleus basalis of Meynert, the dorsal motor nucleus of the vagus, the pedunculopontine nucleus, the raphe nuclei, amygdala and hypothalamus (Dickson 2012; Giguère et al. 2018). Dysfunctions in these systems are likely to contribute to the manifestation of non-motor symptoms of PD such as sleep disorders, depression and cognitive impairment (see chapter 2.3.3).

### 2.3.2.2 Pathological protein accumulation

The other pathological hallmark of PD is the presence of cytoplasmic fibrillar protein inclusions called Lewy bodies (within cell bodies) and Lewy neurites (mostly in dystrophic neuronal processes) (Goedert et al. 2013) (Figure 2.9.B). Lewy inclusions are composed of numerous cellular proteins, the most abundant being misfolded  $\alpha$ -syn, together with membranous material originating from vesicles and fragmented organelles (Shahmoradian et al. 2019; Spillantini et al. 1998, 1997; Wakabayashi et al. 2013; Xia et al. 2008). Misfolded  $\alpha$ -syn has the propensity to acquire a  $\beta$ -sheet-rich structure and form insoluble amyloid fibrils. Several post-translational modifications, including phosphorylation at serine 129, nitration and C-terminal truncation, have been proposed to be responsible for the conformational changes that contribute to the fibrillization of  $\alpha$ -syn (Barrett and Greenamyre 2015). A recent study by Mahul-Mellier et al. (2020) suggests that Lewy body formation involves a complex interplay between  $\alpha$ -syn fibrillization, post-translational modifications and interactions between  $\alpha$ -syn aggregates and membranous organelles. They argued that different stages of Lewy body formation, rather than simply  $\alpha$ -syn fibril formation, lead to molecular events that contribute to neuronal dysfunction and degeneration. Until today, however, the causal role of the Lewy pathology in the neuronal loss has not been confirmed (Kalia and Lang 2015). Clinical PD can also occur in the absence of Lewy pathology, particularly in certain monogenic forms of the disease such as most of the *PRKN* mutation -related cases and a proportion of patients with *LRRK2* mutations (Pouloupoulos et al. 2012; Doherty et al. 2013; Kalia et al. 2015).

As intermediates of the aggregation process  $\alpha$ -syn forms soluble oligomers and protofibrils that are proposed to be the most neurotoxic species potentially damaging cell membranes (Kalia et al. 2013; Karpinar et al. 2009; Winner et al. 2011). Some types of  $\alpha$ -syn oligomers may increase membrane permeability via a pore-forming mechanism which leads to  $\text{Ca}^{2+}$  influx and disruption cellular ion homeostasis (Angelova et al. 2016; Danzer et al. 2007). Notably, misfolded  $\alpha$ -syn loses its physiological functionality in nerve terminals which leads to synaptic dysfunction characterized by compromised neurotransmitter release and enlarged synaptic vesicles (Scott et al. 2010).

A number of cellular mechanisms are implicated in PD pathogenesis, including mitochondrial dysfunction, oxidative stress, ER stress, defective protein degradation systems, impaired

intracellular  $\text{Ca}^{2+}$  homeostasis and neuroinflammation, as described in the following chapters. However, their mechanistic details and interrelations have yet to be elucidated.

### 2.3.2.3 Braak staging system and Lewy pathology spreading

Lewy pathology spreads and afflicts multiple neuroanatomical areas in the brain. In 2003, Braak and colleagues introduced a staging system of Lewy pathology propagation based on semiquantitative assessment of Lewy body inclusions in *postmortem* autopsy samples of incidental and symptomatic sporadic PD cases (Braak et al. 2003). This hypothesis proposes that Lewy pathology spreads in an ascending direction throughout the brain in a sequence of six stages from medullary and olfactory nuclei towards the cortical areas as described in Figure 2.10.A. Generally, the staging system seems to apply to the majority of PD cases studied *postmortem* making it widely accepted (Dickson et al. 2010; Postuma et al. 2012). However, some studies have shown that not all cases fit to the hypothesis questioning its predictive validity and underpinning the heterogenous nature of PD (Kalaitzakis et al. 2008; Parkkinen et al. 2008; Jellinger 2009; Halliday and McCann 2010).

The Braak hypothesis was later revised to postulate that the initial event in sporadic PD is the access of  $\alpha$ -syn pathology to the brain through nasal and intestinal mucosal sites (Braak et al. 2003b; Braak et al. 2006; Hawkes et al. 2007; Hawkes et al. 2009). According to this “dual-hit hypothesis”  $\alpha$ -syn pathology enters the brain from the olfactory bulb via anterograde progression into the temporal lobe, as well as from the enteric plexuses via transsynaptic transmission and retrograde transport along the vagus nerve into the medulla and pons. Indeed, Lewy pathology is not restricted to the brain in PD;  $\alpha$ -syn immunoreactive inclusions have a widespread distribution throughout the spinal cord and PNS as well as multiple peripheral organs including skin, heart, submandibular gland and gastrointestinal system (Beach et al. 2010; Bloch et al. 2006; Wakabayashi and Takahashi 1997). This lends support to the hypothesis that the disease originates from the periphery. Interestingly, an epidemiological study showed that a full truncal vagotomy was associated with a decreased risk of subsequent PD supporting the role of the vagus as a gateway to the brain (Perez-Pardo et al. 2017; Svensson et al. 2015). The dual-hit hypothesis goes well together with hyposmia and constipation being prevalent prodromal symptoms in PD (see chapter 2.3.3).

The Braak hypothesis gained support from the observations that  $\alpha$ -syn aggregates can transfer between neurons. Findings of Lewy bodies in fetal dopamine neurons grafted into the striatum of PD patients showed that Lewy pathology can spread from host to grafted cells, and triggered the idea that the propagation of Lewy pathology is mediated by a prion-like transmission of  $\alpha$ -syn between neurons (Chu and Kordower 2010; Kordower et al. 2008; Li et al. 2008; Visanji et al. 2013). These findings have been confirmed in several *in vivo* experiments where intracerebral inoculation of synthetic recombinant  $\alpha$ -syn fibrils or Lewy body extracts from PD brains induced spreading of  $\alpha$ -syn pathology to anatomically interconnected brain areas and subsequent neurodegeneration (Luk et al. 2012; Recasens et al. 2014; Chu et al. 2019; Henderson et al. 2019). Indeed, mounting evidence has demonstrated the cell-to-cell transmission of small  $\alpha$ -syn aggregates, their ability to seed further aggregation via recruitment of endogenous host  $\alpha$ -syn in the recipient cells and their



bidirectional axonal transport (reviewed in Brundin et al. 2016; Killinger and Kordower 2019). The molecular mechanisms of  $\alpha$ -syn release to the extracellular space and uptake by neighboring cells are not fully elucidated and they may vary between different  $\alpha$ -syn species and conditions (Tyson et al. 2016). Data from cell culture models suggest that oligomeric and fibrillar  $\alpha$ -syn can be released via exocytosis in a process that is enhanced by cellular stress, or through uncontrolled leakage upon cell death. The uptake of  $\alpha$ -syn is believed to be mediated by receptor mediated endocytosis (Karpowicz et al. 2019).

Lewy pathology can also coexist with other proteinaceous inclusions, such as cytoplasmic tau-containing neurofibrillary tangles and extracellular  $\beta$ -amyloid (A $\beta$ ) plaques, suggesting interactions between pathogenic mechanisms of different proteinopathies (Jellinger 2012). Tau and A $\beta$  inclusions have been detected *postmortem* in the cortex of up to half of cognitively impaired or demented PD patients (Compta et al. 2011; Irwin et al. 2013). Thus, comorbidity with AD-type pathologies seems to be associated with the cognitive decline in PD.

#### 2.3.2.4 Dysfunctional protein degradation systems

Abnormal proteins are removed from cells via two main protein clearance systems: ubiquitin-proteasome system (UPS) and autophagy-lysosome pathway (ALP) (Rubinsztein 2006). Both UPS and ALP are responsible for the clearance of intracellular  $\alpha$ -syn (Rocha et al. 2018). Hence, dysfunction in either of these proteolytic machineries is a potential pathogenic mechanism in PD (Olanow and McNaught 2006; Pan et al. 2008; Ebrahimi-Fakhari et al. 2012; Xilouri et al. 2013; Lehtonen et al. 2019). Upon impaired degradation, misfolded  $\alpha$ -syn and other defective proteins accumulate in the cytoplasm contributing to their aggregation, perturbed cellular functions and cytotoxicity. Reciprocally, accumulating aberrant  $\alpha$ -syn species may further aggravate the dysfunction of UPS and ALP forming a vicious feedback loop that leads to neuronal death. The activity of both UPS and ALP declines with aging which may play a role in the neurodegenerative process (Xilouri et al. 2013).

The first evidence of the UPS abnormalities in PD was provided by *postmortem* studies of the SNpc where the proteasomal catalytic activity was found to be significantly reduced when compared to healthy brains (McNaught and Jenner 2001). Apart from the diminished activity, lower expression of different components involved in the normal function of UPS was also identified in the SNpc of PD brains (McNaught et al. 2002, 2003). Additional evidence is provided by genetic studies. Two genes linked to monogenic forms of PD are involved in UPS function: *PRKN* encodes parkin, an E3 ubiquitin ligase, which ubiquitinates substrate proteins and is needed for normal proteasome activation. PD-linked mutations in *PRKN* abolish this function which is thought to lead to the manifestation of the disease (Kitada et al. 1998; Um et al. 2010). A missense mutation in the gene encoding ubiquitin C-terminal hydrolase L1 (UCH-L1; a neuron-specific ubiquitin recycling enzyme) causes partial loss of the enzymatic activity and has been linked to increased risk of familial PD due to impaired de-ubiquitination and negative regulation of UPS (Leroy et al. 1998; Nishikawa et al. 2003).

Animal models of PD also provide support to the notion that impaired UPS is an important etiopathogenic factor in PD. In wild-type rats, selective proteasome inhibition with an intranigral lactacystin injection caused motor impairment, dose-dependent degeneration of nigrostriatal dopamine neurons and cytoplasmic accumulation of  $\alpha$ -syn immunoreactive protein inclusions (McNaught et al. 2002b). In transgenic mice, depletion of 26S proteasomes using a conditional knockout approach led to extensive neurodegeneration of the nigrostriatal pathway and forebrain regions and appearance of intraneuronal Lewy body -like inclusions (Bedford et al. 2008).

Similar to the findings in UPS system, various ALP-related components are differently expressed or dysfunctional in PD. For example, in *postmortem* examinations of the SNpc, chaperone-mediated autophagy-related proteins, such as lysosomal membrane receptors LAMP1 and LAMP2A and heat-shock proteins 70 and 73, were found to be decreased in PD brains as compared to age-matched controls (Alvarez-Erviti et al. 2010; Chu et al. 2009). Several genes linked to ALP have also been associated with PD (Gan-Or et al. 2015). For example, mutant LRRK2 interferes with ALP leading to retardation of  $\alpha$ -syn degradation and, as a consequence, its accumulation (Yue and Yang 2013). Additionally, loss-of-function mutations in the *GBA* gene, which encodes the lysosomal enzyme glucocerebrosidase, are strong genetic risk factors for PD suggesting a role for dysfunctional ALP in the pathogenesis of PD (Sidransky and Lopez 2012).

#### 2.3.2.5 Neuroinflammation

Neuroinflammation, with reactive astrocytosis and microgliosis, is strongly implicated in the pathogenesis of PD. Upon immune response, the morphology of microglia changes from a resting state ramified shape to an amoeboid profile (Kim and Joh 2006; Brück et al. 2016). This morphological change is accompanied with alterations in the expression of cell surface receptors and increased release of nitric oxide, ROS and cytokines. Some of the secreted factors are neuroprotective, such as GDNF and BDNF, but the majority are proinflammatory such as TNF $\alpha$  and IL-1 $\beta$ . Thus, reactive microglia amplify neuroinflammatory response and trigger oxidative damage and cytokine-receptor-mediated apoptosis in surrounding cells. This might contribute to the neurodegeneration in PD. On the other hand, reactive microglia can phagocyte cell debris and pathological protein aggregates, thus promoting neuroprotection.

Increased microglia and complement activation, T-cell infiltration and proinflammatory cytokine levels have been detected *postmortem* in the SNpc and striatum specimens from PD patients as compared to age-matched healthy subjects (McGeer et al. 1988; Hunot et al. 1999; Loeffler et al. 2006; Hirsch and Hunot 2009). Neuroimaging studies using positron emission tomography (PET) with a radiotracer for reactive microglia ( $^{11}\text{C}$ -PK11195) has demonstrated increased and stable microglial activation in the brainstem, basal ganglia and several cortical regions in patients with early-stage PD (Edison et al. 2013; Gerhard et al. 2006; Ouchi et al. 2005). Genetic studies have also identified a single-nucleotide polymorphism within the human leucocyte antigen (HLA) class II region that is associated with increased risk of developing PD, suggesting an immune system-related susceptibility to PD (Hamza et al. 2010; Nalls et al. 2014; Saiki et al. 2010).

Intriguingly, serum immune marker profile seems to be predictive of disease progression in PD patients. Elevated proinflammatory and lower anti-inflammatory marker levels were associated with faster progression of motor symptom and worse cognitive performance (Williams-Gray et al. 2016). In line with this, epidemiological studies have found a potential protective effect of regular non-steroidal anti-inflammatory drug use on the risk of PD (Gagne and Power 2010; Gao et al. 2011).

Although one can consider neuroinflammation to be a secondary phenomenon, accumulating evidence suggests that inflammatory processes *per se* have a contributing role in the pathogenesis of PD. For example, it has been demonstrated in 6-OHDA and MPTP mouse models of PD, that microglial-induced inflammatory events play a role in the neurotoxic effects on nigrostriatal dopamine neurons (He et al. 2001; Wu et al. 2002). In these studies, inhibition of microglia activation with minocycline mitigated dopaminergic cell death in the SNpc and reduced the level of IL-1 $\beta$ . Regardless of whether neuroinflammation has a causal role in neurodegeneration or it is a consequence of neuronal damage, it is clear that the engagement of immune system can exacerbate neuronal dysfunction in PD.

A strong body of evidence suggests that  $\alpha$ -syn can directly activate microglia in the CNS and initiate inflammatory response (Rocha et al. 2018). For example, overexpression of human wild-type  $\alpha$ -syn resulted in early microglial activation in the SNpc of transgenic mice (Su et al. 2008). In primary culture models, extracellular  $\alpha$ -syn triggered microglial activation and increased the expression of proinflammatory cytokines presumably in a receptor mediated manner (Zhang et al. 2005; Klegeris et al. 2008; Su et al. 2008).  $\alpha$ -Syn has also been shown to interact with astrocytes via Toll-like receptor 4 inducing nuclear translocation of NF- $\kappa$ B and subsequent astrocyte activation with increased expression of proinflammatory cytokines (Rannikko et al. 2015; Chavarría et al. 2018).

#### 2.3.2.6 Mitochondrial dysfunction, oxidative stress and ER stress

Mitochondrial dysfunction is closely related to sporadic and familial forms of PD (Moon and Paek 2015; Ryan et al. 2015). It can contribute to dopaminergic cell death due to cellular energy depletion, impaired calcium homeostasis, production of free radicals and increased oxidative stress. Early evidence arose from a group of young drug addicts in California who developed severe parkinsonian symptoms after self-administering a synthetic heroin (pethidine analogue) which was contaminated with a synthesis by-product MPTP (Langston et al. 1999, 1983). Later, *postmortem* examination of these individuals revealed that MPTP had selectively destroyed dopamine neurons in the SNpc. MPTP, when oxidized to MPP<sup>+</sup>, is taken up by dopamine neurons and leads to the inhibition of mitochondrial complex I which is a vital component in the electron transport chain (Nicklas et al. 1985). Other toxins and pesticides that interfere with mitochondrial complex I, like rotenone and paraquat, are also linked to the increased risk of PD (Tanner et al. 2011). Furthermore, in idiopathic PD cases, *postmortem* studies of the SNpc showed deficiency in mitochondrial complex I providing another direct indication of the mitochondrial dysfunction in PD (Schapira et al. 1989).


Importantly, several genes associated with monogenic forms of PD play a role in mitochondrial homeostasis. For example, PINK1 and parkin are the key regulators of mitophagy that is a process to dispose damaged mitochondria (Pickrell and Youle 2015; Ryan et al. 2015). The function of DJ-1 is less well characterized, but it seems to protect mitochondria from oxidative stress and be involved in maintaining mitochondrial homeostasis. Interestingly,  $\alpha$ -syn itself is known to interfere with mitochondrial bioenergetics. In the striatum and SNpc autopsy samples from PD subjects,  $\alpha$ -syn was shown to accumulate in the mitochondria of dopamine neurons where it interacted with the inner membrane, interfered with complex I activity and increased the production of ROS (Devi et al. 2008; Luth et al. 2014). Oligomeric and post-translationally modified species of  $\alpha$ -syn have also been shown to inhibit mitochondrial protein import leading to reduced mitochondrial respiration and enhanced production of ROS (Di Maio et al. 2016).

Accumulation of misfolded proteins can trigger ER stress which leads to UPR activation. ER stress and UPR are also common features of the PD pathogenesis and closely associated with aggregated  $\alpha$ -syn (Mercado et al. 2013). Neuropathological analysis of *postmortem* brain tissue derived from PD patients revealed increased immunoreactivity of the UPR markers phosphorylated PERK, IRE1 $\alpha$  and eukaryotic initiation factor 2 $\alpha$  (eIF2 $\alpha$ ) in dopamine neurons of the SNpc (Heman-Ackah et al. 2017; Hoozemans et al. 2007; Mercado et al. 2018). Phosphorylated PERK and IRE1 $\alpha$  also colocalized with  $\alpha$ -syn inclusions. In line, genetic and neurotoxin rodent models of PD as well as induced pluripotent stem cell -derived neurons overexpressing  $\alpha$ -syn have shown similar ER stress phenotype and activated UPR. Of note, ER-resident CDNF and MANF serve as potential therapeutic agents to mitigate ER stress in PD.

### 2.3.2.7 Selective vulnerability of the SNpc dopamine neurons

The particular susceptibility of the dopamine neurons in the SNpc to the neuronal damage is not fully understood. As described earlier, nigral dopamine neurons are autonomous pacemakers spiking at low frequencies (2-5 Hz) and exhibiting broad action potentials (>2 ms). Unlike most of the other pacemaking neurons that rely on Na<sup>+</sup> channels, the SNpc dopamine neurons have a strong reliance on L-type voltage-dependent Ca<sup>2+</sup> channels with a Ca<sub>v</sub>1.3 subunit leading to increased Ca<sup>2+</sup> entry into the cells (Surmeier and Schumacker 2013). In addition, the SNpc dopamine neurons have low intrinsic Ca<sup>2+</sup> -buffering capacity. In comparison, the dopamine neurons in the VTA, which are also slow pacemakers but less prone to degeneration in PD, have much lower L-type Ca<sub>v</sub>1.3 Ca<sup>2+</sup> channel density, do not manifest significant intracellular Ca<sup>2+</sup> oscillations and express high levels of the Ca<sup>2+</sup> -buffering protein calbindin. The maintenance of Ca<sup>2+</sup> gradient across the plasma membrane is energetically expensive. Because of the high energy demand, nigral dopamine neurons are strongly dependent on the mitochondrial ATP production. Increased mitochondrial oxidative phosphorylation gives rise to high levels of ROS generation and oxidative stress in the neurons. High metabolic demands also increase the risk of bioenergetic crisis in case of an episodic disruption in the mitochondrial function. Reduced Ca<sup>2+</sup> influx might explain the potentially reduced risk of PD associated with the use of calcium channel blockers (Noyce et al. 2012).

The SNpc dopamine neurons have a massively arborized axonal tree with a very high number of axon terminals which causes a great bioenergetic burden on these cells (Giguère et al. 2018). Elevated energetic requirements can deplete cellular antioxidant storages and subject neurons to chronic oxidative stress. It is estimated that up to half of the energy consumed by the SNpc dopamine neurons is used to action potential firing and neurotransmitter release. Thus, stressed neurons may try to adapt to excessive metabolic needs by dysregulating axon terminals as seen in the early phases of PD.



**Figure 2.10. (A) Braak staging system of Lewy pathology spreading.** In stage 1, Lewy inclusions start to appear in the dorsal motor nucleus of the vagus nerve in the medulla oblongata and olfactory system. In stage 2, locus coeruleus and caudal raphe nuclei in the pontine tegmentum become affected. In stage 3, the pathology ascends to the midbrain including substantia nigra, amygdala and the cholinergic nuclei of the basal forebrain. In stage 4, the pathology reaches thalamus, hippocampus and temporal mesocortex. In stages 5 and 6, several cortical areas, including prefrontal, primary sensory and motor cortices, and finally, the entire neocortex become involved. Growing severity of the Lewy pathology is shown by increasing shading (red, violet, black). Reproduced with permission from Springer Nature: Journal of Neural Transmission, Braak et al. (2003b), © Springer Nature 2003 **(B) Time course of Parkinson's disease progression.** Accumulating clinical symptoms (red line) are associated with the loss of nigral dopamine neurons (dashed black line) and Braak stages of Lewy pathology spreading (in violet). Up to two decades long pre-motor/prodromal phase (stages 1 and 2) characterized by specific, mainly autonomic, olfactory and sleep-related non-motor symptoms can precede the diagnosis (time 0 years). During stage 3, the characteristic motor features start to manifest, leading to the diagnosis between stages 3 and 4. Additional non-motor symptoms develop with the disease progression, causing significant disability. Stages 5 and 6 are associated with advanced disease with debilitating axial motor symptoms. Complications of dopaminergic therapy contribute to clinical disability. EDS - excessive daytime sleepiness; MCI - mild cognitive impairment; RBD - REM sleep behavior disorder. Adapted with permission from Elsevier: The Lancet, Kalia and Lang (2015), © Elsevier 2015.

Another hypothesis linking increased oxidative stress to the vulnerability of the SNpc neurons is that oxidation of cytosolic dopamine and its metabolites leads to the excessive production of cytotoxic free radicals, dopamine-quinones, which may disrupt the function of e.g. mitochondrial protein complexes or DAT (Greenamyre and Hastings 2004; Giguère et al. 2018). This may well be a contributing factor in the pathogenesis of PD but probably not the principal culprit. Dysfunctional iron homeostasis causing overload of free iron in the SNpc has also been proposed to trigger oxidative stress in PD. However, this hypothesis has remained controversial thus far (Sian-Hülsmann et al. 2011).

### **2.3.3 Symptoms and diagnosis of Parkinson's disease**

The characteristic motor manifestations of PD include bradykinesia, rigidity, resting tremor, postural instability and gait disturbances (Kalia and Lang 2015). Bradykinesia is the precondition for the clinical diagnosis of PD [UK Parkinson's Disease Society Brain Bank clinical diagnostic criteria (Hughes et al. 1992); revised by Movement Disorder Society Clinical Diagnostic Criteria for Parkinson's disease (Postuma et al. 2015)]. In addition, at least one other motor feature is required for the diagnosis: muscular rigidity, 4-6 Hz resting tremor, and/or postural instability. The diagnosis is confirmed by excluding possible alternative disorders and with the presence of at least three supportive criteria such as unilateral onset, progressive symptoms, marked responsiveness to L-DOPA treatment, L-DOPA-induced dyskinesias and clinical course of 10 years or more. The diagnosis is based on clinical neurological examination but neuroimaging techniques such as single-photon emission computed tomography (SPECT), magnetic resonance imaging (MRI) and PET can be used in differential diagnosis and tracking the progression of the disease (see below chapter 2.3.4) (Politis 2014). Neuropathological overlap and heterogeneity in clinical manifestations complicate the differential diagnosis between early-stage PD and atypical parkinsonian syndromes such as multiple system atrophy (MSA), progressive supranuclear palsy (PSP) and corticobasal degeneration (CBD). Despite advanced diagnostic tools, full certainty of the diagnosis cannot be achieved *pre mortem*; 75-95% of the diagnoses are confirmed by histological identification of Lewy pathology upon autopsy (Postuma et al. 2015).

Apart from the motor impairment, PD patients suffer from heterogeneous non-motor symptoms such as olfactory deficits, cognitive impairment, psychiatric symptoms, sleep disturbances, autonomic dysfunctions, pain and fatigue (Khoo et al. 2013; Kalia and Lang 2015) (Figure 2.10.B). Some of these symptoms are of dopaminergic origin while the others are non-dopaminergic, reflecting the multisystem nature of PD (Fox et al. 2008; Chaudhuri and Schapira 2009). Oftentimes, it is the non-motor symptoms, rather than the motor symptoms, that dominate the decline of patients' health-related quality of life (Barone et al. 2009; Martinez-Martin et al. 2011; Duncan et al. 2014). Cognitive decline is one of the most prevalent and debilitating non-motor feature of PD (Aarsland et al. 2009; O'Callaghan and Lewis 2017). Mild cognitive impairment (MCI) is considered as an intermediate state that typically progresses to PD dementia. In some patients, MCI is recognized already at the earliest stages of the disease. The likelihood of manifesting MCI or dementia considerably grows as the disease progresses. On the whole, the burden of non-motor symptoms increases over time. Psychiatric symptoms such as anxiety, depression and

psychosis, and autonomic dysfunctions including constipation, dysphagia, urinary incontinence and postural hypotension are common features in the later stages of PD and resistant to L-DOPA treatment. These late-stage symptoms are predictive for institutionalization and, eventually, mortality.

Certain non-motor symptoms can precede the onset of the motor impairment and clinical diagnosis by more than a decade (Postuma et al. 2012) (Figure 2.10.B). These prodromal, or pre-motor, symptoms typically include impaired olfaction (hyposmia), constipation, orthostatic hypotension, rapid eye movement (REM) sleep behavior disorder (RBD), depression and excessive daytime sleepiness. Efforts have been made to identify prodromal symptoms that could be used as diagnostic markers to screen for PD before the motor manifestations. Depression, constipation and idiopathic RBD are currently the strongest candidates that, together with family history of PD, could support the diagnosis but the specificity and predictive value of these markers are still insufficient (Howell and Schenck 2015; Noyce et al. 2012).

### **2.3.4 Biomarkers of Parkinson's disease**

Biomarkers can be used to assist in early and differential diagnosis and predict the course of a disease. They can also help to assess response to a therapeutic intervention. An optimal biomarker is easily accessible, inexpensive, sensitive and reproducible (Yilmaz et al. 2019). Currently, there are several biomarker candidates available for PD but none of them is specific or reliable enough to be used alone to diagnose or follow the progression of the disease (Kalia 2018; Parnetti et al. 2019). This shortage mainly arises from the heterogeneity of PD on genetical, pathophysiological and clinical level. The potential approaches to identify new biomarkers include biomolecular, histological and “omics” based analyses of biofluids and peripheral tissue samples (biochemical biomarkers), brain imaging modalities (imaging biomarkers), electrophysiological and digital biomarkers (other biomarker modalities). In general, the diagnostic utility of these approaches remains to be validated in large trials with standardized study protocols before their introduction to clinical practice. The use of a combination of biomarkers could detect multiple pathological aspects and result in improved diagnostic accuracy.

#### **2.3.4.1 Biochemical biomarkers**

Identifying biomarkers from body fluids that are easily accessible (e.g. blood, saliva or urine) would be desirable, but due to the peripheral nature of these matrixes, the sensitivity and specificity issues might limit their utility (Chahine and Stern 2017). Accumulating evidence suggests diagnostic value for cerebrospinal fluid (CSF) and blood biomarkers reflecting the pathophysiology of PD such as  $\alpha$ -syn species, lysosomal enzymes, neurofilaments and markers of A $\beta$  and tau pathology (Parnetti et al. 2019). Especially various forms of  $\alpha$ -syn, total monomeric, phosphorylated, oligomeric and aggregated species, have been investigated intensively (Atik et al. 2016; Visanji et al. 2017). The close contact of the CSF to the brain makes it a potential source for diagnostic markers reflecting the ongoing pathological processes and metabolic changes in the CNS (Constantinescu and Mondello 2013; Eusebi et al. 2017; Parnetti et al. 2019). It is not, however, an ideal matrix for longitudinal monitoring because of the need for repeated lumbar

punctures and risk for blood contamination which may disturb some analyses. Nevertheless, the CSF levels of  $\alpha$ -syn species have provided encouraging results as biomarker candidates for PD. The total concentration of  $\alpha$ -syn in the CSF seems to be lower, whereas the levels of  $\alpha$ -syn oligomers and phosphorylated  $\alpha$ -syn are elevated in PD patients as compared to healthy controls.

Efforts have also been made to quantify plasma levels of  $\alpha$ -syn species, neurofilaments and inflammatory cytokines as potential biomarkers for PD. Plasma sampling would provide a convenient route for early diagnosis and longitudinal disease monitoring. However, the measurement of total  $\alpha$ -syn from plasma has provided inconsistent outcomes, partly because plasma samples are easily contaminated with  $\alpha$ -syn containing erythrocytes (Constantinescu and Mondello 2013; Parnetti et al. 2019). Quantification of  $\alpha$ -syn oligomers and phosphorylated  $\alpha$ -syn from plasma provides an alternative approach which has shown more concordant results with increased levels in PD patients.

Untargeted “omics” techniques in combination with bioinformatic tools provide a powerful approach to detect large amounts of analytes in various biosamples and then compare these analytical “fingerprints” between patients with healthy controls. Metabolomics has been utilized to detect PD-related alterations for example in plasma and CSF levels of amino acids, fatty acids and sugars (LeWitt et al. 2013; Trupp et al. 2014; Willkommen et al. 2018). Also, attempts to use proteomics and transcriptomics data collected from the body fluid samples have been made (Scherzer et al. 2007; Halbgebauer et al. 2016; Santiago et al. 2018). These “omics” based biomarkers have the potential to unravel new aspects of PD pathogenesis and aid in the future diagnostics of the disease.

In consistence with the evidence suggesting that  $\alpha$ -syn pathology appears in the PNS before propagating to the brain, attention has been paid to histological examination of phosphorylated  $\alpha$ -syn deposits in various peripheral tissue biopsies from PD patients (Cersosimo and Benarroch 2012; Lee et al. 2017). For example, submandibular gland biopsy and subsequent detection of phosphorylated  $\alpha$ -syn aggregates in the autonomic nerve fibers have shown promise as an early histological biomarker (Adler et al. 2016; Vilas et al. 2016). Histological analysis of phosphorylated  $\alpha$ -syn in dermal nerve fibers in skin biopsies is also a sensitive candidate for early diagnosis (Doppler et al. 2016).

#### 2.3.4.2 Imaging biomarkers

Brain imaging techniques are widely studied tools that can provide support for early, differential and prognostic diagnosis. Multimodal imaging allows the visualization of structural and functional changes in the brain and can reveal pathophysiological mechanisms underlying the disease processes (Saeed et al. 2017). As imaging modalities are non-invasive, they are well suited for longitudinal tracking of disease progression and therapeutic responses.

Structural MRI is used to assess regional tissue atrophy and serves as a potential aid for PD diagnosis (De Marzi et al. 2016; Frosini et al. 2017; Pyatigorskaya et al. 2017). Neuromelanin-sensitive MRI imaging of the SNpc and locus coeruleus has shown promising diagnostic accuracy



in differentiating PD patients from healthy controls (Castellanos et al. 2015). Functional MRI (fMRI) can be used to detect changes in brain network activities by measuring cerebral oxygen-rich blood flow. Significantly reduced functional connectivity in resting-state fMRI have been reported within the basal ganglia of patients with early-stage PD (Rolinski et al. 2015; Szewczyk-Krolikowski et al. 2014). Indeed, resting-state fMRI of basal ganglia connectivity is a promising diagnostic tool that may help to identify patients at risk of developing PD (Rolinski et al. 2016).

SPECT is a useful technique to assess the integrity of the nigrostriatal dopamine pathway (Saeed et al. 2017). Gamma-emitting cocaine analogues, such as  $^{123}\text{I}$ -FP-CIT and  $^{123}\text{I}$ - $\beta$ -CIT, bind selectively to presynaptic DAT and can be used to measure the density of nigrostriatal fibers. Reduced striatal uptake in DAT-SPECT is a characteristic readout in PD patients when compared to healthy controls. A meta-analysis suggests that DAT-SPECT is useful in the diagnosis of early PD and in differentiating PD from essential tremor and vascular parkinsonism (Vlaar et al. 2007). DAT-SPECT can also be utilized to follow disease progression because striatal binding seems to correlate to disease severity stages (Benamer et al. 2000; Brücke et al. 1997).

PET is another imaging modality taking advantage of radioactive ligands. It can be utilized to gauge the integrity of dopaminergic system, cerebral glucose metabolism, pathological A $\beta$  and tau protein accumulation and neuroinflammation (Saeed et al. 2017). 6- $^{18}\text{F}$ -fluoro-L-dopa ( $^{18}\text{F}$ -DOPA) is a PET tracer that is converted to  $^{18}\text{F}$ -dopamine by AADC. Thus,  $^{18}\text{F}$ -DOPA measures the activity of AADC and provides an indirect estimation of the presynaptic dopamine storage pools. In PD, striatal  $^{18}\text{F}$ -DOPA uptake is decreased and associated with the progression of the disease. PET imaging of presynaptic DAT, using for example  $^{18}\text{F}$ -FP-CIT or  $^{11}\text{C}$ -methylphenidate, shows reduced striatal uptake in PD patients, in line with DAT-SPECT findings. However, compensatory mechanisms, such as upregulation of AADC activity and downregulation of DAT, in response to neurodegeneration may inflict biases in AADC-PET, DAT-PET or DAT-SPECT-based estimates of the nigrostriatal injury (Lee et al. 2000). VMAT2 imaging, instead, for example with  $^{11}\text{C}$ -dihydrotetrabenazine ( $^{11}\text{C}$ -DTBZ) PET, seems to be less prone to the compensatory changes, thus producing more reliable readouts of the nigrostriatal degeneration in PD. Despite considerable efforts to develop an  $\alpha$ -syn-specific radiotracer, a direct method for  $\alpha$ -syn imaging *in vivo* is still lacking (Eberling et al. 2013). Such a tracer would allow for tracking  $\alpha$ -syn pathology spreading over time and monitoring the efficacy of  $\alpha$ -syn targeting therapies.

#### 2.3.4.3 Other biomarker modalities

Electrophysiological techniques, such as electroencephalography (EEG), can be used to identify candidate biomarkers for PD. For example, lower background rhythm frequency and increased relative power of delta and theta oscillations in resting-state EEG show promise as predictive biomarkers for cognitive deterioration in PD (Caviness et al. 2015; Klassen et al. 2011). In a prospective EEG study, slower background frequency during REM sleep together with increased delta and theta band powers predicted the later development of PD dementia (Latreille et al. 2016).

With recent technological advancements, digital biomarkers form a rapidly emerging field of research to improve the longitudinal tracking of neurodegenerative diseases (Espay et al. 2016; Hansen et al. 2018; Baker et al. 2019; Kourtis et al. 2019; Mahadevan et al. 2020). Digital biomarkers refer to the use of inbuilt sensors in portable (e.g. smartphone), wearable (e.g. smart watch) or implantable devices allowing for continuous active and/or passive data collection of biological (e.g. blood glucose), physiological (e.g. heart rate or blood pressure) or functional (e.g. motor activities, speech or facial expression) parameters (Dorsey et al. 2017).

### **2.3.5 Current treatments for Parkinson's disease**

Today, there is no cure available for PD patients. The current therapies ameliorate the disease associated motor symptoms by replenishing striatal dopamine deficiency with the BBB-penetrating dopamine precursor L-DOPA or boosting dopaminergic transmission with dopamine receptor agonists or drugs that prevent the degradation of dopamine (Armstrong and Okun 2020; Fox et al. 2018). The efficacy of these dopamine-based therapies, however, gradually diminishes as the disease progresses and they are unable to delay or reverse the neurodegenerative processes underlying the disease. Thus, there is an unmet medical need for a disease-modifying therapy for PD. The currently used treatments for motor and non-motor symptoms of PD are presented in Table 2.2.

L-DOPA has been the cornerstone of PD treatment since the seminal experiments in 1961 by Walther Birkmayer and Oleh Hornykiewicz who showed the striking effect of the first intravenous injections of L-DOPA on akinetic PD patients (Birkmayer and Hornykiewicz 1961; Hornykiewicz 2002). L-DOPA, given in combination with a peripheral AADC inhibitor, has remained the “the gold standard” of PD treatment until today due to its efficacy, tolerability and low cost (Lewitt 2015).

Although L-DOPA has remarkable therapeutic efficacy in the early-stage PD, the progressive loss dopamine neurons results in narrowing therapeutic window (Armstrong and Okun 2020; Chou et al. 2018; Connolly and Lang 2014). The required dose escalations, the short half-life of L-DOPA and its intermittent availability after oral administration cause fluctuations in drug levels in the brain. Eventually, the effect of a L-DOPA dose starts to wear off faster causing the re-emerge of parkinsonian motor and non-motor symptoms (“off” state) while the highest L-DOPA concentrations cause troublesome dyskinesias, i.e. involuntary, nonrhythmic chorea-type movements.

The motor fluctuations between the “off” state and the “on” state, when the symptoms are controlled but often associated with peak-dose dyskinesias, are proposed to result from the reduced capacity of the degenerating synaptic terminals to store extra dopamine and alterations in the striatal postsynaptic signaling cascades (Chou et al. 2018; Marsden and Parkes 1976). The pathogenic mechanisms of dyskinesias remain to be fully elucidated but appear to include pulsatile stimulation of dopamine receptors, conversion of L-DOPA to dopamine in serotonin neurons and subsequent unphysiological release of dopamine from the striatal serotonergic terminals as a “false neurotransmitter”, corticostriatal glutamatergic overdrive, stimulation of

nAChRs on dopaminergic terminals and aberrant activation of direct-pathway MSNs (Chase and Oh 2000; Espay et al. 2018).

Over time, the motor fluctuations become increasingly difficult to manage and their prevalence increases (Armstrong and Okun 2020; Chou et al. 2018; Connolly and Lang 2014). A common strategy to reduce them is to divide L-DOPA dosage into smaller but more frequent doses. Various combinations of medications are also commonly used together with L-DOPA to increase therapeutic benefit while limiting the high dose-related side-effects. The most common adjunct medications are dopamine receptor agonists and MAO-B and COMT inhibitors (Fox et al. 2018). Subcutaneous apomorphine (D1 and D2 agonist) injections, self-administered via an injection pen, can be used to achieve faster medication response for those who suffer from severe “off” periods.

The abovementioned strategies are effective to a certain extent, but as the disease advances, some patients become unresponsive to conventional medication adjustments. In these cases, device-aided interventions such as intracranial deep brain stimulation (DBS), small dose infusion of L-DOPA-carbidopa gel into jejunum or continuous subcutaneous apomorphine infusion may be considered to treat difficult motor fluctuations and dyskinesias (Armstrong and Okun 2020; Fox et al. 2018; Martinez-Martin et al. 2015; Volkmann et al. 2013). The selection between these advanced therapy options is made based on case-by-case evaluation.

In addition to the pharmacological interventions, physical exercise regimens, physiotherapy and occupational therapy may be useful in improving motor performance or delaying the disease progression (Armstrong and Okun 2020; Fox et al. 2018; Mak et al. 2017). Strength and aerobic endurance training programs appear to produce long-lasting positive effects on the physical functioning of PD patients. Balance, gait, Tai chi and dance exercises have also been shown to reduce falls and improve mobility and walking capacity. The physical exercise interventions should be sustained, intensive, multimodal and start at an early stage of the disease in order to produce maximal benefits. Additionally, speech therapy seems to be a useful strategy for managing hypophonia and dysphagia that are common symptoms in PD.

The non-motor symptoms substantially increase the clinical and financial burden of PD but remain largely refractory to the existing dopaminergic drugs (Armstrong and Okun 2020; Connolly and Lang 2014; Seppi et al. 2019). The management of non-motor symptoms is symptomatic and extrapolated from the general treatment guidelines for these symptoms in non-PD individuals with certain disease-specific elaborations. Some symptoms, such as MCI, have remained without evidence-based treatment options.

**Table 2.2. Currently used treatments for motor and non-motor symptoms of Parkinson's disease.** References provided in the main text

Category	Interventions	Mechanism or mode of action	Indications	Common adverse effects
L-DOPA preparations	L-DOPA-carbidopa/ benserazide, <i>p.o.</i> immediate release	L-DOPA: Precursor of DA, crosses the BBB → AADC converts to DA in the CNS → increases DA content in the remaining DA neurons	Early symptomatic monotherapy, motor fluctuations	Nausea, vomiting, diarrhea, loss of appetite, orthostatic hypotension, dizziness, cardiac arrhythmia, dyspnea, sleepiness, insomnia, restlessness, depression, confusion, hallucinations, strange dreams, motor fluctuations, peak-dose dyskinesias
	L-DOPA-carbidopa/ benserazide, <i>p.o.</i> extended release	Carbidopa/ benserazide: peripheral AADC inhibitor → prevents the conversion of L-DOPA to DA outside the brain → increases L-DOPA concentration in the CNS and reduces peripheral side-effects of DA		
	L-DOPA-carbidopa intestinal gel infusion	See above, reduces fluctuations in the plasma concentration of L-DOPA	Advanced motor fluctuations and dyskinesias that are unresponsive to conventional medication adjustments	See above + weight loss, constipation, falls, anxiety, neuropathy + complications of surgical device insertion, abdominal pain, irritation or infection at the stoma, tube dislocation or obstruction
Non-ergot DA agonists	Pramipexole	Activates postsynaptic D2-like DA receptors	Early symptomatic monotherapy, L-DOPA adjunct, motor fluctuations, depression	Nausea, vomiting, constipation, sleepiness, orthostatic hypotension, dizziness, edema, hallucinations, impulsive-compulsive disorders*
	Ropinirole		Like Pramipexole - depression	
	Rotigotine (transdermal patch)			
	Apomorphine, intermittent s.c. injections	Activates postsynaptic D2- and D1-like DA receptors	Motor fluctuations	The same as other non-ergot DA agonists. In addition, yawning, runny nose, heart palpitation, dyskinesias, priapism, injection/infusion site reactions and nodules
	Apomorphine, continuous s.c. infusion		Advanced motor fluctuations that are unresponsive to conventional medication adjustments	
MAO-B inhibitors	Selegiline	Blocks the degradation of DA to DOPAL in the CNS → increases DA concentration in the CNS	Early symptomatic monotherapy, L-DOPA adjunct, motor fluctuations	Headache, exacerbation of L-DOPA adverse effects
	Safinamide			Headache, arthralgia, dyspepsia, constipation, flu-like syndrome, exacerbation of L-DOPA adverse effects
	Rasagiline			
COMT inhibitors	Entacapone	Blocks the degradation of L-DOPA to 3-OMD in periphery → increases L-DOPA concentration in the CNS	Motor fluctuations	Dark-colored urine, GI complications, dry mouth, increased sweating, angina pectoris, exacerbation of L-DOPA adverse effects
	Opicapone			Like Entacapone + muscle spasms + increased plasma levels of creatine kinase
	Tolcapone	Blocks the degradation of L-DOPA to 3-OMD in periphery as well as L-DOPA and DA degradation to 3-OMD and 3-MT in the CNS, respectively → increases L-DOPA and DA concentration in the CNS		Like Entacapone + hepatotoxicity
	Anticholinergics <sup>a</sup>	Biperiden		Centrally active muscarinic receptor (especially M1) antagonist → reduces the relative cholinergic overactivity in basal ganglia
Trihexyphenidyl				
NMDA receptor antagonist	Amantadine	Not completely understood. Blocks NMDA and nAChRs → decreases corticostriatal glutamatergic activity; increases DA and NA release and blocks reuptake of DA	Early symptomatic monotherapy <sup>b</sup> , L-DOPA adjunct <sup>b</sup> , dyskinesias	Orthostatic hypotension, nausea, dry mouth, blurred vision constipation, urinary retention, edema, insomnia, confusion, hallucinations, livedo reticularis
Atypical antipsychotic	Clozapine	5-HT <sub>2</sub> receptor favoring antagonist with lower affinity for D1 and D2 receptors	Dyskinesias	Sleepiness, dizziness, sedation, tachycardia, constipation, orthostatic hypotension, drooling, risk of agranulocytosis → regular monitoring of the blood count

## Treatments for non-motor symptoms

Functional neurosurgery	Bilateral STN DBS <sup>a</sup>	Not completely understood. High frequency electric stimulation via implanted electrodes causes global inhibition of the target nucleus → mitigates abnormal oscillations in basal ganglia	Advanced motor fluctuations and dyskinesias that are unresponsive to conventional medication adjustments	Dysarthria, impaired working memory, falls, surgery related complications, infections, device related problems, depression
	Bilateral GPI DBS <sup>a</sup>			Like Bilateral STN DBS - depression
	Unilateral VIM DBS <sup>a</sup> or thalamotomy	Mitigates abnormal oscillations in the thalamus	Medication-refractory tremor as the main symptom	Like Bilateral STN DBS
Cholinesterase inhibitors	Rivastigmine	Decrease the hydrolysis of ACh → increases ACh concentration at cholinergic synapses in the brain	Dementia	GI complications, weight loss, bradycardia, vivid dreams, risk for exacerbation of rest tremor
	Donepezil			
	Galantamine			
Antidepressants	Fluoxetine, paroxetine, citalopram, sertraline	SSRI	Depression	Nausea, anorexia, drowsiness, sexual dysfunction, akathisia
	Venlafaxine	SNRI		Drowsiness, insomnia, sexual dysfunction, gastrointestinal symptoms
	Nortriptyline, desipramine, amitriptyline	Tricyclic antidepressants		Anticholinergic side-effects**, orthostatic hypotension, ventricular arrhythmias, heart block, sleepiness, sexual dysfunction, weight gain
Atypical antipsychotics	Clozapine	5-HT <sub>2</sub> receptor favoring antagonists with lower affinity for D1 and D2 receptors (First, potentially hazardous medications such as anticholinergics, amantadine, DA agonists, MAO-B and COMT inhibitors are reduced)	Psychosis, hallucinations	See above
	Quetiapine			Extrapyramidal symptoms, sedation
	Pimavanserin			Nausea, constipation, edema, confusion, weight gain
Hypnotic	Eszopiclone	Positive allosteric modulator of GABA <sub>A</sub> receptor → increases Cl <sup>-</sup> influx into the neurons	Insomnia	Headache, dry mouth, nausea, and dizziness
Hormone	Melatonin	Melatonin receptor agonist → regulates physiological sleep-wake cycle	Insomnia, RBD	Not common, rarely daytime sleepiness, dizziness, headache
Tranquilizer	Clonazepam <sup>a</sup>	Positive allosteric modulator of GABA <sub>A</sub> receptor → increases Cl <sup>-</sup> influx into the neurons	RBD	Sedation, drowsiness confusion
Mineralocorticoid	Fludrocortisone	Increases the reabsorption of Na <sup>+</sup> and water in the kidney → increases blood volume and blood pressure	Orthostatic hypotension	Hypertension, metabolic abnormalities, GI complications, myopathy
Sympathomimetics	Midodrine	Activates adrenergic α <sub>1</sub> -receptors in the vasculature → increases vascular tone and blood pressure		Hypertension, nausea, weakness, heartburn, headache, chills
	Droxidopa	Precursor of NA, AADC converts to NA in the PNS → maintains blood pressure and blood flow		Hypertension, tachycardia, nausea, vomiting, headache
Osmotic laxative	Macrogol	Inert substance that osmotically retains water in the bowel → softens and increases the volume of the stool	Constipation	Electrolyte imbalance, increased bowel gas, abdominal pain, nausea, diarrhea
Dietary supplements	Probiotics and prebiotic fiber	Stimulate the growth and activity of beneficial micro-organisms in the gut → aid digestion		Bloating, increased bowel gas, diarrhea
Prostaglandin E1 derivative	Lubiprostone	Activates Cl <sup>-</sup> -channels on the apical surface of GI epithelial cells → increases fluid secretion, softens the stool and promotes bowel motility		Nausea, diarrhea, headache, bloating, abdominal pain, flatulence
Antidiuretic hormone analog	Desmopressin	Increases the presence of aquaporin channels in the distal nephron by binding to V2 receptors → increases water reabsorption from urine → decreases the total amount of urine	Neurogenic bladder dysfunction	Hyponatremia, hypertension, edema, urinary tract complications, headache, dizziness, GI complications, sleepiness
Neurotoxin	Botulinum toxin injection into salivary glands	Prevents the release of ACh from the postganglionic parasympathetic axon terminals → reduction of saliva production	Drooling	Dysphagia, dry mouth, injection-related discomfort

3-OMD, 3-O-methyldopa; ACh, acetylcholine; DA, dopamine; GI, gastrointestinal; NA, noradrenaline; nAChR, nicotinic acetylcholine receptor; p.o., peroral; s.c., subcutaneous; SNRI, serotonin and noradrenaline reuptake inhibitor; SSRI, selective serotonin reuptake inhibitor; VIM, ventralis intermedius nucleus of the thalamus; see text for other abbreviations.

\* Impulsive-compulsive disorders include pathological gambling, hypersexuality, binge eating, compulsive shopping and overuse of dopaminergic medications.

\*\* Anticholinergic side-effects include cognitive impairment, confusion, tachycardia, vertigo, impaired accommodation, increased intraocular pressure, dry mouth, urinary retention and constipation.

<sup>a</sup> For cognitively intact patients; <sup>b</sup> Not commonly used anymore

## **2.4 Animal models of Parkinson's disease**

In spite of fast developments in *in vitro* disease models, organoids and computer-aided drug development tools, none of these techniques have been able to completely replace the pre-clinical studies in animal models. *In vivo* experiments still form a crucial step where the efficacy and safety of a novel therapy can be tested in a complex, whole-organism setting before entering into the first human trials. Furthermore, animal models provide a powerful means to gain deeper understanding of pathogenic mechanisms which, in turn, contribute to establishing novel targets for disease-modifying therapies. The 3R principles lay the groundwork for performing responsible and humane animal research: they aim at replacing, reducing and refining animal testing whenever possible.


Reflecting the complex nature of the disease, a wide array of strategies and organisms has been used to produce *in vivo* models of PD (Dauer and Przedborski 2003; Gubellini and Kachidian 2015). They all model certain aspects of the disease, but none of the current models can fully recapitulate the neuropathology and symptoms of PD. Although non-mammals provide simple and powerful systems to model the pathobiology of PD, the focus here is on the most widely used rodent and NHP models. They can be classified into four main categories: pharmacological, neurotoxin-based, genetic and  $\alpha$ -syn pathology spreading models. The main features of these models are summarized in Figure 2.11.

### **2.4.1 Pharmacological models**

Reserpine and alpha-methyl-*p*-tyrosine (AMPT) are pharmacological agents that have been used to induce parkinsonism in rodents and NHPs (Carlsson et al. 1957; Windle and Cammermeyer 1958; Bezard and Przedborski 2011). They deplete brain dopamine and other catecholamines by inhibiting VMAT2 (reserpine) or TH (AMPT) while leaving dopamine neurons intact. Striatal dopamine depletion leads to PD-like motor deficits.

### **2.4.2 Neurotoxin-based models**

Neurotoxins produce a prominent and rapid degeneration of the nigrostriatal dopamine system resulting in a clear motor impairment in experimental animals. 6-OHDA is a widely used catecholaminergic neurotoxin (Simola et al. 2007). 6-OHDA is a hydrophilic structural analogue of catecholamines that is unable to cross the BBB. Thus, it needs to be directly injected into the target brain structure in a stereotaxic surgery. Prophylactic peripheral administration of a noradrenaline transporter blocker, such as desipramine, is common to prevent the uptake of 6-OHDA into noradrenergic nerve terminals.



**Figure 2.11. Animal models of Parkinson's disease.** The main characteristics of pharmacological, neurotoxin-based, genetic and alpha-synuclein pathology spreading models of PD. AMPT - Alpha-methyl-*p*-tyrosine; α-syn+/- - alpha-synuclein positive/negative; BAC - bacterial artificial chromosome; BBB - blood-brain barrier; DA - dopamine; DAT - dopamine transporter; ER - endoplasmic reticulum; LB - Lewy body-like; MAO-B - monoamine oxidase B; MFB - medial forebrain bundle; NHP - non-human primate; PFF - α-syn preformed fibrils; ROS - reactive oxygen species; SN - substantia nigra; TFAM - mitochondrial transcription factor A; Tg - transgenic, TH - tyrosine hydroxylase; VMAT2 - vesicular monoamine transporter 2. References are provided in the text. Figure drawn by the author and Fanni-Sofia Renko.

Urban Ungerstedt developed a widely used unilateral 6-OHDA lesion model of PD (Ungerstedt 1968; Ungerstedt and Arbuthnott 1970; Ungerstedt 1971). Unilateral injection of 6-OHDA into the SNpc, medial forebrain bundle (MFB) or dorsal striatum induces degeneration of nigrostriatal dopamine neurons on the injected side accompanied with contralateral motor impairment (Perese et al. 1989; Sauer and Oertel 1994; Kirik et al. 1998; Deumens et al. 2002). In severe lesion models (SNpc and MFB lesions), marked dopamine depletion in the striatum leads to supersensitization of postsynaptic dopamine receptors. Unilateral lesions produce asymmetrical motor deficits that can be easily assessed with behavioral tests. For example, amphetamine-induced dopamine release in the intact striatum elicits ipsilateral rotational behavior, whereas activation of supersensitized dopamine receptors in the lesioned striatum by apomorphine induces turning to the contralateral direction. Important advantages of unilateral lesion model include increased survival of the experimental animals as compared to bilateral lesions and the possibility to use contralateral hemisphere as an internal control in the analyses.

MPTP is another dopaminergic neurotoxin most commonly used to model PD in mice and NHPs, but not in rats which were found to be resistant to the toxin (Bezard and Przedborski 2011; Dauer and Przedborski 2003). Lipophilic MPTP readily crosses the BBB, and thus is suitable for systemic administration. In the brain, MAO-B converts MPTP to the toxic metabolite MPP<sup>+</sup>. Utilizing mitochondrial transmembrane potential MPP<sup>+</sup> concentrates into the mitochondria where it inhibits the complex I of the electron transport chain (Chiba et al. 1984; Nicklas et al. 1985). MPTP intoxication also causes ER stress, neuroinflammation and upregulation of pro-apoptotic Bax and downregulation of anti-apoptotic Bcl-2 in dopamine neurons (Vila et al. 2001; Mercado et al. 2013; Lee et al. 2019). In mice, strain, gender, age and body weight affect the sensitivity to MPTP (Jackson-Lewis and Przedborski 2007). Prolonged subcutaneous or intraperitoneal administration of low or moderate doses of MPTP to C57/Bl6 mice produces a robust PD-like phenotype with defined apoptosis of the SNpc dopamine neurons and impaired motor functions. A more recent model was characterized by Prediger and colleagues who administered MPTP to rodents via an intranasal infusion (Prediger et al. 2011, 2010, 2006). Intranasal MPTP delivery resulted in olfactory, cognitive, emotional and motor defects, loss of TH-ir cells in the olfactory bulb and SNpc, decreased fiber density in the striatum as well as dopamine depletion in the olfactory bulb, striatum and prefrontal cortex. These features appear analogous to many of those observed during the prodromal and early stages of PD making the model a valuable tool for testing novel therapeutic strategies to restore sensory and cognitive deficits of early PD. The MPTP monkey model, however, remains the gold standard model in the preclinical assessment of novel treatment strategies as a final step before proceeding to clinical trials (Burns et al. 1983; Emborg 2007; Porras et al. 2012).

Rotenone is a natural cytotoxic compound used as an insect and fish poison. Systemically administered rotenone reproduces many features of PD but can cause high mortality in experimental animals (Heikkilä et al. 1985b; Betarbet et al. 2000; Gubellini and Kachidian 2015). The dopaminergic damage caused by rotenone can be variable or difficult to reproduce. Paraquat (N,N'-dimethyl-4,4'-bipyridinium), a structural analog of MPP<sup>+</sup>, is a potent and widely used herbicide that produces progressive, but poorly reproducible, dopaminergic neurodegeneration and accumulation of  $\alpha$ -syn immunoreactive inclusions after systemic administration (Cochemé



and Murphy 2008; Manning-Bog et al. 2002; McCormack et al. 2002; Ossowska et al. 2005; Rappold et al. 2011; Richardson et al. 2005; Shimizu et al. 2001).

Lactacystin is a selective proteasome inhibitor. When stereotactically injected into the nigrostriatal pathway, it disturbs cellular protein homeostasis, causes dose-dependent formation of cytoplasmic,  $\alpha$ -syn and ubiquitin containing protein inclusions and triggers dopaminergic neurodegeneration (McNaught et al. 2002b; Bentea et al. 2017). Other proteasome inhibitors, for example systemic administration of PSI (*N*-carbobenzoxy-L-isoleucyl-L- $\gamma$ -*t*-butyl-L-glutamyl-L-alanyl-L-leucinal), have also been tested to model PD in rodents and NHPs with equivocal results (Bentea et al. 2017).

### 2.4.3 Genetic models

A number of the genetic causes of familial PD have been modeled in mice and rats (Dawson et al. 2010). The most common genetic models are derived from transgenic overexpression of autosomal dominant genes (*SNCA* and *LRRK2*) or knockout of autosomal recessive genes (*PRKN*, *PINK1* and *DJ-1*). Even though these models represent direct etiological causes of PD, and thus should replicated relevant pathogenic mechanisms, they fail to produce robust nigrostriatal degeneration, striatal dopamine depletion or motor impairments. Nevertheless, they provide valuable tools to dissect the pathophysiological roles of these proteins.

Duplication or triplication of *SNCA* gene leads to a rare form of familial early onset PD with a clear gene dosage effect on the severity of the disease. This suggests that the level of  $\alpha$ -syn dictates its toxicity and has spurred the creation of several animal models based on the overexpression of human or murine wild-type  $\alpha$ -syn (Dawson et al. 2010; Visanji et al. 2016). Other  $\alpha$ -syn transgene models overexpress PD-related missense mutations of *SNCA* (e.g. A30P or A53T). Viral vectors, such as adeno-associated virus (AAV) or lentivirus, offer another approach to deliver  $\alpha$ -syn transgene into the brain of wild-type animals which has been shown to recapitulate the essential neuropathological features of PD (Kirik et al. 2003, 2002; Lo Bianco et al. 2002). It is worthwhile to note, however, that transgenic  $\alpha$ -syn models rely on the expression of supraphysiological levels of  $\alpha$ -syn which might reduce their validity in mimicking sporadic PD pathology.

Knockout of *PRKN*, *PINK1* or *DJ-1* genes seems to be insufficient to perturb the survival or function of nigrostriatal dopamine neurons or to induce formation of  $\alpha$ -syn positive protein inclusions (Dawson et al. 2010; Blesa and Przedborski 2014). Due to the absence of consistent parkinsonian phenotype in single knockout mice, a triple knockout model with parkin, DJ-1 and PINK1 deletion was created (Kitada et al. 2009). However, even the inactivation of all three recessive PD genes was insufficient to induce loss of nigral dopamine neurons within the mouse lifespan. Only bacterial artificial chromosome (BAC) -mediated overexpression of truncated human *PRKN* (Q311X) in dopamine neurons has been shown to produce age-dependent degeneration of nigrostriatal dopamine system, accumulation of  $\alpha$ -syn positive protein inclusions and late-onset motor deficits in mice (Lu et al. 2009).

MitoPark is an example of a genetic PD model that is not derived from any of the known PD-related genetic risk factors. In MitoPark mice, *TFAM* gene encoding mitochondrial transcription factor A (TFAM) is conditionally deleted from dopaminergic neurons (Ekstrand et al. 2007; Galter et al. 2010; Good et al. 2011). TFAM is required for the transcription of mitochondrial DNA coding for essential subunits of the respiratory chain. MitoPark mice show apparent PD-like phenotype both on pathological and behavioral level.

#### **2.4.4 Alpha-synuclein pathology spreading model**

Recombinant  $\alpha$ -syn preformed fibrils (PFFs) can be inoculated into various brain regions to induce propagating Lewy body-like pathology both in mice with different genetic backgrounds, rats and NHPs (Luk et al. 2012; Luk et al. 2012b; Rey et al. 2016; Abdelmotilib et al. 2017; Karampetsou et al. 2017; Thakur et al. 2017; Shimozawa et al. 2017; Chu et al. 2019; Henderson et al. 2019). PFF-induced  $\alpha$ -syn pathology triggers reactive microgliosis prior to dopaminergic degeneration in the SNpc suggesting that neuroinflammation contributes the vulnerability of dopamine neurons (Duffy et al. 2018). One advantage of this model is that the PD-like phenotype develops slower than in neurotoxin or AAV- $\alpha$ -syn overexpression models offering a therapeutic window for experimental interventions (Okuzumi et al. 2018).

#### **2.4.5 Behavioral and histological assessment in animal models**

As described in chapter 2.2, the nigrostriatal dopamine system plays a key role in regulating motor behavior. Thus, the degree of nigrostriatal degeneration in animal models of PD can be followed longitudinally by measuring motor deficits with specific behavioral tests (Asakawa et al. 2016; Dunnett and Lelos 2010). The behavioral tests that are commonly used to measure motor impairment in rodents include, but are not restricted to, drug-induced rotational behavior (rotametry), limb-use asymmetry (cylinder), open field, rotarod, stepping, skilled paw-reaching (staircase) and adhesive removal tests. The non-motor symptoms of PD can also be measured with various behavioral tests which are not, however, specifically designed for rodent models of PD. The brains of NHPs are anatomically and physiologically closest to the human brains making NHPs invaluable subjects in PD research (Emborg 2007; Porras et al. 2012). NHPs replicate almost all of the human parkinsonian motor symptoms after neurotoxin administration enabling much more versatile behavioral assessments as compared to other species, for example the use of various rating scales that imitate the clinically used Unified Parkinson's disease Rating Scale (UPDRS).

When studying disease-modifying therapies in animal models of PD, the decisive neuropathological outcomes (e.g. neuroprotective or restorative effects) can be obtained only with cellular and molecular level investigation after tissue collection. Histologically, the integrity of the nigrostriatal pathway can be measured by immunolabelling dopaminergic markers such as TH, DAT or VMAT2 and quantifying the number of immunoreactive cell bodies in the SNpc and the density of immunoreactive fibers in the dorsal striatum. Lewy body-like pathology can be visualized for example by immunostaining phosphorylated  $\alpha$ -syn. In histological assessments, it is important to note that the expression levels of individual molecular markers are under divergent

regulation. Thus, a combination of markers should be favored in order to draw reliable conclusions.

## **2.5 Preclinical studies of neurotrophic factors for Parkinson's disease**

Due to the dopaminotrophic properties of GDNF, NRTN, CDNF and MANF, their potential as a disease-modifying therapy for PD has been examined extensively both *in vitro* and *in vivo*. Here, an overview of the key preclinical studies, with the main focus on *in vivo* experiments in normal and lesioned nigrostriatal dopamine system, will be provided.

### **2.5.1 Effects of neurotrophic factors on dopaminergic function of intact nigrostriatal system**

NTF delivery into the brain of a PD patient would be likely to affect both degenerating and intact neurons. Therefore, knowledge on the effects of exogenous recombinant NTFs not only on the pathological processes but also on the physiological functions of the brain is pivotal. Characterization of the effects of NTFs on normal, non-lesioned basal ganglia circuitry allows for improving their therapeutic efficacy and safety profile.

#### **2.5.1.1 Neurochemical and functional effects of exogenous GDNF**

The exogenous administration of GDNF elicits long-lasting neurochemical and functional changes in the intact nigrostriatal system. GDNF augments dopamine, DOPAC and HVA levels in tissue samples and induces spontaneous motor activity in young and aged animals (Hudson et al. 1995; Hebert et al. 1996; Martin et al. 1996; Hebert and Gerhardt 1997; Kobayashi et al. 1998; Grondin et al. 2003; Georgievska et al. 2004; Hadaczek et al. 2010). Dopamine turnover, measured as HVA/dopamine ratio, reflects the strength of dopaminergic transmission and is enhanced in GDNF-treated animals. The abovementioned effects, however, are not specific to the dopaminergic system, since nigrostriatal serotonin, 5-hydroxyindoleacetic acid (5-HIAA) and noradrenaline levels are also altered. Intracranial GDNF injections are also associated with reduced food intake and body weight in experimental animals. Noteworthy, unilateral striatal delivery of GDNF has been shown to produce bilateral effects on dopamine neurochemical markers, neuronal activity and motor functions in normal animals as well as in PD patients (Grondin et al. 2003; Salvatore et al. 2009; Slevin et al. 2005; Stanford et al. 2007).

GDNF has been shown to modulate the activity of dopaminergic nerve terminals. GDNF application increases stimulus-evoked dopamine release in cultured midbrain dopamine neurons as well as in striatal synaptosomes and slices (Feng et al. 1999; Gomes et al. 2006). Amperometric recordings showed that in primary cultures GDNF exposure increased the quantal size of dopamine-containing synaptic vesicles (Pothos et al. 1998). With patch clamp technique, GDNF was demonstrated to acutely potentiate the excitability and synaptic transmission of dopamine neurons in midbrain cultures and slices by inhibiting A-type K<sup>+</sup> channels and potentiating high voltage-gated Ca<sup>2+</sup> channel currents (Bourque and Trudeau 2000; Yang et al. 2001; Wang et al.

2003). Similar effects have also been observed *in vivo* in brain microdialysis measurements. Intracranial administration of GDNF augmented stimulus-evoked overflow of striatal dopamine in non-lesioned rats (Hebert et al. 1996; Hebert and Gerhardt 1997; Cass et al. 1999; Salvatore et al. 2004; Cass and Peters 2010) and NHPs (Gash et al. 1995; Grondin et al. 2003). These microdialysis studies, however, were conducted under general anesthesia which is known to have profound effects on neuronal activity, neurotransmitter synthesis, release, reuptake and metabolism (Marinelli and McCutcheon 2014; Müller et al. 2011). In a microdialysis study with freely-moving rats, no significant differences in potassium-evoked dopamine output was detected between GDNF- and vehicle-injected animals (Xu and Dluzen 2000).

GDNF-induced enhancements in dopaminergic function are associated with a sustained increase in TH phosphorylation and activity. Exogenous GDNF has been shown to downregulate the expression of TH but increase its phosphorylation and activity (Beck et al. 1996; Rosenblad et al. 2003; Georgievska et al. 2004; Kobori et al. 2004; Salvatore et al. 2004, 2009). The downregulation of TH may be a compensatory response to its enhanced activity. In line with this, *in vivo* TH activity is also elevated in MEN2B (multiple endocrine neoplasia type 2B) mice with constitutively active RET (Mijatovic et al. 2008).

#### 2.5.1.2 Neurochemical and functional effects of other NTFs

Data on the molecular and functional effects of other exogenously administered GFLs, CDNF or MANF on the intact nigrostriatal system are scarce. Similarly to GDNF, striatal administration of NRTN has been shown to provoke some behavioral and neurochemical changes associated with functional upregulation of dopamine neurons. In non-lesioned rats, NRTN delivery augmented amphetamine-induced locomotor activity and caused a persistent increase in dopamine turnover (Horger et al. 1998; Hadaczek et al. 2010). A single nigral injection of NRTN increased stimulus-evoked dopamine release in the striatum and tissue levels of dopamine in the SN of normal rats (Cass and Peters 2010). In naïve monkeys, striatal delivery of an AAV2 vector encoding human NRTN (CERE-120, see Table 2.3) increased the number of TH-ir cells in the SNpc and fibers in the striatum (Herzog et al. 2008, 2007). PET imaging demonstrated that  $^{18}\text{F}$ -dopa uptake was increased in the NTRN-transduced striatum as compared to the control side.

#### 2.5.1.3 Diffusion properties of GDNF, NRTN, CDNF and MANF in the intact brain

The volume of distribution of exogenous GDNF and NRTN in the brain parenchyma is limited due to their high-affinity binding to extracellular matrix-associated heparan sulfates (Hamilton et al. 2001; Piltonen et al. 2009; Runeberg-Roos et al. 2016). Poor diffusion of GDNF has also been demonstrated in the normal rhesus monkey brain (Lapchak et al. 1998; Salvatore et al. 2006). Intrastratially administered MANF, instead, diffuses significantly better than GDNF in the intact rat brain distributing throughout the striatum and frontal cortex (Voutilainen et al. 2011, 2009). CDNF also diffuses rapidly in the intact rat brain resulting in widespread distribution that extends almost over the entire hemisphere at the level of the striatal infusion site (Mätlik et al. 2017).

## 2.5.2 Effects of neurotrophic factors in animal models of Parkinson's disease

According to the initial findings, GDNF selectively promotes the survival and morphological differentiation of dopamine neurons in rat embryonic midbrain cultures and enhances their high-affinity dopamine uptake (Lin et al. 1993). GDNF is also able to support the viability of nigral dopamine neurons in postnatally derived primary cultures by inhibiting natural cell death (Burke et al. 1998). In *in vitro* neurotoxin models, GDNF prevents the loss of cultured midbrain dopamine neurons in response to MPP+ or 6-OHDA (Eggert et al. 1999) and promotes the recovery and regrowth of damaged dopamine neurons after the removal of the toxin challenge (Hou et al. 1996; Kramer et al. 1999). Likewise, NRTN supports the survival of dopamine neurons in embryonic midbrain cultures with similar potency and efficacy to that observed for GDNF (Cacalano et al. 1998; Horger et al. 1998; Runeberg-Roos et al. 2016). MANF was shown to selectively increase the survival and sprouting of dopamine neurons in embryonic midbrain cultures (Petrova et al. 2003). CDNF protected primary neurons in embryonic midbrain cultures against cytotoxicity induced by 6-OHDA and  $\alpha$ -syn oligomers (Latge et al. 2015). The promising outcomes in *in vitro* studies have encouraged testing the neuroprotective and regenerative effects of GDNF, NRTN, CDNF and MANF in various animal models of PD. An overview of these *in vivo* studies is provided in Table 2.3.

Notably, according to toxin-induced neurorestoration models, GDNF delivery into the striatum seems to provide better protection on TH-ir fibers in the striatum and result in more robust functional recovery as compared to nigral delivery. GDNF delivery to its physiological site of expression (i.e. the striatum) may be an underlying cause for this discrepancy (Kirik et al. 2004; Ibáñez and Andressoo 2017).

A combined lentiviral vector-mediated transduction of CDNF and MANF into the SNpc led to synergistic neuroprotective effects in 6-OHDA-lesioned rats with reduced amphetamine-induced turning behavior and robust preservation of dopaminergic fibers in the striatum and cell bodies in the SNpc (Cordero-Llana et al. 2015c). Also, when CDNF protein was injected into the striatum of 6-OHDA-lesioned rats together with GDNF, the co-administration resulted in stronger neurorestorative effects than either of the NTFs could produce alone (Voutilainen et al. 2017). The additive effects of the NTFs may stem from different mechanisms of action: both proteins activated the survival promoting PI3K/Akt signaling pathway and CDNF reduced the expression of ER stress-related markers GRP78, ATF6 and phosphorylated eIF2 $\alpha$ .

Continuous NTF infusion may cause ligand-dependent receptor desensitization decreasing the responsiveness of the target tissue to the treatment (Lohse 1993). To circumvent this risk, intermittent convection-enhanced delivery (CED), which enables target engagement in a pulsatile fashion, has been developed and tested in preclinical studies (Bobo et al. 1994; Gash et al. 2005). CED significantly enhances tissue distribution of macromolecules by delivering brief bolus infusions of a treatment solution in addition to low-rate basal infusion.

**Table 2.3. Disease-modifying effects of neurotrophic factors (NTFs) in animal models of Parkinson's disease (PD).**

NTF	Species; PD model	Para- digm <sup>a</sup>	NTF delivery	NTF dose	Neuropathological and neurochemical outcomes	Functional outcomes	Reference
70 GDNF	Rats; Unilateral 6-OHDA-lesion in MFB	Neuro- restoration	Single ipsilateral injection into SN	100 µg	Preserved nigral DA and metabolite levels; increased survival of TH+ neurons and neurites in SN	Reduced APO-induced rotations	(Hoffer et al. 1994); (Bowenkamp et al. 1995)
	Mice; Repeated MPTP s.c. injections	Neuro- protection Neuro- restoration	Single unilateral injection into SN or STR	10 µg	Preserved nigrostriatal DA and metabolite levels, increased survival of TH+ neurons in SN and TH+ fibers in STR Partially preserved nigrostriatal DA and metabolite levels, increased survival TH+ fibers in STR	Increased locomotor activity	(Tomac et al. 1995)
	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- protection	Single ipsilateral injection into SN, STR or LV	25 µg, 25 µg or 50 µg	Increased survival of TH+ neurons in SN (STR and SN delivery), increased survival TH+ fibers in STR (STR delivery)	Normalized contralateral forelimb akinesia in stepping test, reduced APO- and AMPH- induced rotations (STR delivery)	(Kirik et al. 2000)
	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- protection	AAV vector-mediated ipsilateral transduction into SN, STR or SN+STR	4x10 <sup>9</sup> vg (SN), 9x10 <sup>9</sup> vg (STR)	Increased survival of TH+ neurons in SN (SN, STR and SN+STR delivery), increased survival TH+ fibers in STR (STR delivery)	Reduced AMPH-induced rotations, improved spontaneous contralateral paw use in cylinder and staircase test (STR delivery)	(Kirik et al. 2000b)
	Rats; Unilateral 6-OHDA-lesion in SN	Neuro- restoration	Lentiviral vector-mediated ipsilateral transduction into STR	2.6x10 <sup>5</sup> TU	Increased survival TH+ fibers in STR	Improved spontaneous contralateral paw use in staircase test	(Brizard et al. 2006)
	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- protection Neuro- restoration	Single ipsilateral injection into STR	10 µg	Increased survival of TH+ neurons in SN and TH+ fibers in STR Increased survival of TH+ neurons in SN	Reduced AMPH-induced rotations	(Yue et al. 2014)
	Rats; Unilateral lentiviral vector- transduction of A30P ha-syn to SN	Neuro- protection	Lentiviral vector-mediated ipsilateral transduction into SN	?	Lack of protective effect on TH+ neurons in SN and TH+ fibers in STR	N.D.	(Lo Bianco et al. 2004a)
	Rats; Unilateral AAV vector-transduction of wild-type ha-syn to SN	Neuro- protection	Lentiviral or AAV vector- mediated ipsilateral transduction into SN, STR or SN+STR	3x10 <sup>4</sup> TU (SN), 4.5x10 <sup>4</sup> TU (STR)	Lack of protective effect on TH+ neurons in SN and TH+ and VMAT2+ fibers in STR, no effect on the number of α-syn+ aggregates	Failed to reduce AMPH-induced rotations	(Decressac et al. 2011)
	Monkeys; Unilateral 6-OHDA-lesion in MFB	Neuro- protection	AAV vector-mediated ipsilateral transduction into SN and STR	2.4x10 <sup>9</sup> vg (SN), 6x10 <sup>9</sup> vg (STR)	Increased survival of TH+ and VMAT2+ neurons in SN, increased size of TH+ neurons in SN	Improved rating scale scores, reduced AMPH- and APO-induced rotations	(Eslamboli et al. 2003)
	Monkeys; Unilateral MPTP infusion via car. art.	Neuro- restoration	Ipsilateral injection into SN, STR or LV	1x150 µg, 1x450 µg or 3x100/450 µg	Increased cell size and neurite density of TH+ neurons in SN (SN and STR delivery), preserved DA levels in ventral midbrain (ICV delivery)	Improved rating scale scores in bradykinesia, rigidity, posture and balance	(Gash et al. 1996)
	Monkeys; Unilateral MPTP infusion via car. art.	Neuro- restoration	Single ipsilateral injection into LV	300 µg	Preserved nigrostriatal DA and metabolite levels	Improved rating scale scores	(Gerhardt et al. 1999)
	Monkeys; Unilateral MPTP infusion via car. art.	Neuro- restoration	Lentiviral vector-mediated ipsilateral transduction into SN+STR	?	Increased survival of TH+ neurons in SN and TH+ fibers in STR	Increased uptake in <sup>18</sup> F-DOPA-PET, improved rating scale scores, reversed motor deficits in a hand-reach task	(Kordower et al. 2000); (Emborg et al. 2009)
	Monkeys; Unilateral MPTP infusion via car. art.	Neuro- restoration	Chronic infusion to ipsilateral LV or bilaterally into STR	5 or 15 µg/24h	Increased cell size and survival of TH+ neurons in SN, increased survival TH+ fibers in STR, preserved DA and metabolite levels in STR and GP	Improved rating scale scores in bradykinesia, rigidity, posture and balance	(Grondin et al. 2002)
NRTN	Monkeys; Unilateral MPTP infusion via car. art.	Neuro- restoration	Pulsed, CED infusion into ipsilateral SN	7.5 or 22.5 µg/24h	Increased cell size and survival of TH+ neurons in SN	Improved rating scale scores, increased movement speed	(Gash et al. 2005)
	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- restoration	Single or repeated ipsilateral injection into SN	1x1 or 1x10 µg or 7x5 µg (every 3 <sup>rd</sup> day)	Increased survival of TH+ (single injection) and FG+ neurons in SN (single and repeated injections)	N.D.	(Horger et al. 1998)
	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- restoration	Multiple ipsilateral injections into STR or LV	7x5 or 7x10 µg (every 3 <sup>rd</sup> day)	Increased survival of TH+ and FG+ neurons in SN (STR delivery)	No effect on AMPH-induced rotations	(Rosenblad et al. 1999)

CDNF	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- protection Neuro- restoration	Single ipsilateral injection into STR	5 µg	Increased survival of TH+ neurons in SN, preserved striatal DA and metabolite levels Increased survival TH+ fibers in STR, preserved striatal DA and metabolite levels	Reduced MetAMPH-induced rotations, increased DA turnover (restoration study)	(Oiwa et al. 2002)
	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- protection	AAV2 vector (CERE-120 <sup>b</sup> )-mediated ipsilateral transduction into STR	1.6x10 <sup>8</sup> , 8x10 <sup>8</sup> or 4x10 <sup>9</sup> vg	Increased survival of TH+ neurons in SN, increased survival of VMAT2+ fibers in STR	Reduced AMPH-induced rotations, no effect on spontaneous forepaw use in the cylinder test	(Gasmi et al. 2007) (Gasmi et al. 2007b)
	Monkeys; Unilateral MPTP infusion via car. art.	Neuro- restoration	AAV2 vector (CERE-120 <sup>b</sup> )-mediated ipsilateral transduction into SN+STR	Total dose 1.7x10 <sup>11</sup> TU	Increased survival of TH+ neurons in SN and TH+ fibers in STR	Improved rating scale scores	(Kordower et al. 2006)
	Monkeys; Unilateral MPTP infusion via car. art.	Neuro- restoration	Chronic infusion into ipsilateral STR	30 µg/24h	Preserved DA metabolite levels in GP	Improved rating scale scores, increased locomotor activity	(Grondin et al. 2008)
	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- protection Neuro- restoration	Single ipsilateral injection into STR	1-10 µg	Increased survival of TH+ neurons in SN and TH+ fibers in STR (3-10 µg)	Reduced AMPH-induced rotations (3-10 µg)	(Lindholm et al. 2007)
				10 µg	Partially (N.S.) increased survival TH+ neurons in SN		
	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- restoration	Chronic infusion into ipsilateral STR	1.5-4.5 µg/24h	Increased survival of TH+ neurons in SN (3-4.5 µg/24h) and TH+ fibers in STR (1.5-4.5 µg/24h)	Reduced AMPH-induced rotations (3 µg/24h)	(Voutilainen et al. 2011)
	Mice; Repeated MPTP i.p. injections	Neuro- protection Neuro- restoration	Single bilateral injections into STR	10 µg/side	Increased survival of TH+ neurons in SN and TH+ fibers in STR	Increased locomotor activity	(Airavaara et al. 2012)
	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- protection	AAV vector-mediated ipsilateral transduction into STR	4x10 <sup>7</sup> , 2x10 <sup>8</sup> or 1x10 <sup>9</sup> vg	Partially (N.S.) increased survival TH+ neurons in SN (2x10 <sup>8</sup> and 1x10 <sup>9</sup> vg), partially (N.S.) increased survival TH+ fibers in STR (1x10 <sup>9</sup> vg)	Reduced AMPH-induced rotations (2x10 <sup>8</sup> and 1x10 <sup>9</sup> vg)	(Back et al. 2013)
	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- restoration	AAV vector-mediated ipsilateral transduction into STR	?	Increased survival of TH+ neurons in SN and TH+ fibers in STR, preserved TH expression in SN, preserved TH and DAT expression in STR	Reduced AMPH-induced rotations, increased locomotor activity, increased DAT-PET uptake in STR	(Ren et al. 2013)
	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- protection	Lentiviral vector-mediated ipsilateral transduction into SN or STR	6x10 <sup>5</sup> TU	Increased survival of TH+ fibers in STR (SN delivery), lack of protective effect on TH+ neurons in SN (SN and STR delivery) and TH+ fibers in STR (STR delivery)	Reduced AMPH-induced rotations (SN delivery), no effect on AMPH- or APO induced rotations (STR delivery)	(Cordero-Llana et al. 2015c)
	Rats; Unilateral 6-OHDA-lesion in MFB	Neuro- restoration	Single ipsilateral injection into SN	1-100 µg	No effect on TH+ neurons in SN and TH+ fibers in STR	Modestly reduced APO- induced rotations (10 µg)	(Huotari et al. 2018)
	Monkeys; Mild unilateral 6-OHDA-lesion in STR	Neuro- restoration	Ipsilateral injection into STR (two injection sites)	2x10 µg	No effects on TH+ fibers or neurons	Increased DAT-SPECT uptake in STR (proper negative control group and between group comparisons are lacking)	(Garea-Rodríguez et al. 2016)

MANF	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- protection Neuro- restoration	Single ipsilateral injection into STR	3-30 µg	Increased survival of TH+ neurons in SN (10 µg)	Reduced AMPH-induced rotations (10 µg)	(Voutilainen et al. 2009)
				10 µg	Partially (N.S.) increased survival TH+ neurons in SN		
	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- restoration	Chronic infusion into ipsilateral STR	1.5-4.5 µg/24h	Lack of protective effect on TH+ neurons in SN and TH+ fibers in STR	No effect on AMPH-induced rotations	(Voutilainen et al. 2011)
	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- protection	Lentiviral vector-mediated ipsilateral transduction into SN or STR	1.6x10 <sup>6</sup> TU	Increased survival of TH+ neurons in SN (SN delivery), lack of protective effect on TH+ neurons in SN (STR delivery) and TH+ fibers in STR (SN and STR delivery)	No effect on AMPH- or APO induced rotations	(Cordero-Llana et al. 2015c)
	Rats; Unilateral 6-OHDA-lesion in STR	Neuro- restoration	AAV vector-mediated ipsilateral transduction into STR	2x10 <sup>10</sup> vg	Increased survival of TH+ neurons in SN and TH+ fibers in STR, increased striatal DA and metabolite levels	Reduced AMPH-induced rotations	(Hao et al. 2017)

AAV, adeno-associated virus; AMPH, amphetamine; APO, apomorphine; car. art., carotid artery; CED, convection-enhanced delivery; DA, dopamine; DAT, dopamine transporter; FG, fluorogold; GP, globus pallidus; hα-syn, human α-synuclein; i.p., intraperitoneal; LV, lateral ventricle; MFB, medial forebrain bundle; N.D., not determined; N.S., not significant; s.c., subcutaneous; SN, substantia nigra; STR, striatum; TH, tyrosine hydroxylase; TU, transducing units (lentiviral titer); vg, viral genomes (AAV titer); VMAT2, vesicular monoamine transporter 2

<sup>a</sup> In neuroprotection paradigm, NTF is administered prior to, or at the same time as the lesion. A neuroprotective therapy slows down or prevents neurodegeneration. In neurorestoration paradigm, NTF is applied after the lesion when neuronal damage has occurred. Neurorestoration refers to recovery of damaged or impaired neuronal connections, somas, functionality or phenotype.

<sup>b</sup> CERE-120 is an AAV2-derived vector encoding a hybrid form of human NRTN cDNA where the NRTN pre-pro domain is replaced with that of human NGF to ensure enhanced secretion (Ramirez et al. 2004; Gasmi et al. 2007)

## 2.6 Neurotrophic factors in clinical trials for Parkinson's disease

The encouraging outcomes in preclinical studies spurred the assessment of intracerebrally delivered recombinant NTFs in PD patients. To date, four different trophic factors have entered clinical trials in PD: GDNF, NRTN, platelet-derived growth factor (PDGF) and most recently CDNF. A summary of these trials is provided in Table 2.4., excluding the placebo-controlled phase I/II study by Paul et al. (2015) assessing the safety and tolerability of intracerebroventricularly (ICV) infused PDGF-BB, which lies beyond the scope of this thesis.

### 2.6.1 Clinical studies with GDNF

The safety and efficacy of GDNF protein therapy as well as AAV2 vector -mediated gene therapy have been evaluated in several clinical trials in patients with advanced PD (Table 2.4.). In two separate open-label phase I trials, intraputamenal infusion of GDNF led to significantly improved motor function associated with increased  $^{18}\text{F}$ -DOPA uptake in the posterior putamen after one year of treatment and caused no serious adverse effects (Gill et al. 2003; Patel et al. 2005; Slevin et al. 2007, 2005). A case report described that these functional improvements sustained up to three years after the cessation of the protein infusion (Patel et al. 2013). Nevertheless, the primary endpoints in two double-blind, placebo-controlled phase I/II trials were not met (Nutt et al. 2003; Lang et al. 2006). ICV injections of GDNF failed to improve patients' UPDRS scores and caused severe side effects like nausea, vomiting, weight loss, anorexia and paresthesias (Nutt et al. 2003). The lack of efficacy was possibly due to a suboptimal delivery method; a *postmortem* histological analysis of one GDNF-treated subject from the study suggested the lack of target engagement by showing little penetration of GDNF into the brain parenchyma from the lateral ventricle (Kordower et al. 1999). Likewise, intraputamenal infusion of GDNF was unable to improve the off-medication UPDRS motor score although  $^{18}\text{F}$ -DOPA uptake was increased in GDNF-infused patients after six months of treatment (Lang et al. 2006). Intraputamenal infusion of GDNF, however, seemed to be better tolerated as compared to ICV injections: treatment-related adverse events included mainly paraesthesias, headache and upper respiratory tract infections in some subjects. In a few patients, neutralizing antibodies against the recombinant GDNF were detected.

The discrepancies in the outcomes of these first trials could be attributed to insufficient bioavailability of GDNF in the target area (Sherer et al. 2006). The continuous low-rate intraputamenal infusion protocol utilized in Lang et al. (2006) study resulted in spatially limited and heterogenous distribution in the putamen (Salvatore et al. 2006). Intermittent intraputamenal CED delivery, instead, allows much wider and homogeneous exposure across the putamen and avoids receptor desensitization (Bobo et al. 1994; Lohse 1993). Thus, Whone and colleagues set out to evaluate the efficacy and safety of recombinant GDNF administered via intermittent intraputamenal CED infusion in a randomized, placebo-controlled phase II study (Whone et al. 2019; Whone et al. 2019b). Although there was a clinically significant motor improvement in 43% of the GDNF-treated patients and overall increase in putamenal  $^{18}\text{F}$ -DOPA uptake in the GDNF



group, the trial failed to meet its primary efficacy endpoint. The treatment administration setup used in this trial is illustrated in Figure 2.12.

**Table 2.4. Neurotrophic factors (NTFs) in clinical trials for Parkinson's disease (PD).**


NTF	Delivery	Dosing	Phase / design	Patients	Key outcomes	References
GDNF	rhGDNF; lateral ventricle; unilateral bolus injections 1/month	Dose-escalation: 25, 75, 150, 300 or 4000 µg for 6-8 months + 20 months open-label extension	I/II, randomized, double-blind, placebo-controlled	50; moderate to severe PD H&Y 3-5 (off)	No improvement in UPDRS motor scores, various SAEs e.g. paresthesias, nausea, anorexia, weight loss and vomiting	(Nutt et al. 2003)
	rhGDNF; putamen; uni- or bilateral continuous infusion	11–43 µg/putamen/24h for 12 months + 12 months extension at 14-29 µg/putamen/24h	I, open-label	5; advanced PD (>6 years from diagnosis)	57% improvement in UPDRS motor score (off) at 24 months, increased <sup>18</sup> F-DOPA-PET uptake in putamen, sprouting of dopaminergic fibers, no therapy-related SAEs	(Gill et al. 2003) (Patel et al. 2005) (Love et al. 2005)
	rhGDNF; putamen; unilateral CED infusion	Dose-escalation: 3, 10 and 30 µg/day for 6 months + 12 months extension at 30 µg/24h	I, open-label	10; moderate to severe PD H&Y 3-4 (off)	45% bilateral improvement in UPDRS motor score (off) at 12 months, no therapy-related SAEs, anti-GDNF antibodies in some patients	(Slevin et al. 2005) (Slevin et al. 2007)
	rhGDNF; putamen; bilateral continuous infusion	15 µg/putamen/24h for 6 months + open-label extension	II, randomized, double-blind, placebo-controlled	34; advanced PD (>5 years from diagnosis)	No improvement in UPDRS motor score (off), increased <sup>18</sup> F-DOPA-PET uptake in putamen, well tolerated, anti-GDNF antibodies in some patients	(Lang et al. 2006)
	rhGDNF; putamen; bilateral intermittent CED infusions 1/month	120 µg/putamen for 9 months + 9 months open-label extension	II, randomized, double-blind, placebo-controlled	41; moderate PD H&Y 2-3 (off)	No improvement in UPDRS motor score (off), increased <sup>18</sup> F-DOPA-PET uptake in putamen, no therapy-related SAEs	(Whone et al. 2019) (Whone et al. 2019b) NCT03652363 <sup>a</sup>
	AAV2-GDNF; putamen; bilateral CED infusion	Dose-escalation: 9x10 <sup>10</sup> , 3x10 <sup>11</sup> or 9x10 <sup>11</sup> vg	I, open-label	13; moderate to severe PD H&Y 3-4 (off)	No improvement in UPDRS motor scores, increased <sup>18</sup> F-DOPA-PET uptake in putamen, no therapy-related SAEs, anti-GDNF and anti-AAV2 antibodies in some patients	(Heiss et al. 2019) NCT01621581 <sup>a</sup>
NRTN	AAV2-NRTN <sup>b</sup> ; putamen; bilateral multi-site injections	Dose-escalation: 1.3x10 <sup>11</sup> or 5.4x10 <sup>11</sup> vg	I, open-label	12; moderate to severe PD H&Y 3-4 (off)	36% improvement in UPDRS motor score (off) at 12 months, no increase in <sup>18</sup> F-DOPA-PET uptake, no therapy-related SAEs	(Marks et al. 2008) NCT00252850 <sup>a</sup>
	AAV2-NRTN <sup>b</sup> ; putamen; bilateral multi-site injections	5.4x10 <sup>11</sup> vg + double-blind extension up to 18 months (n=30)	II, randomized, double-blind, sham-surgery controlled	58; advanced PD (>5 years from diagnosis)	No improvement in UPDRS motor score (off) at 12 months, no increase in <sup>18</sup> F-DOPA-PET uptake, improved PDQ-39 score at 12 months, improved UPDRS motor score (off) at 18 months (n=30), no therapy-related SAEs	(Marks et al. 2010) NCT00400634 <sup>a</sup>
	AAV2-NRTN <sup>b</sup> ; putamen+SN; bilateral multi-site injections	Dose-escalation: 2x10 <sup>11</sup> vg/SN + 2.7x10 <sup>11</sup> vg/putamen or 2x10 <sup>11</sup> vg/SN + 1.2x10 <sup>12</sup> vg/putamen	I, open-label	6; moderate PD H&Y 2-3 (off)	Well-tolerated, no therapy-related SAEs	(Bartus et al. 2013) NCT00985517 <sup>a</sup>
	AAV2-NRTN <sup>b</sup> ; putamen+SN; bilateral multi-site injections	2x10 <sup>11</sup> vg/SN + 1x10 <sup>12</sup> vg/putamen	II, randomized, double-blind, sham-surgery controlled	51; moderate PD H&Y 2-3 (off)	No improvement in UPDRS motor score (off), no therapy-related SAEs, anti-AAV2 antibodies in some patients	(Olanow et al. 2015) NCT00985517 <sup>a</sup>
CDNF	rhCDNF; putamen; intermittent CED infusions 1/month	Low-dose or high-dose for 6 months + 6 months open-label extension	I/II, randomized, double-blind, placebo-controlled	17; moderate PD H&Y 2.5-3 (off)	Well-tolerated, no treatment-related SAEs, potentially improved UPDRS motor score in some patients at 12 months, increased DAT-PET uptake in putamen of some patients, ongoing analyses of exploratory endpoints	(Huttunen and Saarma 2019) NCT03295786 <sup>a</sup>

AAV2, adeno-associated virus serotype 2; CED, convection-enhanced delivery; H&Y, Hoehn and Yahr scale of Parkinson's disease symptoms; off, off medication; PDQ-39, a 39-item Parkinson's disease Questionnaire to assess health and quality of life; rh, recombinant human; SAE, serious adverse event; SN, substantia nigra; UPDRS, Unified Parkinson's Disease Rating Scale; vg, viral genomes

<sup>a</sup> ClinicalTrials.gov identifier; <sup>b</sup> CERE-120

Viral vector -mediated gene transfer provides another opportunity for targeted delivery of therapeutic proteins into restricted brain regions in a safe and sustained manner. Gene therapy

may also enhance the tissue distribution of the expressed protein. An open-label phase I trial sought to assess the safety and effectiveness of bilateral intraputamenal AAV2-GDNF gene therapy in advanced PD (Heiss et al. 2019). Delivery of AAV2-GDNF using intraoperative MRI-guided CED infusion was safe and well tolerated. It also increased  $^{18}\text{F}$ -DOPA uptake in the putamen suggesting a neurotrophic effect on dopaminergic neurons but failed provide any clinically significant improvements in the UPDRS scores.



**Figure 2.12. Intracranial administration of GDNF to Parkinson's disease patients in a clinical trial.** (A) A manikin head demonstrating the drug delivery system used in the study. An application set coming from infusion pumps was attached to a skull-mounted transcutaneous port. Intracranially implanted catheters were also connected to the skull-mounted port. (B) The skull-mounted port was relatively imperceptible. It was the only external component when a patient was not receiving an infusion. (C) An axial MRI image at the level of the putamen shows two of catheters entering either side of the brain posteriorly and reaching the putamen. Gadolinium test infusate can be seen distributed throughout both putamen. (D) A single patient receiving GDNF infusion. (E) Three patients receiving their monthly GDNF infusions in a study infusion suite. Image reproduced from (Whone et al. 2019), under the terms of the Creative Commons Attribution 4.0 International.

## 2.6.2 Clinical studies with NRTN

Supported by the promising safety and efficacy outcomes in preclinical animal studies, intraparenchymal injections of AAV2 vector encoding human NRTN (CERE-120) have been evaluated in several clinical trials (Table 2.4.). In an open-label phase I study, AAV2-NRTN injected into the putamen showed evidence for safety and produced a 36% increase in the off-medication UPDRS motor score as compared to the baseline at 12 months (Marks et al. 2008). However, in a 12-month double-blind phase II trial, the intraputamenal AAV2-NRTN gene transfer failed to show superiority to sham surgery when assessed using the off-medication UPDRS motor score (Marks et al. 2010). Interestingly, a subgroup of patients, who were followed in a double-blind manner up to 18 months, revealed a small but significant clinical benefit in favor of AAV2-NRTN suggesting a possibility for a delayed effect. In addition, the gene therapy was well tolerated and improved several secondary outcome measures both at 12 and 18 months.

In contrast to the findings in MPTP-treated monkeys (Kordower et al. 2006; Herzog et al. 2007), brain autopsy data from four PD subjects who died from unrelated causes after intraputamenal administration of AAV2-NRTN showed impaired axonal transport of NRTN from the putamen to SNpc (Bartus et al. 2011; Bartus et al. 2015). To overcome this axonal transport deficiency, the next studies utilized a new dosing paradigm: AAV2-NRTN was injected into the SNpc in combination with putamenal delivery to activate repair genes directly in the cell bodies, and thereby enhance the potential trophic effects. Nonetheless, after encouraging safety outcomes in an open-label phase I study (Bartus et al. 2013), a sham-surgery controlled double-blind phase II trial failed to meet the primary endpoint assessed by the off-state UPDRS motor score at 15-month time point (Olanow et al. 2015). *Postmortem* analysis of two patients from these studies demonstrated that AAV2-NRTN delivery provided persistent, but spatially restricted, transgene expression in the putamen and SN still at 8 and 10 years after the transfection (Chu et al. 2020). NRTN expression was associated with focal sprouting and upregulation of TH expression, but not with antiparkinsonian effects such as reduced Lewy pathology when compared to untreated PD control subjects.

### **2.6.3 The first clinical study with CDNF**

The first-in-human, double-blind, placebo-controlled phase I/II study investigating the safety and efficacy of intraputamenal CDNF application in 17 patients with moderate PD was recently completed (Huttunen and Saarma 2019; ClinicalTrials.gov identifier NCT03295786). The study consisted of two parts: the initial 6-month main study in which patients were assigned to receive either placebo or two dose levels of CDNF, and the 6-month open-label extension part during which all patients received one of the two doses of CDNF. The company announced in a press release that the study achieved its primary endpoint of safety and tolerability at 12 months (Herantis Pharma Plc, press release 27 August 2020). Two patients had to discontinue the study due to adverse events related to the implanted dose delivery system. Treatment-related adverse events, however, were transient and mild. The preliminary non-statistical assessment of the secondary endpoints suggests an improvement in the UPDRS motor score in some patients at 12 months supporting the potential efficacy of CDNF therapy. The exploratory endpoints included DAT-PET imaging which showed encouraging response in the putamen of some CDNF-treated individuals. Noteworthy, the final analyses of the secondary and exploratory outcomes including CSF proteomics and  $\alpha$ -syn measurements are still underway.

### **3 AIMS OF THE STUDY**

The promising preclinical results of GDNF and NRTN have failed to translate into an effective disease-modifying treatment for PD. Therefore, to fulfill this unmet medical need, new approaches for neurotrophic therapies have to be explored. The studies were executed to support the preclinical characterization of unconventional NTFs, CDNF and MANF, and novel small molecule RET agonists. We aimed to uncover the functional effects of intracranially administered CDNF and MANF in the normal rat brain and to clarify the neuroprotective potential of RET agonists in a rat model of PD.

The specific objectives were to determine:

1. the effects of exogenously administered GDNF, CDNF and MANF on dopamine and GABA release and dopamine synthesizing and metabolizing enzymes in the normal rat brain (Study I)
2. the diffusion and transport properties of CDNF after injection to the substantia nigra of non-lesioned rats (Study II)
3. the neuroprotective and neurorestorative effects of small molecule RET agonists in a 6-OHDA model of PD in rats (Studies III and IV).

## 4 MATERIALS AND METHODS

The main methods used by the author in studies I-IV are shortly described here. More comprehensive descriptions of the materials and methods can be found in the original publications and their supplementary materials which are attached to the thesis.

### 4.1 Animals

Adult Wistar male rats (RccHan:WIST, Harlan, the Netherlands) were used for all experiments in study I and II, for the neuroprotection experiment in study III and for the neurorestoration experiment in study IV. Rats were normally housed in groups of 3-4 per cage under a 12h light/dark cycle (lights on 06:00-18:00). All experiments were performed during the light period. Rats had *ad libitum* access to rat chow and tap water at all times. In study I, rats were moved to individual microdialysis cages (30 × 30 × 30 cm) after the surgery. In studies III and IV, rats were housed in individual cages during the brain infusion of the treatments but regrouped again after removing the infusion cannulas and minipumps. The animal experiments were reviewed and approved by the National Animal Experiment Board of Finland (ESLH-2009-05234/Ym-23; ESAVI/5459/04.10.03/2011; ESAVI/7551/04.10.07/2013; ESAVI/ 198/04.10.07/2014 and ESAVI/7812/04.10.07/2015) and carried out in accordance with the European Union directive 2010/63/EU on the protection of animals used for scientific purposes.

### 4.2 Drugs, toxins and neurotrophic therapies

#### 4.2.1 Neurotrophic factors

The following NTFs were used: recombinant human (rh) GDNF (ProSpec-Tany TechnoGene Ltd., Israel) (study I and unpublished experiment), rhGDNF (PeproTech Inc., USA) (studies III and IV), rhCDNF (Biovian Oy, Finland) (studies I, II and unpublished experiment) and rhMANF (Icosagen AS, Estonia) (study I and unpublished experiment). NTFs were diluted in sterile phosphate-buffered saline, pH 7.4 (PBS) which also served as a vehicle control in studies I, IV and unpublished experiment.

#### 4.2.2 Small molecule RET agonists

BT13 (N,N-diethyl-3-(4-(4-fluoro-2-(trifluoromethyl)benzoyl)piperazin-1-yl)-4-methoxy-benzene-sulfonamide), MW 517.54 g/mol (study III), and BT44 ((4-5-((3,4-dihydroisoquinolin-2(1H)-yl)sulfonyl)-2-methoxy-phenyl)piperazin-1-yl(4-fluoro-2-(trifluoromethyl)phenyl)-methanone), MW 577.59 g/mol (study IV), were synthesized by EvoBlocks Ltd. (Hungary). The structure and integrity of the compounds were verified by NMR spectrometry, and the purity by liquid chromatography–mass spectrometry (BT13, 98.6%) or high-performance liquid chromatography (HPLC) (BT44, 97.3%). BT13 and BT44 were dissolved in 100% propylene glycol (PG) for the *in vivo* neuroprotection and -restoration experiments due to limited solubility in aqueous solutions

(studies III and IV). Their long-term stability in PG at 37°C was confirmed before the *in vivo* experiments. PG was used as a vehicle control in the studies.

#### 4.2.3 Neurotoxin

In studies III and IV, dopaminergic lesions were induced using intracranial injections of 6-OHDA (6-hydroxydopamine hydrochloride, Sigma-Aldrich GmbH, Germany), dissolved in ice-cold, de-oxygenated saline with 0.02% ascorbic acid. Desipramine hydrochloride 15 mg/kg, i.p. (calculated as free base; Sigma-Aldrich) was administered 30 min before the 6-OHDA injection to prevent the uptake of 6-OHDA into noradrenergic nerve terminals.

#### 4.2.4 Drugs

Isoflurane (Vetflurane® 1000 mg/g, Virbac SA, France) was administered via inhalation to induce and maintain general anesthesia during stereotaxic surgeries. Lidocaine-adrenalin solution (Lidocain 10 mg/ml c. adrenalin, Orion Pharma Oyj, Finland) was used for local anesthesia. To relieve postoperative pain, rats received tramadol 1 mg/kg, s.c. (Tramal® 50 mg/ml, Orion Pharma) at the end of the surgeries (study I), or buprenorphine 0.05 mg/kg, s.c. (Temgesic® 0.3 mg/ml, Indivior UK Limited, United Kingdom) before and carprofen 5 mg/kg, s.c. (Rimadyl Vet® 50mg/ml, Zoetis Inc., USA) immediately after the surgeries (studies II-IV). Additional doses of the analgesics were given one day later.

D-amphetamine sulfate (Division of Pharmaceutical Chemistry and Technology, University of Helsinki, Finland) was used to stimulate dopamine overflow from striatal nerve terminals in study I and induce turning behavior in unilaterally 6-OHDA lesioned rats in studies III and IV. In GABA microdialysis experiment (unpublished), nipecotic acid (Sigma-Aldrich) served as a universal inhibitor of GABA transporters (IC<sub>50</sub> ~18 µM). For *in vivo* TH activity experiment in study I, rats were administered with 3-hydroxybenzylhydrazine (NSD1015) (Sigma-Aldrich) to inhibit AADC in the brain. Clorgyline (IC<sub>50</sub> ~1.2 nM) and pargyline (IC<sub>50</sub> ~8.2 nM) (included in the Monoamine Oxidase Assay Kit, MAK136, Sigma-Aldrich) were used to irreversibly inhibit MAO-A and MAO-B activities, respectively, in MAO activity experiment in study I. Before transcatheter perfusions in studies II-IV, rats were deeply anesthetized with sodium pentobarbital (Mebunat Vet® 60 mg/ml, Orion Pharma).

### 4.3 Stereotaxic surgeries

All stereotaxic surgeries were performed under isoflurane anesthesia (3.5–4.5% during induction and 2.0–3.5% during maintenance) using a stereotaxic frame (Stoelting Co., USA). The skull was exposed and burr holes were made using a high-speed drill (Freedom Electric Co., USA). NTFs, RET agonists, vehicles and 6-OHDA were administered into the target brain regions as described below. Brain injections were made using an electronic microinjector (Quintessential stereotactic injector, Stoelting) and 10-µl syringes with 26G (Hamilton Co., USA) (studies I-III) or 33G (NanoFil, World Precision Instruments, USA) (study IV) blunt tapered needle attached. At the completion of the injections, the needle was kept in place for 4-5 min to minimize backflow of the solution. All

coordinates were determined relative to the bregma, according to the rat brain atlas (Paxinos and Watson 1998). After the surgeries, rats were allowed to recover at least one week before microdialysis or behavioral experiments.

#### **4.3.1 Neurotrophic factor injections and guide cannula implantation**

In study I, a unilateral injection of GDNF, CDNF or MANF (10 µg in 5 µl) or PBS (5 µl) was made into the left dorsal striatum (A/P +1.0, M/L +2.7, D/V -5.0 mm). The injection rate was 1.0 µl/min. For the enzyme activity assays, the incision was sutured after the injection. For the microdialysis experiments, a guide cannula (BASi MD-2250, Bioanalytical Systems Inc., USA) was implanted after the NTF injection. The tip of the cannula was placed into the left dorsal striatum (A/P +1.0, M/L +2.7, D/V -4.0 mm) for dopamine release measurements, or into the left GPe (A/P -1.4, M/L +3.5, D/V -5.5 mm) for GABA release measurements. Thereafter, the cannula was fixed to the skull with three stainless steel screws and polycarboxylate cement (Aqualox, Voco GmbH, Germany). In study II, CDNF (3 µg in 4 µl, or 3 µg in 1 µl) was injected into the left SN (A/P -5.4, M/L +2.0, D/V -7.2 mm) with the flow rate of 0.5 µl/min. The incision was sutured after the injection.

#### **4.3.2 6-OHDA injections**

In study III, rats received a single unilateral injection of 6-OHDA (16 µg in 4 µl; calculated as free base) into the left dorsal striatum (A/P +1.0, M/L +2.7, D/V -4.0 mm) at the flow rate of 1.0 µl/min. In study IV, 6-OHDA injections were made into three different sites in the right dorsal striatum (A/P +1.6, M/L -2.8, D/V -6.0 mm; A/P 0.0, M/L -4.1, D/V -5.5 mm; and A/P -1.2, M/L -4.5, D/V -5.5 mm) with the dose of 3 µg in 1.5 µl per site and flow rate of 0.5 µl/min (Penttinen et al. 2016). The needle was kept at a 10° angle to avoid lateral ventricles. Lastly, the scalp incision was closed with sutures.

#### **4.3.3 Infusions of RET agonists and GDNF with osmotic minipumps**


Osmotic infusion pumps (Alzet Osmotic pump model 2002, Durect Co., USA), connected with catheter tubing to brain cannulas (Alzet Brain infusion kit no. 2, Durect), were used to deliver RET agonists, GDNF and vehicles into the lesioned dorsal striatum in studies III and IV. The pumps were aseptically filled with study solutions and allowed to reach the steady-state pumping rate by incubating overnight in isotonic saline at 37°C. The pumps and cannulas were implanted in a stereotaxic surgery one hour (study III) or two weeks (study IV) after the 6-OHDA injections. The pump was placed into a subcutaneous pocket between the scapulae. The tip of the cannula was lowered to coordinates A/P +1.0, M/L +2.7, D/V -4.0 mm (study III) or A/P +0.2, M/L -3.0, D/V -5.0 mm (study IV) after which the cannula was secured to the skull with three stainless steel screws and polycarboxylate cement (Aqualox).

In study III, the pumps constantly delivered BT13 ~3-6 µg/24h (0.25-0.5 µg/µl), GDNF 3 µg/24h (0.25 µg/µl) or 100% PG into the dorsal striatum at the flow rate of 0.5 µl/h for seven days. The dose for BT13 could not be determined accurately because the solubility of BT13 in 100% PG was limited to 0.5 µg/µl in stable test tube conditions at 37°C. In study IV, the pumps infused BT44 0.1

$\mu\text{g}/24\text{h}$  (8.3 ng/ $\mu\text{l}$ ), BT44 0.3  $\mu\text{g}/24\text{h}$  (25 ng/ $\mu\text{l}$ ), GDNF 3  $\mu\text{g}/24\text{h}$  (0.25  $\mu\text{g}/\mu\text{l}$ ), 100% PG or PBS at the flow rate of 0.5  $\mu\text{l}/\text{h}$  for 14 days. The doses for BT44 were selected based on the potency of the compound in neuronal survival assay *in vitro* where it showed at least 10 times higher potency as compared to BT13. Therefore, 10-20 and 30-60-times lower doses of BT44 were chosen for study IV than were used for BT13 in study III. After the treatment infusions, the pumps, cannulas and screws were removed, and the incision was cleaned, disinfected and sutured.

## 4.4 Microdialysis experiments

Microdialysis experiments were carried out in freely-moving rats one and three weeks after the stereotaxic injection of NTFs and implantation of guide cannulas (study I and unpublished experiment). The one-week time point was selected based on an earlier brain microdialysis experiment with GDNF in freely-moving rats (Xu and Dluzen 2000). The three-week time point served to probe the long-term effects of NTFs on dopamine and GABA transmission. Proper function of the microdialysis probes (BASi MD-2200, Bioanalytical Systems, membrane length 2 mm) was verified with *in vitro* recovery test at room temperature before the experiments but *in vivo* dialysate concentrations were not corrected for *in vitro* recoveries. In each experiment, the probe was inserted into the guide cannula and perfusion of the microdialysis membrane was started with modified Ringer solution (147 mM NaCl, 2.7 mM KCl, 1.2 mM  $\text{CaCl}_2$ , 1.0 mM  $\text{MgCl}_2$ , 0.04 mM ascorbic acid) at the flow rate of 2.0  $\mu\text{l}/\text{min}$ . After a two-hour stabilization period, dialysate samples were collected every 15 min for 270 min (study I) or 150 min (unpublished experiment) as described below and in Figure 4.1. To minimize variation between the subjects, the results were analyzed as percent changes of the analyte concentrations from the baseline levels instead of actual analyte concentrations. In addition, the conventional microdialysis (as opposed to no-net-flux microdialysis) used here is not a suitable method to measure exact analyte concentrations in the extracellular fluid. After the first microdialysis experiment at week 1, the probe was removed from the brain. After the second experiment at week 3, rats were sacrificed and the brains were collected to verify the correct placements of the probes. Data only from the rats with accurate probe placements were included in the analyses.



**Figure 4.1. Time course of the dopamine and GABA microdialysis experiments.** The lag time between the administration of a stimulation solution and the corresponding response is due to slow flow rate of the perfusion solution (2.0  $\mu\text{l}/\text{min}$ ) and large dead volume of the long tubes. Figure drawn by the author.

In the dopamine microdialysis experiment (study I), dopamine, DOPAC and HVA concentrations in the first four samples were used to calculate the baselines. To depolarize the striatal nerve terminals and evoke dopamine release, 100 mM potassium solution (27.5 mM NaCl, 100 mM KCl,



1.2 mM CaCl<sub>2</sub>, 1.0 mM MgCl<sub>2</sub>, 0.04 mM ascorbic acid) was administered for 15 min via reverse dialysis starting at 15 min. To stimulate dopamine release for the second time by draining synaptic dopamine vesicles and reversing DAT function, 100 µM D-amphetamine (in modified Ringer) was administered via reverse dialysis for 15 min starting at 120 min. The analyte concentrations were quantified right after collecting the samples using a HPLC system with a C18 reverse-phase column (Kinetex C18, Phenomenex Inc., USA) and an electrochemical detector (Coulochem II, ESA Biosciences Inc., USA).

In the GABA microdialysis experiment (unpublished), reuptake of GABA into neurons and glial cells was blocked with 100 µM nipecotic acid which was included in the Ringer solution during the whole experiment (Vihavainen et al. 2008). The baseline level for GABA was calculated as an average in the first three samples. To stimulate GABA release, modified Ringer solution with 50 mM potassium (100 µM nipecotic acid, 99.7 mM NaCl, 50 mM KCl, 1.2 mM CaCl<sub>2</sub>, 1.0 mM MgCl<sub>2</sub>) was delivered for 15 min starting at 3 min (reverse dialysis). The samples were stored at -80°C until analyzing with a HPLC system equipped with a C18 reverse-phase column (Kinetex C18, Phenomenex) and a fluorescence detector (Jasco FP-1520, Jasco Co., United Kingdom) as described in detail elsewhere (Julku et al. 2016; Piepponen and Skujins 2001). To enable sensitive fluorometric detection of GABA, it was derivatized with a mixed *o*-phthaldialdehyde (Sigma-Aldrich) and 2-mercaptoethanol (Sigma-Aldrich) reagent before the samples were injected into the column.

## 4.5 Enzymes activity measurements

Seven days after the intrastriatal injections of NTFs or vehicle in study I, tissue samples were collected from the dorsal striatum for enzyme activity measurements. Rats were decapitated, and the brains were excised rapidly and rinsed in ice-cold saline. Bilateral striatal punches were collected from coronal slices with a 3-mm sample corer, snap frozen on dry ice and stored at -80°C. Later, the samples were weighed, homogenized and centrifuged.


For *in vivo* TH activity measurement, rats were administered with a BBB penetrating AADC inhibitor NSD1015 100 mg/kg, i.p., 30 min before tissue collection (Vihavainen et al. 2008). After sample preparation, the S1 supernatants filtered with Vivaspin® 500 filter concentrators (10,000 MWCO PES; Sartorius Stedim Biotech GmbH, Germany) were analyzed with a HPLC system equipped with a C18 reverse-phase column (Kinetex XD-C18, Phenomenex) and an electrochemical detector (ESA CoulArray Electrode Array, ESA Biosciences) to measure the amount of accumulated L-DOPA as ng/g wet weight of the tissue specimen (Valros et al. 2015).

For COMT, MAO-A and MAO-B activity assays, the total protein concentration was determined using bicinchoninic acid method (Pierce™ BCA Protein Assay Kit, ThermoFisher Scientific Inc., USA). Total COMT activity was measured as described in (Schendzielorz et al. 2012; Smith et al. 2014). The reaction products, vanillic and isovanillic acid, were analyzed with HPLC using an electrochemical detector (Coulochem II, ESA Biosciences). MAO-A and MAO-B activities were determined with Monoamine Oxidase Assay Kit (MAK136, Sigma-Aldrich). To differentiate MAO-A and MAO-B activities, isoform-specific inhibitors (5 µM of clorgyline for MAO-A and 5 µM of

pargyline for MAO-B) were used. The fluorescence of the samples was measured with a multi-well plate reader (Victor2, 1420 Multilabel Plate Reader, PerkinElmer Inc., USA) using wavelengths 530 nm and 590 nm for excitation and detection, respectively.

## 4.6 Behavioral tests

In studies III and IV, behavioral tests were conducted at defined timepoints as indicated in Figure 4.2. which shows the designs of the neuroprotection (A; study III) and neurorestoration (B; study IV) experiments. All behavioral assessments were carried out in a blinded manner.



**Figure 4.2. Experimental design for the neuroprotection and neurorestoration experiments.** The neuroprotective effect of BT13 (A) and neurorestorative effect of BT44 (B) were investigated in a unilateral 6-OHDA model of PD in rats. Amph I-III: amphetamine-induced (2.5 mg/kg, i.p.) rotational asymmetry tests, Cylinder I and II: cylinder tests. Figure drawn by the author.

### 4.6.1 Amphetamine-induced rotational behavior

Amphetamine-induced rotational asymmetry test was used to measure motor deficits arising from the unilateral 6-OHDA lesion of the nigrostriatal dopamine system. Rotational behavior was monitored in automated rotometer bowls (Med Associates Inc., USA) as described previously in (Ungerstedt and Arbuthnott 1970; Lindholm et al. 2007). After a 30-min habituation period, rats received a single injection of D-amphetamine 2.5mg/kg, i.p. (calculated as free base). The number of full (360°) uninterrupted clockwise and counterclockwise turns was recorded for 120 min. Net

ipsilateral turns to the lesion side were calculated by subtracting contralateral turns from ipsilateral turns.

#### 4.6.2 Cylinder test

In study IV, cylinder test was used to evaluate spontaneous limb-use asymmetry. Cylinder tests were carried out before the amphetamine-induced rotational asymmetry tests using a protocol described by (Schallert et al. 2000; Tillerson et al. 2001). Briefly, rats were placed in a transparent plexiglass cylinder (diameter ~20 cm, height ~30 cm) and their exploratory activity was recorded for 10 min with a video camera placed under the cylinder. The number of weight-shifting movements against the wall was scored individually for the ipsilateral (non-impaired) and contralateral (impaired) forepaws. A simultaneous placement of both forepaws on the wall as well as lateral exploration along the wall by alternating right and left forepaw placements were scored as “both paw contacts”. After a rear, the first weight-receiving contact to the ground was scored as a landing contact for the forepaw used for landing, or as a “both paw contact” if both paws contacted the ground simultaneously. Percentual limb-use asymmetry score was first calculated separately for wall exploration and landing using the following formula (Schallert and Woodlee 2005):

$$\frac{\text{ipsilateral contacts} + 0.5 \times \text{both paw contacts}}{\text{ipsilateral} + \text{contralateral} + \text{both paw contacts}} \times 100\%$$

Finally, the mean of these scores was used as an averaged asymmetry score. Scores >50% indicate greater reliance on the ipsilateral (non-impaired) forelimb and <50% on the contralateral (impaired) forelimb.

### 4.7 Perfusion and brain sectioning

After completing the microdialysis (study I) or behavioral experiments (studies III and IV), or 2-24 h after CDNF injections in the diffusion study (study II), rats were deeply anesthetized with sodium pentobarbital 90 mg/kg, i.p., and transcardially perfused with PBS and 4% paraformaldehyde (PFA) in PBS. The brains were removed, post-fixed in 4% PFA overnight and stored in 20% sucrose in PBS at 4°C until snap freezing in dry ice-cooled isopentane. The frozen brains were cut into 40 µm-thick coronal sections with a cryostat (Leica CM3050, Leica Biosystems, Germany). The sections were collected in a series of six sections and stored in cryoprotective buffer at -20°C. For the diffusion study (study II), some brains were embedded in paraffin and cut to 5 µm sagittal sections (M/L 1.4-3.4 mm), taking every 10<sup>th</sup> section. The paraffin sections were collected on microscope slides and stored at 4°C.

### 4.8 Immunohistochemistry

Immunohistochemical labelling was performed on free-floating coronal (studies II-IV) or paraffin-embedded sagittal (study II) sections using standard immunohistochemical procedures as described in detail in the original publications. The primary and secondary antibodies used in the experiments are listed in Table 4.1.

**Table 4.1. Primary and secondary antibodies used for immunohistochemical labelling.**

Primary antibodies				Secondary antibodies				Studies
Antibody	Cat#, Supplier	Reactivity	Concentration; Incubation	Antibody	Conjugation	Cat#, Supplier	Concentration; Incubation	
Rabbit polyclonal anti-hCDNF	300-100, Icosagen	Human, mouse	Free-floating: 1:500; +4°C o/n	Goat anti-rabbit	Biotin	BA-1000, Vector Laboratories Inc., USA	1:200; RT 1h	II
			Paraffin: 1:1000; +4°C o/n	Horse anti-rabbit	Biotin	BA-1100, Vector Laboratories	1:200; RT 1h	II
				Goat anti-rabbit	AlexaFluor488	A-11034, ThermoFischer Scientific	1:200; RT 2h	II
Mouse monoclonal anti-parvalbumin	MAB1572, Merck Millipore KGaA, Germany	Rat, mouse, rabbit, human etc.	Paraffin: 1:1000; +4°C o/n	Goat anti-mouse	AlexaFluor568	A-11004, ThermoFischer Scientific	1:200; RT 2h	II
Mouse monoclonal anti-TH	MAB318, Merck Millipore	Rat, mouse, human, zebrafish etc.	Free-floating: 1:2000; RT o/n	Horse anti-mouse	Biotin	BA-2001, Vector Laboratories	1:200; RT 2h	III
			Paraffin: 1:500; +4°C o/n	Goat anti-mouse	AlexaFluor568	A-11004, ThermoFischer Scientific	1:200; RT 2h	II
Rat monoclonal anti-DAT	MAB369, Merck Millipore	Rat, mouse, human, monkey	Free-floating: 1:2000; +4°C o/n	Rabbit anti-rat	Biotin	BA-4000, Vector Laboratories	1:200; RT 1h	III
Rabbit polyclonal anti-TH	AB152, Merck Millipore	Rat, mouse, human, fruit fly etc.	Free-floating: 1:2000; RT o/n	Protein A (binds to IgG:s of rabbit, mouse, human, guinea pig etc.)	Biotin	MP Biomedicals, USA (protein A), Sigma-Aldrich (N-hydroxysuccinimido-biotin), conjugated in-house	1:100; RT 1h	IV
Rabbit monoclonal anti-DAT	ab184451, Abcam Plc., United Kingdom	Rat, mouse	Free-floating: 1:500; +4°C o/n	Goat anti-rabbit	Biotin	BA-1000, Vector Laboratories	1:500; RT 1h	IV

RT, room temperature; o/n, overnight; TH, tyrosine hydroxylase; DAT, dopamine transporter; IgG, immunoglobulin G

## 4.9 Histological analyses

The striatum and midbrain sections were stained for TH and DAT to assess the density of dopaminergic fibers in the striatum and the number of dopamine neurons in the SNpc.

### 4.9.1 Fiber density in the striatum

The optical densities of TH-ir and DAT-ir fibers in the dorsal striatum were measured bilaterally from three different rostro-caudal levels through the striatum from each rat (studies III and IV). The analyses were performed under blinded conditions. First, digital images of the immunostained sections were acquired with an automated bright field microscope whole slide scanner Panoramic P250 Flash II (3DHitech Ltd., Hungary). The images were converted to 8-bit gray scale, and colors were inverted. The dorsal striata were outlined, and integrated optical densities divided by the size of the outlined areas were measured with Fiji ImageJ software (Media Cybernetics Inc., USA). All density values were corrected for nonspecific background staining measured from the corpus callosum. The data are expressed as percentage of the fiber density on the lesioned hemisphere as compared to the intact hemisphere. The total magnitude of striatal denervation was estimated as an average reduction in the fiber density at the three levels measured.

### 4.9.2 Cell counts in the SNpc

In studies III and IV, the number TH-ir cells in the SNpc was estimated in a blinded manner with the optical fractionator method and dissector principle according to unbiased counting rules (West et al. 1991). Olympus BX51 microscope (Olympus Co., Japan) was connected to a computer running Stereo Investigator platform (MBF Bioscience Inc., USA), and cell counting was done bilaterally under 60x magnification (Olympus PlanApo 60x 1.40 Oil /0.17 objective) from three sections per brain at approximately the same rostro-caudal levels as described earlier (Voutilainen et al. 2009). The results are presented as percentage of cells in the lesioned SNpc as compared to the intact SNpc.

Additionally, in study IV, TH-ir cells bodies in the SNpc were counted using an automated deep convolutional neural networks (CNN) algorithm in a cloud-embedded image processing platform (Aiforia Technologies Oy, Finland). This computer-assisted cell counting method, that is based on supervised machine learning and automated image recognition, is described in detail and validated in (Penttinen et al. 2018). Briefly, digitized images of TH-immunolabelled sections were uploaded to Aiforia™ platform. The SNpc was demarcated bilaterally in six sections from each brain. CNN algorithm, that was trained to recognize dopaminergic cell bodies from digital images, counted the number of TH-ir neurons within the demarcated areas. The number of neurons was summed up from the six sections separately for both hemispheres. The performance of the algorithm was confirmed against stereological assessment of TH-ir cell counts in 20 randomly selected brains in study IV. The results obtained with CNN algorithm and stereological counting method had a strong positive correlation (Pearson's  $r = 0.83$ ,  $p < 0.001$ ).

## 4.10 Statistical analyses

All results are presented as mean  $\pm$  standard error of the mean (SEM). The statistical analyses were conducted with SPSS® Statistics 24 software (IBM SPSS Inc., USA). The group sizes and NTF doses were selected on the basis of our earlier experience (Airavaara et al. 2006; Käenmäki et al. 2010; Lindholm et al. 2007; Voutilainen et al. 2011, 2009). Paired two-tailed Student's *t*-test was used for two-sample within-subject comparisons and unpaired two-tailed Student's *t*-tests for two-sample between-subject comparisons. Group-wise analyses were done using one-way analysis of variances (ANOVA) in case of between-group comparisons, or repeated measures ANOVA in case of within-group comparisons, followed by Ryan-Einot-Gabriel-Welsch *F* (REGWF; study I) or Tukey HSD (studies II-IV) *post hoc* tests. To assess the strength of a linear association between two variables, Pearson correlation test was applied. The results were considered to be statistically significant at  $p < 0.05$ .


## 5 RESULTS

### 5.1 Injection of neurotrophic factors into the intact rat brain

#### 5.1.1 Effects of exogenous neurotrophic factors on dopamine release and turnover in the striatum (I)

In the microdialysis experiment one week after the intrastriatal NTF injections, the baseline levels of extracellular dopamine, DOPAC or HVA did not differ between the treatment groups. Both potassium and amphetamine-evoked release of dopamine was significantly elevated in MANF-injected rats as compared to vehicle or GDNF-injected rats (Figure 5.1.A and B). There was also a trend for increased stimulus-evoked dopamine release in CDNF-injected rats, but this effect did not reach statistical significance when compared to vehicle. At three weeks post injection, there were no more significant differences in dopamine release between the treatment groups (Study I, supplementary material).


Reflecting enhanced dopamine metabolism, MANF induced a significant increase in dopamine turnover in the striatum as evaluated by calculating DOPAC/dopamine ratio in the microdialysis samples at one week, but not anymore at three weeks post injection (Figure 5.1.C). There were no significant differences in HVA/dopamine ratio between the treatment groups at one or three weeks after the injection.



**Figure 5.1. Striatal dopamine release and turnover measured with brain microdialysis one week after an intrastriatal injection of neurotrophic factors.** (A) Potassium- and amphetamine-evoked (75–135 min and 180–255 min, respectively) dopamine release was increased in MANF group as compared to vehicle and GDNF groups (\* $p < 0.05$ ; REGWF after repeated measures ANOVA; 75–135 min:  $F_{3,36} = 4.874$ ,  $p = 0.006$ ; 180–255 min:  $F_{3,36} = 3.683$ ,  $p = 0.021$ ). In addition, the overall (15–270 min) extracellular concentration of dopamine was elevated in MANF group as compared to vehicle and GDNF groups (\* $p < 0.05$ ) and in CDNF group as compared to GDNF group (# $p < 0.05$ ; REGWF after repeated measures ANOVA; 15–270 min:  $F_{3,36} = 4.678$ ,  $p = 0.007$ ). (B) Averaged total potassium- and amphetamine-evoked dopamine overflow was augmented in MANF group when compared to vehicle and GDNF groups (\* $p < 0.05$ ; REGWF after one-way ANOVA; K<sup>+</sup> response:  $F_{3,36} = 4.874$ ,  $p = 0.006$ ; Amphetamine response:  $F_{3,36} = 3.683$ ,  $p = 0.021$ ). (C) DOPAC/dopamine ratio was higher in MANF-injected rats as compared to vehicle and GDNF-injected rats during the whole experiment and in the baseline samples (\* $p < 0.05$ ; REGWF after repeated measures ANOVA; 15–270 min:  $F_{3,36} = 3.065$ ,  $p = 0.040$ ; 15–60 min:  $F_{3,36} = 3.868$ ,  $p = 0.017$ ). The dose of each NTF was 10  $\mu$ g. Results are shown as % of the baseline (A and B) or concentration ratio (C); mean  $\pm$  SEM;  $n = 10$ . Figures adapted from (Renko et al. 2018), under the terms of the Creative Commons Attribution 4.0 International.

### 5.1.2 Effects of exogenous neurotrophic factors on dopamine neurochemistry (I)

The amount of L-DOPA, the precursor of dopamine, accumulated into striatal tissue samples after AADC inhibition by NSD1015 provides a direct measure for L-DOPA production rate indicating the *in vivo* TH activity of the nigrostriatal pathway (Carlsson et al. 1972). At one week post NTF injection, TH activity was increased approximately by 60% in the dorsal striatum of rats administered with GDNF as compared with the vehicle-administered controls (Figure 5.2.A). MANF also tended to increase TH activity (approximately by 50%), but the effect was not statistically significant.



**Figure 5.2. Effects of neurotrophic factors on dopamine neurochemistry-regulating enzymes one week after an intrastriatal injection.** (A) *In vivo* TH activity was increased in GDNF-injected rats when compared with vehicle-injected rats as analyzed by measuring the amount of accumulated L-DOPA in the striatal samples after AADC inhibition with NSD1015 (\* $p < 0.05$ ; REGWF after one-way ANOVA;  $F_{3,22} = 3.780$ ,  $p = 0.025$ ). (B) DOPAC/HVA ratio was significantly smaller in GDNF group than in the other groups during the whole experiment (# $p < 0.05$ ; REGWF after repeated measures ANOVA; 15–270 min:  $F_{3,36} = 7.397$ ,  $p = 0.001$ ). (C) GDNF injection increased total COMT activity in the striatum when compared to the vehicle-injected striatum (\*\* $p < 0.001$ ; unpaired two-tailed Student's *t*-test  $t(13) = -5.159$ ) as well as compared to the non-injected striatum (# $p < 0.001$ ; paired two-tailed Student's *t*-test  $t(7) = 6.041$ ). (D) MAO-A activity in the GDNF-injected striatum was reduced as compared to the vehicle-injected striatum (\* $p = 0.011$ ; unpaired two-tailed Student's *t*-test  $t(14) = 2.944$ ). The dose of the NTFs was 10  $\mu$ g. Mean  $\pm$  SEM;  $n = 6-7$  (A),  $n = 10$  (B),  $n = 7-8$  (C, D). Figures adapted from (Renko et al. 2018), under the terms of the Creative Commons Attribution 4.0 International.


Interestingly, the concentration of dopamine metabolites in the striatal microdialysis samples collected at one week post NTF injections showed that DOPAC/HVA ratio was significantly reduced in rats injected with GDNF as compared to all other groups (Figure 5.2.B). This suggests that GDNF injection modulates the activity of dopamine metabolizing enzymes. Indeed, GDNF significantly increased the total COMT activity by 155% in the injected striatum as compared to vehicle-injected striatum and by 170% when compared to the non-injected striatum (Figure 5.2.C). GDNF also significantly reduced the activity of MAO-A, the isoform primarily responsible for the oxidation of



dopamine in rats (Meiser et al. 2013), in the injected striatum by 27% as compared to the vehicle-injected controls (Figure 5.2.D).

### 5.1.3 Effects of exogenous neurotrophic factors on GABA release in the GPe (unpublished data)

To test whether intrastrially injected NTFs modulate the GABAergic neurotransmission in the striatopallidal or striatonigral projection pathways, we collected microdialysis samples from the GPe and SNr of freely-moving rats at one and three weeks after the NTF injections. To stimulate GABA release, the MSN nerve terminals were depolarized by administering high concentration of potassium through the microdialysis probe. According to the pilot experiment, the overall extracellular concentration of GABA did not differ significantly between the treatment groups at one or three weeks post injection, nor were there any significant effects on stimulus-evoked GABA output (Figure 5.3.). At three weeks post injection, there seemed to be a tendency toward increased potassium-evoked release of GABA in CDNF-injected animals, but this difference did not reach statistical significance. We also measured the amount of GABA in the GPe tissue samples collected one week after the intrastriatal NTF injections, but no significant differences were found between the groups. GABA release in the SNr could not be reliably quantified because the concentration of GABA in the dialysates was at the limit of quantitation of the HPLC system and sometimes even below it (we used 1 mm long membranes for the SNr microdialysis).




**Figure 5.3. GABA release in the GPe measured with brain microdialysis one and three weeks after an intrastriatal injection of neurotrophic factors.** There were no significant differences in GABA release between the treatment groups at one week (A) or three weeks (B) post injection. The dose of each NTF was 10  $\mu$ g. Results are shown as % of the baseline; mean  $\pm$  SEM;  $n = 4-6$ .

### 5.1.4 Distribution of CDNF after nigral injection (II)


To characterize the diffusion properties of supra-nigraly injected CDNF (3  $\mu$ g) in naïve rats, the brains were collected for immunohistochemical analysis at 2, 6 and 24 h after the injection. We observed robust CDNF staining throughout the ipsilateral midbrain including the SNpc at 2 and 6 h post injection, but not anymore at the 24-h time point (Figure 5.4.A-C). Clear CDNF labelling was also detected in the hippocampus and amygdala as well as thalamic and hypothalamic regions. There was no CDNF signal observed in the striatum at any of the time points. Additionally, increasing the injection volume from 1 to 4  $\mu$ l while keeping all other parameters, such as the amount of CDNF, constant resulted in larger diffusion volume of CDNF around the SNpc at the 2-

h time point as shown in the representative sagittal images of the CDNF staining (Figure 5.4.D). These qualitative data suggest that CDNF readily diffuses to the brain areas surrounding the SN but is eliminated within 24 h after the nigral injection.




**Figure 5.4. Diffusion of human CDNF (hCDNF) after an injection to the substantia nigra (SN) of intact rat brain.** (A-C) Representative hCDNF immunostaining on coronal sections collected 2, 6 or 24 h after a nigral hCDNF (3 µg/4 µl) injection. CDNF readily diffuses in the brainstem, in particular around the SN, to the hippocampus, amygdala, thalamus and hypothalamus within 2 to 6 h after the injection. The panel on the right shows the rostro-caudal levels of the stained sections. (D) Representative hCDNF immunolabelling on sagittal sections collected 2 h after a nigral hCDNF injection in the concentration of 3 µg/1 µl or 3 µg/4 µl. CDNF diffuses farther with the 4-µl injection volume. The arrowheads indicate the approximate injection sites (added afterwards to the original figure); n = 2 at each time point or volume. Figures adapted from (Albert et al. 2019), under the terms of the Creative Commons Attribution 4.0 International.

To further elucidate the localization of CDNF after the nigral injection, the midbrain sections were double stained for CDNF and TH or PV. We observed colocalization of CDNF with TH-ir neurons in the SN, while the colocalization with PV-ir neurons was not evident. Multiple TH-ir cells were detected in the nigral area of each brain, and approximately two thirds of them had CDNF-ir puncta inside their cell body on the injected side (Figure 5.5.). Although these observations are only qualitative, they are in good accordance with the earlier results by (Mätlik et al. 2017).



**Figure 5.5. Immunofluorescent double staining for human CDNF (hCDNF) and tyrosine hydroxylase (TH) in the substantia nigra (SN).** Representative confocal images from a rat brain injected with hCDNF (3  $\mu$ g) to the SN and perfused 2 h later (right panel). Control side images (left panel) are taken with identical settings to account for possible background staining. White arrows on the injected side indicate hCDNF-ir puncta (cyan dots) within TH-ir cells (magenta) demonstrating the colocalization of hCDNF with dopamine neurons. Scale bars 10  $\mu$ m;  $n$  = 6. Figures adapted from (Albert et al. 2019), under the terms of the Creative Commons Attribution 4.0 International.

When  $^{125}$ I-labeled CDNF was injected to the SN and several brain areas were dissected 24 h later, no radioactivity was detected in distant brain areas such as the striatum, globus pallidus or cortex suggesting the absence of axonal transport to these areas (Figure 5.6.). The highest radioactivity after the nigral injection of  $^{125}$ I-CDNF was measured in the ipsilateral STN. Concurrent administration of either 2000 or 10 000-fold molar excess of unlabeled (“cold”) CDNF to the SN together with  $^{125}$ I-CDNF to compete for the active transport mechanisms did not block the spread of  $^{125}$ I-CDNF to the STN.



**Figure 5.6. Spread of  $^{125}$ I-labeled CDNF after an injection to the substantia nigra (SN) of intact rat brain.** Percentage of the total radioactivity (=summed radioactivity in all brain areas measured) in the ipsilateral (left panel) and contralateral (right panel) brain areas measured from tissue samples 24 h after the nigral injection of  $^{125}$ I-CDNF (5 ng). For a subset of animals, 2000 (10  $\mu$ g) or 10 000-fold (50  $\mu$ g) molar excess of cold (unlabeled) CDNF was injected together with  $^{125}$ I-CDNF. No transport into the striatum or other brain areas was detected.  $^{125}$ I-CDNF spread into the ipsilateral STN, but this spread was not affected by the excess of cold CDNF. FCTX, frontal cortex; CTX, cortex; STR, striatum; HC, hippocampus; GP, globus pallidus; STN, subthalamic nucleus; mean  $\pm$  SEM;  $n$  = 5-6 per brain region. Figures adapted from (Albert et al. 2019), under the terms of the Creative Commons Attribution 4.0 International.

## 5.2 Activation of neuronal pro-survival signaling pathways by RET agonists

To facilitate the comparison of pharmacological properties between RET agonists and positive controls (GDNF and NRTN), the molar concentrations of the ligands used in different *in vitro* experiments are provided in Table 5.1. From this comparison it is obvious that the potencies of GDNF and NRTN are at the level of four orders of magnitude larger than those of BT-compounds in the phosphorylation assays, which reflect the increase in protein phosphorylation status after a short-term (15 min) exposure to the ligands. In the neuronal survival and protection assays in cultured dopamine neurons, the potency differences are at the level of one to three orders of magnitude, because the integrated effect of prolonged (5 days) RET activation allowed to use much lower ligand concentrations. Direct potency and efficacy comparisons based on these experiments, however, must be done with caution due to putative differences in experimental conditions, such as transfection efficiencies of GFR $\alpha$  co-receptors, which may have an impact on the intensity readouts.

**Table 5.1. Comparison of molar concentrations used in *in vitro* experiments in Studies III and IV.**


Assay	Ligand	Concentration	Effect
RET phosphorylation in MG87RET fibroblasts	BT13	25-100 $\mu$ M	+
	BT44	18-75 $\mu$ M	+
	GDNF	6.6 nM	+
	NRTN	4.2 nM	+
Akt phosphorylation in MG87RET fibroblasts	BT13	25-100 $\mu$ M	?
	BT44	36-75 $\mu$ M	+
	GDNF	6.6 nM	+
	NRTN	4.2 nM	+
ERK phosphorylation in MG87RET fibroblasts	BT13	25-50 $\mu$ M	+
	BT44	7.5-75 $\mu$ M	+
	GDNF	6.6 nM	+
	NRTN	4.2 nM	+
Survival of cultured dopamine neurons (wt)	BT13	0.1-1 $\mu$ M	+
	BT44	7.5 nM - 3.5 $\mu$ M	+
	GDNF	0.33 nM	+
Survival of cultured dopamine neurons (RET knockout)	BT13	1-5 $\mu$ M	–
	BT44	7.5-75 nM	–
	GDNF	0.33 nM	–
Protection of cultured dopamine neurons against MPP <sup>+</sup> induced cytotoxicity (wt)	BT13	1 $\mu$ M	+
	BT44	75 nM	+
	GDNF	0.33 nM	+
Protection of cultured dopamine neurons against MPP <sup>+</sup> induced cytotoxicity (RET knockout)	BT13	0.1-1 $\mu$ M	–
	GDNF	0.33 nM	–

+, significant increasing effect compared to vehicle; –, no significant effects compared to vehicle; ?, no quantitative data but seems to increase phosphorylation compared to vehicle; wt, wild-type neurons

### 5.2.1 RET agonists phosphorylate RET and induce intracellular signaling cascades (III and IV)


A novel small molecule RET agonist BT13 stimulated the phosphorylation of RET in immortalized MG87RET fibroblasts transfected with GFR $\alpha$ 1 or GFR $\alpha$ 2 co-receptors as demonstrated by representative Western blot images (Figure 5.7.A and B). The effect of BT13 was concentration dependent and did not require the presence of GFR $\alpha$  co-receptors since RET phosphorylation was also elicited in GFP-transfected MG87RET fibroblasts incubated with BT13 (Figure 5.7.C). This suggests that BT13 functions as a direct RET agonist.

When the downstream effects of BT13 were investigated, we saw that RET phosphorylation activated the intracellular targets Akt and ERK which are imperative for neuronal survival and neurite outgrowth (Figure 5.7.D-F). Akt and ERK phosphorylation also seemed to be dependent on the concentration of BT13 but independent on the expression of GFR $\alpha$ 1/2.




**Figure 5.7. BT13 induces phosphorylation of RET and activates its downstream targets Akt and ERK in MG87RET fibroblasts.** Representative Western blots show the increased level of RET phosphorylation in GFR $\alpha$ 1 (A), GFR $\alpha$ 2 (B) and GFP-transfected (C) cells in response to BT13 in a concentration-dependent manner. BT13 also increased the phosphorylation of Akt and ERK in a concentration-dependent manner in cells transfected with GFR $\alpha$ 1 (D), GFR $\alpha$ 2 (E) and GFP (F). GDNF, NTRN and soluble GFR $\alpha$ 1/GDNF complex produced expected responses when used as positive controls for GFR $\alpha$ 1, GFR $\alpha$ 2 and GFP transfected cells, respectively. Their concentrations are provided in ng/ml. IP, immunoprecipitation; WB, Western blotting; GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

In line with the *in vitro* data, BT13 (750  $\mu$ g) increased the level of phosphorylated ERK (pERK) and ribosomal protein S6 (pS6; downstream target of Akt) when injected into the mouse striatum. pERK and pS6 immunolabelling was significantly enhanced in the BT13-injected striata as compared to the vehicle-injected striata (Figure 5.8.). GDNF (5 or 10  $\mu$ g) was used as a positive control and produced a significant increase in pERK and pS6 staining in the striatum.



**Figure 5.8. BT13 activates intracellular survival-promoting signaling pathways in the mouse brain.**

(A) Representative images of pERK and pS6 immunostained striatal sections collected after an intrastriatal injection of GDNF (10 μg), BT13 (750 μg) or vehicle. (B) Relative optical density of pERK labeling in the dorsal striatum as compared to the vehicle-injected side. BT13 (750 μg) increased the level of pERK (\* $p = 0.048$ ;  $t(2) = 4.38$ ). GDNF elevated the level of pERK at the dose of 5 μg (\* $p = 0.027$ ;  $t(2) = 6.00$ ) and 10 μg (\*\* $p = 0.009$ ;  $t(3) = 6.19$ ). (C) Relative optical density of pS6 labeling in the dorsal striatum as compared to the vehicle-injected side. BT13 (750 μg) increased the level of pS6 (\* $p = 0.040$ ;  $t(3) = 3.47$ ). GDNF elevated the level of pS6 at the dose of 5 μg (\*\* $p = 0.004$ ;  $t(2) = 15.23$ ). Significant differences between the vehicle and BT13/GDNF-injected sides were determined with paired two-tailed Student's  $t$ -test. Scale bar 200 μm;  $n = 3-4$ .




**Figure 5.9. BT44 induces phosphorylation of RET and activates its downstream targets Akt and ERK in MG87RET fibroblasts.** Representative Western blots show the increased level of RET phosphorylation in GFRα1 (A), GFRα2 (B) and GFP-transfected (C) cells in response to BT44 in a concentration-dependent manner. BT44 also increased the phosphorylation of Akt and ERK in a concentration-dependent manner in cells transfected with GFRα1 (D) and GFRα2 (E). In GFP-transfected cells, BT44 seemed to phosphorylate only ERK but not Akt (F). GDNF and NRTN (ng/ml) served as positive controls in cells transfected with GFRα1 and GFRα2, respectively. IP, immunoprecipitation; WB, Western blotting; GAPDH, glyceraldehyde-3-phosphate dehydrogenase.



Analogously to the parental compound BT13, BT44 activated RET and its downstream target pathways in MG87RET fibroblasts. BT44 stimulated RET phosphorylation in cells transfected with the co-receptors GFR $\alpha$ 1 and GFR $\alpha$ 2 as well as in GFP-transfected cells suggesting a direct, co-receptor independent, agonistic mechanism (Figure 5.9.A-C). BT44 also induced the phosphorylation of Akt and ERK in GFR $\alpha$ 1-RET and GFR $\alpha$ 2-RET expressing fibroblasts (Figure 5.9.D and E). Interestingly, in GFP-transfected cells, BT44 induced ERK phosphorylation but did not have a clear effect on Akt phosphorylation (Figure 5.9.F).

## 5.2.2 RET agonists promote the survival of cultured midbrain dopamine neurons (III and IV)

Both BT13 and BT44 supported the survival of primary dopamine neurons *in vitro* as compared to vehicle. The number of TH-ir cells was increased in BT13 (0.1 and 1  $\mu$ M), BT44 (7.5, 75 and 3500 nM) and GDNF (10 ng/ml  $\approx$  0.33 nM) -treated midbrain cultures from wild-type, but not from RET knockout, mice indicating that the survival-promoting effect of BT13 and BT44 on dopamine neurons was RET dependent (Figure 5.10.A-D). BT13 (1  $\mu$ M) (Mahato et al. 2020) and BT44 (75 nM) also protected MPP $^{+}$  -challenged dopamine neurons in culture with a similar efficacy as GDNF (10 ng/ml  $\approx$  0.33 nM) (Figure 5.10.E).



**Figure 5.10. BT13 and BT44 promote the survival of cultured dopamine neurons from wild-type, but not from RET knockout, mice and protect them against MPP $^{+}$  -induced cell death.** (A) The number of TH-ir cells in wild-type midbrain cultures on 5<sup>th</sup> day *in vitro* (DIV) presented as percentage of the vehicle-treated samples. BT13 0.1  $\mu$ M ( $^{***}$ p < 0.0001), BT13 1  $\mu$ M ( $^{***}$ p = 0.0002) and GDNF 10 ng/ml ( $^{*}$ p = 0.021; Dunnett after repeated measures ANOVA;  $F_{2,35,16.46} = 6.55$ , p = 0.006) increased the number of TH-ir neurons in the culture. N = 8. (B) The number of TH-ir cells in RET knockout midbrain cultures on 5<sup>th</sup> DIV presented as percentage of the vehicle-treated samples. N = 8. (C) The number of TH-ir cells in wild-type midbrain cultures on 5<sup>th</sup> DIV. BT44 7.5 nM ( $^{***}$ p = 0.0008), BT44 75 nM ( $^{**}$ p = 0.006), BT44 3500 nM ( $^{**}$ p = 0.004) and GDNF 10 ng/ml ( $^{***}$ p = 0.0009; Dunnett after repeated measures ANOVA;  $F_{6,18} = 5.734$ , p = 0.0018) increased the number of TH-ir neurons as compared to the vehicle. N = 4. (D) The number of TH-ir cells in RET knockout midbrain cultures on 5<sup>th</sup> DIV. N = 2. (E) BT44 75 nM ( $^{*}$ p = 0.016) and GDNF 10 ng/ml ( $^{*}$ p = 0.019; Dunnett after repeated measures ANOVA;  $F_{3,15} = 5.410$ , p = 0.01) increased the number of surviving TH-ir cells in wild-type midbrain cultures exposed to MPP $^{+}$  neurotoxin. N = 6. Mean  $\pm$  SEM; the number of TH-ir cells is normalized to the total number of cells in the culture.


### 5.3 Delivery of RET agonists into the striatum of hemiparkinsonian rats

#### 5.3.1 BT13 protects against motor dysfunction in 6-OHDA model of Parkinson's disease (III)

The neuroprotective effect of intrastriatally infused BT13 was compared with that of GDNF in a unilateral 6-OHDA model of PD in rats. The experimental design is presented in Figure 4.2.A.

##### 5.3.1.1 Motor deficits

Amphetamine-induced rotational behavior was similar in all treatment groups at two weeks following the toxin administration (Figure 5.11.A). In animals infused with BT13 (3-6  $\mu$ g/24h) and GDNF (3  $\mu$ g/24h), the number of ipsilateral turns was significantly reduced at four and six weeks after the lesion as compared to the vehicle infusion (Figure 5.11.B and C). Noteworthy, there was spontaneous recovery in the turning behavior of vehicle-treated rats at four ( $p = 0.085$  for 2 weeks vs. 4 weeks:  $t(11) = 1.895$ ) and six weeks post lesion ( $p = 0.034$  and  $p = 0.020$  for 2 weeks vs. 6 weeks:  $t(10) = 2.460$  and 4 weeks vs. 6 weeks:  $t(10) = 2.756$ , respectively; paired two-tailed Student's  $t$ -test), an observation that is accordance with previously published data on this particular toxin administration scheme (Penttinen et al. 2016).




**Figure 5.11. BT13 normalizes amphetamine-induced turning behavior in a time-dependent manner in 6-OHDA lesioned rats.** (A) Effect of BT13 and GDNF infusions on amphetamine-induced turning behavior at two weeks post lesion. (B) At four weeks post lesion, the number of turns was lower in BT13 3-6 $\mu$ g/24h (\* $p = 0.013$ ) and GDNF 3 $\mu$ g/24h (\*\* $p = 0.001$ ) groups as compared to the vehicle group (Tukey HSD after one-way ANOVA;  $F_{2,53} = 7.344$ ,  $p = 0.002$ ). (C) The turning behavior was further reduced in BT13 3-6 $\mu$ g/24h (\* $p = 0.011$ ) and GDNF 3 $\mu$ g/24h (\*\* $p = 0.007$ ) groups as compared to the vehicle group at six weeks post lesion (Tukey HSD after one-way ANOVA;  $F_{2,51} = 5.882$ ;  $p = 0.005$ ). Mean  $\pm$  SEM; vehicle  $n = 11-12$ , GDNF 3 $\mu$ g/24h  $n = 15-16$ , BT13 3-6 $\mu$ g/24h  $n = 25-27$ .

##### 5.3.1.2 Dopaminergic neurons in the SNpc and fibers in the striatum

The immunohistochemical assessment of nigral and striatal sections collected after the last behavioral test revealed that GDNF 3  $\mu$ g/24h infusion partially protected TH-ir fibers in the lesioned striatum as compared to the vehicle treatment (Figure 5.12.A and D). BT13 3-6  $\mu$ g/24h, however, failed to produce a significant protective effect on dopaminergic fibers. Neither BT13 nor GDNF were able to protect TH-ir cells in the SNpc (Figure 5.12.B and E). Since GDNF delivery



into the striatum can downregulate the expression of TH (Georgievska et al. 2004; Salvatore et al. 2004), we used DAT as another dopaminergic marker to analyze the fiber density in the striatum. The density of DAT-ir fibers was comparable to that of TH-ir fibers in all treatment groups suggesting unaltered TH expression in the striatum (Figure 5.12.C and F).




**Figure 5.12. Effect of BT13 and GDNF on dopaminergic fibers in the striatum and cell bodies in the SNpc of 6-OHDA lesioned rats.** Representative images of TH-ir fiber density in the striatum (A), TH-ir neurons in the SNpc (B) and DAT-ir fiber density in the striatum (C) in different treatment groups at six weeks post lesion. The dashed line in (A) shows the area in the dorsal striatum used for optical density measurements and in (B) the area used for stereological cell counting. The arrowheads denote the lesion side. (D) Quantification of TH-ir fiber density in the striatum. GDNF protected TH-ir fibers in the lesioned striatum as compared to the vehicle treatment (\* $p = 0.024$ ; Tukey HSD after one-way ANOVA  $F_{2,51} = 3.708$ ,  $p = 0.032$ ). (E) Number of TH-ir cell bodies in the SNpc. (F) Densitometric quantification of DAT-ir fibers in the striatum. Scale bars: 1 mm (A and C) and 0.5 mm (B). Mean  $\pm$  SEM; in TH staining (D and E) vehicle  $n = 11-12$ , GDNF  $3\mu\text{g}/24\text{h}$   $n = 15-16$ , BT13  $3-6\mu\text{g}/24\text{h}$   $n = 24-25$ ; in DAT staining (F) vehicle  $n = 8$ , GDNF  $3\mu\text{g}/24\text{h}$   $n = 13$ , BT13  $3-6\mu\text{g}/24\text{h}$   $n = 19$ .

### 5.3.2 BT44 shows neurorestorative potential in 6-OHDA model of Parkinson's disease (IV)

The neurorestorative effect of intrastratially infused BT44 was studied in a rat model of PD with progressive unilateral 6-OHDA lesion (Penttinen et al. 2016) utilizing amphetamine-induced rotational behavior and limb-use asymmetry tests. The experimental design is presented in Figure 4.2.B. Rats were assigned into equal treatment groups according to their amphetamine-induced rotation rate at 12 days after the 6-OHDA injections (Figure 5.13.A).

### 5.3.2.1 Motor deficits

Both GDNF 3  $\mu\text{g}/24\text{h}$  and BT44 0.3  $\mu\text{g}/24\text{h}$  were able to produce a functional recovery in 6-OHDA lesioned animals. Amphetamine-induced rotational asymmetry was significantly reduced in GDNF-infused rats at six and 12 weeks post lesion and in BT44 0.3  $\mu\text{g}/24\text{h}$  -infused rats at 12 weeks post lesion as compared to the vehicle-treated rats (Figure 5.13.B and C). Spontaneous limb-use asymmetry in the cylinder test, however, was not balanced by any of the treatment infusions (Figure 5.13.E and F). This is in line with earlier literature where GDNF therapy has been reported to induce functional recovery in amphetamine-induced rotation test but not in the cylinder test (Georgievska et al. 2002; Gasmi et al. 2007b; Yue et al. 2014).




**Figure 5.13. BT44 reduces amphetamine-induced rotations in 6-OHDA lesioned rats in a time-dependent manner.** (A) Amphetamine-induced rotation rate at two weeks (12 days) post lesion was used to assigned rats into equal treatment groups. (B) At six weeks post lesion, GDNF 3  $\mu\text{g}/24\text{h}$  alleviated amphetamine-induced rotational asymmetry as compared to PBS (# $p = 0.025$ ) and PG (\* $p = 0.046$ ; Tukey HSD after one-way ANOVA;  $F_{4,45} = 3.638$ ,  $p = 0.012$ ). (C) At 12 weeks, BT44 0.3  $\mu\text{g}/24\text{h}$  significantly reduced the number of ipsilateral turns as compared to PBS (# $p = 0.046$ ) and PG (\* $p = 0.010$ ). GDNF-treated rats rotated significantly less than PBS (#### $p < 0.001$ ), PG (\*\*\* $p < 0.001$ ) and BT44 0.1  $\mu\text{g}/24\text{h}$  -treated rats (††† $p < 0.001$ ; Tukey HSD after one-way ANOVA;  $F_{4,44} = 11.365$ ;  $p < 0.001$ ). (D) Rotation rate per 5 min at 12 weeks post lesion. From 60-min time point onwards, the turning response was smaller in BT44 0.3  $\mu\text{g}/24\text{h}$  -infused rats as compared to PBS (# $p = 0.029$ ) and PG -infused rats (\*\* $p = 0.003$ ; Tukey HSD after repeated measures ANOVA; 60–120 min:  $F_{4,44} = 8.491$ ,  $p < 0.001$ ). (E-F) There were no differences between the treatment groups in spontaneous limb-use asymmetry score measured in the cylinder test at six and 12 weeks post lesion. Mean  $\pm$  SEM;  $n = 8-11$ .

Further analysis of the amphetamine-induced rotation rate per 5 min at 12 weeks post lesion revealed a distinct profile in the turning behavior of BT44 0.3  $\mu\text{g}/24\text{h}$  -treated rats as compared to the other treatment groups (Figure 5.13.D). They seemed to have a faster and nearly full recovery from the strong amphetamine-response in the beginning of the experiment: from 60-min time point onwards, the rotation rate was significantly reduced in BT44 0.3  $\mu\text{g}/24\text{h}$  group as compared to the vehicle groups.

### 5.3.2.2 Dopaminergic neurons in the SNpc and fibers in the striatum


The density of TH-ir fibers in the striatum was significantly higher in BT44 0.3  $\mu\text{g}/24\text{h}$  and GDNF 3  $\mu\text{g}/24\text{h}$  -infused rats than in PBS-infused animals (Figure 5.14.A). In accordance with this, BT44 0.3  $\mu\text{g}/24\text{h}$  and GDNF 3  $\mu\text{g}/24\text{h}$  seemed to increase the density of DAT-ir fibers in the striatum, but only the effect of GDNF 3  $\mu\text{g}/24\text{h}$  was significant when compared to the vehicles (Figure 5.14.C). Intrastriatal infusion of GDNF, but not BT44, was also able to prevent the loss of dopamine neurons in the SNpc. The number of TH-ir and DAT-ir cell bodies in GDNF-treated rats was significantly higher as compared to the vehicle-treated rats (Figure 5.14.B and D).



**Figure 5.14. BT44 restores dopamine fibers in the striatum but is not able to protect dopamine neurons in the SNpc of 6-OHDA lesioned rats.** (A) BT44 0.3 $\mu\text{g}/24\text{h}$  (# $p = 0.025$ ) and GDNF 3 $\mu\text{g}/24\text{h}$  (# $p = 0.010$ ) significantly increased the density of TH-ir fibers in the striatum as compared to PBS, while BT44 0.1 $\mu\text{g}/24\text{h}$  showed a trend for fiber restoration ( $p = 0.065$ ; Tukey HSD after one-way ANOVA;  $F_{5,46} = 3.671$ ,  $p = 0.007$ ). (B) The number of TH-ir cells was higher in the SNpc of GDNF-treated rats as compared to all other treatment groups (\*\*\* $p < 0.001$  vs. PG; ### $p < 0.001$  vs. PBS; †† $p < 0.01$  vs. BT44 0.1 $\mu\text{g}/24\text{h}$  and BT44 0.3 $\mu\text{g}/24\text{h}$ ; Tukey HSD after one-way ANOVA;  $F_{5,48} = 8.898$ ,  $p < 0.001$ ). (C) The density of striatal DAT-ir fibers was higher in GDNF-treated rats as compared to PBS (## $p = 0.001$ ), PG (\*\* $p = 0.006$ ) and BT44 0.1 $\mu\text{g}/24\text{h}$  -treated rats († $p = 0.046$ ; Tukey HSD after one-way ANOVA;  $F_{5,45} = 4.746$ ,  $p = 0.001$ ). (D) GDNF-treated rats had a greater number of DAT-ir cells in the SNpc than PBS (# $p = 0.012$ ) or BT44 0.1 $\mu\text{g}/24\text{h}$  -treated rats (†† $p = 0.008$ ; Tukey HSD after one-way ANOVA;  $F_{5,46} = 3.752$ ,  $p = 0.006$ ). The brains of the lesion control group were analyzed at two weeks post lesion to verify the extent of the lesion at the time of the treatment initiation. Mean  $\pm$  SEM; lesion control group  $n = 4$ , other groups  $n = 8-11$ .

### 5.3.2.3 Correlation of TH immunohistological measures with amphetamine-induced rotational behavior

Due to the apparent discrepancies between amphetamine-induced turning behavior and TH immunohistological outcomes at 12 weeks post lesion, we analyzed the correlations of these measures quantitatively. Amphetamine-induced rotations showed low negative correlation with TH-ir fiber density in the striatum (Pearson's  $r = -0.43$ ,  $p < 0.01$ ) and moderate negative correlation with TH-ir cell count in the SNpc (Pearson's  $r = -0.64$ ,  $p < 0.001$ ) when data from all treatment groups were pooled together (Figure 5.15.A and B). Noteworthy, GDNF-treated rats rotated substantially less as compared to all other rats, but this behavioral improvement was not accompanied by fiber restoration in the striatum to the same extent. This is visualized in Figure 5.15.A as a grouping of GDNF data points (black squares) close to the y-axis but not higher than the other data points. We also analyzed the correlation between the striatal fiber densities and nigral cell numbers from the pooled dataset and found a low correlation between these two histological read-outs (Pearson's  $r = 0.45$ ,  $p < 0.01$ ) (Figure 5.15.C).



**Figure 5.15. Correlations between TH immunohistochemical measures and amphetamine-induced turning rate.** (A) Amphetamine-induced turning rate at 12 weeks post lesion plotted against striatal TH-ir fiber density for each experimental animal. (B) Amphetamine-induced turning rate at 12 weeks post lesion plotted against nigral TH-ir neuronal number for each experimental animal. (C) TH-ir cell numbers in the SNpc plotted against TH-ir fiber densities in the striatum.  $r$  - Pearson correlation coefficient, all treatment groups analyzed together.

## 6 DISCUSSION

This work aimed to support the preclinical characterization of neurotrophic therapies for PD by testing the effects of unconventional NTFs, CDNF and MANF, in the non-lesioned rat brain and novel small molecule RET agonists, BT13 and BT44, in an animal model of PD. Patients await a disease-modifying treatment for their debilitating condition. Therefore, efforts have to be made to overcome the lack of translatability of NTF-based treatments that has overshadowed the field of research until today.

The main outcomes suggest that exogenously administered GDNF, CDNF and MANF divergently modify dopamine release and dopamine synthesizing and metabolizing enzymes in the nigrostriatal system of the normal rat brain. We saw that nigrally delivered CDNF readily diffuses around the brainstem and colocalizes with TH-ir, but not PV-ir, neurons in the SN. No active transport of CDNF into distal brain areas was detected after a nigral injection. BT13 and BT44 were shown to activate RET and its downstream intracellular target pathways responsible for neuronal survival and regeneration *in vitro*. They also supported the survival of cultured midbrain dopamine neurons in a RET-dependent manner. BT13 promoted functional recovery in a neuroprotection model of PD, and BT44 alleviated motor impairment and showed potential to restore striatal dopaminergic fibers in a neurorestoration model of PD in rats.

### 6.1 Effects of exogenously administered neurotrophic factors on dopamine release in the striatum

A single intrastriatal injection of 10 µg of GDNF, CDNF and MANF has been shown to produce robust neuroprotective effects on nigrostriatal dopamine neurons at four weeks after the injection (Lindholm et al. 2007; Voutilainen et al. 2009). Also in normal rats, a single 10 µg -injection of GDNF into the SNpc was shown to elicit neurochemical changes on dopaminergic system that persisted for at least three weeks (Hudson et al. 1995). We set out to characterize the effects of GDNF, CDNF and MANF on dopaminergic neurotransmission at one and three weeks after the striatal delivery. Importantly, our experiments were conducted in freely-moving animals in order to avoid the confounding effects of anesthetics on neuronal functions (Marinelli and McCutcheon 2014; Müller et al. 2011). Our brain microdialysis results provide the first insight into the long-lasting and divergent biological effects of exogenously administered CDNF and MANF on nigrostriatal dopamine system in the normal rat brain. These are relevant data when considering these NTFs as a potential therapeutic approach for PD. It is important to confirm that they do not decrease the release of dopamine in the striatum, and therefore have apparent interactions with the dopaminergic medications of PD.

We discovered that an intrastriatal injection of MANF significantly elevated stimulus-evoked dopamine output and enhanced DOPAC/dopamine turnover in the striatum one week after the injection. MANF may modulate presynaptic release mechanisms or storage pools of dopamine resulting in enhanced neurotransmission. Using two distinct stimuli (high concentration of K<sup>+</sup> and

amphetamine) to evoke dopamine release, we pursued to dissect the contribution of these factors to the final output. Hypertonic  $K^+$  solution depolarizes nerve terminals and causes  $Ca^{2+}$ -dependent exocytosis of vesicles close to the presynaptic membrane (Westerink et al. 1989). This pool of presynaptic dopamine is considered to be readily releasable. Amphetamine stimulus, on the contrary, gives an estimate of the total amount of dopamine stored in the synaptic terminals by depleting vesicular dopamine stores and causing  $Ca^{2+}$ -independent release of dopamine (Westerink et al. 1989; Sulzer 2011). We saw an increase both in potassium- and amphetamine-evoked dopamine overflow in MANF-injected rats. Thus, MANF seems to replenish presynaptic dopamine stores and enhance  $Ca^{2+}$ -mediated exocytosis or increase the proportion of readily releasable pool of vesicles. In primary neuron cultures and acute brain slices, GDNF has been shown to facilitate synaptic transmission by increasing the quantal size of dopamine release, potentiating  $Ca^{2+}$  influx and inhibiting A-type  $K^+$  channels (Pothos et al. 1998; Bourque and Trudeau 2000; Yang et al. 2001; Wang et al. 2003). It is possible that MANF exerts corresponding modulatory effects on the synaptic terminals. The spatiotemporal resolution of microdialysis, however, is not good enough to draw conclusions about precise mechanisms behind the observed effects. For that, further studies focusing on the cellular-level mechanisms are needed. Studies with *in vivo* voltammetry would also offer a better temporal and spatial resolution to clarify the mechanisms of enhanced dopamine release in MANF-treated rats.

As discussed below in chapter 6.4. MANF and CDFN have better diffusion properties in the brain parenchyma as compared to GDNF (Mätlik et al. 2017; Voutilainen et al. 2011, 2009). They can reach dopaminergic terminals in the whole striatum and enhance their function which can contribute to the overall dopamine outflow. GDNF, instead, readily binds to the heparan sulfate side chains of the extracellular matrix which mostly likely restricts its distribution close to the injection site and reduces the effects on distant nerve terminals. Increased sprouting of dopaminergic fibers in the striatum could also account for the enhanced dopaminergic neurotransmission in MANF-treated animals, but this seems unlikely since an earlier study demonstrated no effect of intrastriatal infusion of MANF on TH-ir fibers in the striatum of intact rats (Voutilainen et al. 2011).

Microdialysis is a highly invasive method causing mechanical damage and gliosis around the sampling site (Benveniste and Hansen 1991). ER stress and inflammation are inevitable consequences of the mechanical insult. The capability of MANF and CDFN to mitigate ER stress and neuroinflammation could partly explain their augmented effects as compared to GDNF in the microdialysis experiment. Furthermore, the implantation of the guide cannula after the NTF injection disrupts the BBB. Hypothetically, this enables NTF-neutralizing antibodies to invade to the brain which can cause unexpected variation to the effects of the NTFs.

It was unexpected that GDNF did not enhance stimulus-evoked dopamine release. By contrast, CDFN had a more pronounced effect than GDNF in spite of the fact that the half-life of exogenous GDNF in the brain has been estimated to be 3-4 days whereas the half-life of CDFN is only approximately 5.5 h (Granhölm et al. 2000; Mätlik et al. 2017). We showed that COMT activity was increased after GDNF injection. It can be speculated that the smaller effect of GDNF on stimulus-evoked dopamine release results from the enhanced dopamine metabolism which contributes to

the clearance of extracellular dopamine. GDNF/RET signaling is also shown to function as a negative regulator of the cell surface trafficking of DAT (Boger et al. 2007; Kopra et al. 2017; Littrell et al. 2012; Zhu et al. 2015). This could partly explain why amphetamine-stimulated dopamine release was not enhanced in GDNF-treated rats.

The recombinant NTFs used in Study I were produced in different cell lines which might have contributed to their divergent effects. GDNF was produced in *E. coli*, CDNF in Sf9 insect cells and MANF in mammalian Chinese hamster ovary cells. NTFs produced in mammalian cells may have stronger activity than NTFs produced in other cell lines. Proteins produced in insect or mammalian cells can be post-translationally glycosylated unlike proteins produced in bacterial cell lines. Glycosylation has an impact on the biological properties of recombinant proteins. According to mass spectrometer analysis, however, CDNF and MANF used in our experiment were not glycosylated making them comparable with GDNF in that regards.

Finally, it is tempting to speculate that the differences seen in dopamine release may originate from other neurotransmitter systems. NTFs may well affect, for example, cholinergic interneurons or glutaminergic terminals in the striatum. Increased activation of presynaptic nAChRs or NMDA receptors on dopaminergic nerve terminals would facilitate the release of dopamine. Thus, it would be informative to measure stimulus-evoked release of other neurotransmitters such as glutamate and acetylcholine in the striatum after the injection of NTFs.

## 6.2 GABA release in the globus pallidus after exogenously administered neurotrophic factors

An essential question that remains is what are the effects of NTFs on other neuronal populations outside the nigrostriatal dopamine pathway (Aron and Klein 2011). The non-dopaminergic effects are equally important to be explored because they might cause adverse events in PD patients or give rise to positive therapeutic benefits alleviating, for example, the non-motor symptoms of PD. Despite its importance, this area of research has been overshadowed by the studies focusing on the dopaminergic effects of NTFs. The effects of NTFs on the GABAergic system, for instance, are substantive since they directly affect the output of the basal ganglia and motor control. Salvatore and colleagues (2009) reported long-lasting effects of a single unilateral injection of GDNF on striatal proteins regulating GABAergic neuronal function in the non-lesioned rat brain. They described for example bilateral changes in GABA synthesizing enzyme glutamic acid decarboxylase (GAD) 65/67 and increased levels of D1 receptor and phosphorylated DARPP-32 (dopamine and cAMP regulated phosphoprotein 32 kDa) in the contralateral striatum after the striatal delivery of GDNF. Patch clamp recordings from rat midbrain slices suggest that MANF can acutely increase GABA release and GABA<sub>A</sub> receptor-mediated inhibitory postsynaptic currents in dopamine neurons of the SNpc (Zhou et al. 2006). The physiological relevance of these phenomena, however, has remained obscure.

To address the question whether GDNF, CDNF and MANF have divergent effects also on GABAergic MSNs, we performed another microdialysis experiment where we measured potassium-evoked release of GABA in the striatopallidal terminal area GPe and striatonigral terminal area SNr one

and three weeks after an intrastriatal NTF injection.  $K^+$ -stimulus caused a marked increase in the extracellular concentration of GABA in the GPe, but we did not detect significant differences between the treatment groups at either of the time points. Unfortunately, GABA concentration in the dialysates collected from the SNr was too low to be quantified reliably with our HPLC system. In the future experiments probing the SNr, either the perfusion rate of the dialysis solution has to be slower or the dialysis membrane has to be longer to yield higher concentration of GABA in the samples. No-net-flux microdialysis would also permit the determination of the actual extracellular concentration of GABA at the baseline. However, this method is not suitable for the detection of stimulus-evoked rapid concentration changes.

Due to the lack of significant differences between the treatment groups, speculations on the possible effects of NTFs on GABAergic neurotransmission are avoided here. Overall, this pilot study should be repeated in order to draw reliable conclusions from the results.

### **6.3 Effects of exogenously administered neurotrophic factors on dopamine synthesis and metabolism**

We provided novel data on the effects of CDNF and MANF on TH activity and the effect of GDNF on the activity of dopamine metabolizing enzymes COMT and MAO. To clarify if the elevated dopamine release in MANF-injected rats was due to enhanced synthesis of dopamine, we measured *in vivo* TH activity in the striatum one week after the intrastriatal NTF delivery. We saw significantly increased TH activity only in GDNF-injected rats. This observation is in accordance with earlier results showing increased TH phosphorylation and enzymatic activity in rats administered with exogenous GDNF or MEN2B mice with constitutively active RET (Beck et al. 1996; Rosenblad et al. 2003; Georgievska et al. 2004; Salvatore et al. 2004; Mijatovic et al. 2008; Salvatore et al. 2009). Perhaps, out of the NTFs studied here, only GDNF has the ability to stimulate the phosphorylation of TH. It should be noted, however, that we investigated only one time point, and therefore the data are not sufficient to draw conclusions whether the NTFs can modify TH activity at different time points. It would be worth testing if MANF and CDNF are also able to modify the phosphorylation of TH as well as the time-dependence of the effects of NTFs on TH activity. Taken together, TH activity data could not provide an explanation for the increased stimulus-evoked dopamine overflow seen in MANF-treated rats. Other possible mechanisms are discussed above in chapter 6.1.

An intriguing outcome from the microdialysis experiment was the significantly decreased DOPAC/HVA ratio in GDNF-injected rats. This phenomenon triggered us to further elucidate the effect of GDNF on the activity of COMT, MAO-A and MAO-B. Indeed, we detected increased COMT activity in the striatum of GDNF-injected rats together with reduced MAO-A activity which was, perhaps, a compensatory response to the increased COMT activity. This finding is well in line with an earlier study where a 10- $\mu$ g nigral injection of GDNF elicited a dose-dependent increase in dopamine turnover measured as HVA/dopamine ratio in the SN and striatum tissue samples one week after the injection to normal rats (Hudson et al. 1995). The observed changes in COMT and MAO-A activity direct dopamine degradation towards 3-MT-mediated pathway and away from DOPAC-mediated pathway giving a logical explanation for why DOPAC was formed relatively less



as compared to HVA. The putative changes in 3-MT levels would help to further clarify the effects of NTFs on the metabolic pathways of dopamine, but our HPLC system was not optimized to detect 3-MT in the dialysates or tissue samples.

LPS-induced microglia activation has been shown to enhance total COMT activity in the rat brain (Helkamaa et al. 2007). In our study, we controlled the possibility for microglia activation due to the surgical procedure by comparing the vehicle-injected striatum to the non-injected striatum. No differences in COMT activity between the two hemispheres were detected. Thus, the observed effect on COMT activity is not due to the injection procedure *per se*, but rather related to the injected GDNF solution. However, the rhGDNF used in the study was produced in an *E. coli* cell line. Although it was tested by the manufacturer to be compliant for cell culture use in terms of LPS counts, we cannot exclude the possibility that there were some bacterial LPS residues in the product which could induce microglia activation and result in enhanced COMT activity. To confirm the results, GDNF originating from a mammalian cell line should be tested in the same setup and microglia activation after GDNF-injection should be ruled out using e.g. Iba1 (ionized calcium-binding adapter molecule 1) immunohistochemistry. Furthermore, follow-up experiments should test whether GDNF injection has an effect on the expression level of COMT which could contribute to the increased enzyme activity. The time-dependence of COMT activation and the effects of MANF and CDNF on the activity of dopamine metabolizing enzymes would also be interesting to investigate in the future.

The neurotrophic therapies should be compatible with the current PD medications that are described in chapter 2.3.5 and Table 2.2. Thus, the increased COMT activity after GDNF treatment is an important aspect to take into consideration if GDNF is administered to PD patients. This might call for adjustments to the existing dopamine-replacing drugs; for example, the dosage of a COMT inhibitor might need to be increased. The enhanced *in vivo* TH activity in GDNF-treated rats and elevated stimulus-evoked dopamine release in MANF-treated rats, on the other hand, could translate into reduced L-DOPA doses in PD patients.

## 6.4 Spreading properties of neurotrophic factors after administration into the basal ganglia

Insufficient brain distribution and bioavailability have limited the clinical benefits of the NTFs tested in phase II trials (Lapchak et al. 1998; Hamilton et al. 2001; Salvatore et al. 2006). Knowing the diffusion and transportation properties of a neurotrophic treatment is of paramount importance when delineating the optimal site of administration for a therapy that is intracranially delivered to a single site in a condition which afflicts normal neuroanatomical connections and is accompanied with disturbed axonal transport (Chu et al. 2012; De Vos et al. 2008; O’Keeffe and Sullivan 2018). Intrastrially administered MANF has been shown to diffuse significantly better in the rat brain when compared to GDNF (Voutilainen et al. 2011, 2009). CDNF also diffuses readily and spreads to the cortical areas and hippocampus after an injection or infusion into the striatum (Mätlik et al. 2017; Voutilainen et al. 2011). GDNF and CDNF have been demonstrated to undergo retrograde transport to the SNpc along nigrostriatal dopamine neurons when injected into the striatum (Tomic et al. 1995b; Voutilainen et al. 2011; Mätlik et al. 2017). MANF, instead, seems

to be actively transported from the striatum to the frontal cortex but not to the SN (Voutilainen et al. 2009).

Our pilot experiment suggested that a 3- $\mu$ g injection of CDFN into the SN had comparable effects with a 10- $\mu$ g striatal injection of CDFN on behavioral and histological read-outs in a 6-OHDA rat model of PD (unpublished data). Therefore, we wanted to further examine how CDFN behaves when injected into the SN; how well it diffuses into the surrounding brain areas, how different variables affect the diffusion properties, and is it transported or taken up by specific types of neurons. As expected, CDFN had a widespread diffusion in the brainstem of normal rats after the nigral injection. CDFN immunolabelling was detected at 2 and 6 h, but not anymore at 24 h post injection. This suggests that the exogenous CDFN was degraded between 6 and 24 h which goes well in line with the 5.5-h half-life of CDFN (Mätlik et al. 2017). The volume of distribution of injected CDFN was increased with the increased injection volume which is clearly an important factor to be optimized when the delivery protocol for a therapeutic protein is being established. A larger injection volume results in the diffusion of the protein to more distant brain regions which may give rise to enhanced therapeutic effects or emerge of adverse effects depending on the protein in question, the affected brain regions and the clinical condition.

We saw similar punctate CDFN-ir staining inside the TH-ir neurons of the SNpc as was previously shown after striatal injection and retrograde transportation of CDFN to the SNpc (Mätlik et al. 2017). CDFN did not seem colocalize with PV-ir neurons in the SNpc suggesting selective uptake into dopamine neurons. This can possibly contribute to the neuroprotective effects of CDFN on dopamine neurons and its favorable safety profile. Further experiments with quantitative outcomes, however, are needed to draw decisive conclusions of the localization of exogenously administrated CDFN.

When radiolabeled CDFN was injected into the SN we observed prominent spread to the ipsilateral STN, but no active transport to the striatum or other distal brain areas. It is possible that upon cellular uptake CDFN is targeted to lysosomal degradation instead of axonal transportation when injected near the cell bodies as is the case with NGF (Butowt and von Bartheld 2001). Noteworthy, only one time point was analyzed after the injection of radiolabeled CDFN. As seen in the diffusion experiment, there is little CDFN immunoreactivity left in the brain at 24 h post injection, probably due to the short half-life of CDFN. It is possible that CDFN is transported from the SN but not detected in other brain areas 24 h after the injection anymore. Thus, it would be good to check if there are signs of active transport at earlier time points (e.g. at 2 h and 6 h) after the injection. Co-administration of unlabeled CDFN did not significantly reduce the radioactive signal in the STN suggesting that CDFN was passively diffused to this proximal nucleus instead of actively transported. Active transportation, however, cannot be completely ruled out because the prominent diffusion of radiolabeled CDFN to the area covering the STN potentially masks the effect of the transportation.

## 6.5 Potential of neurotrophic factor mimetics as a therapeutic strategy for Parkinson's disease

Since the clinical proof-of-concept studies with NTFs have remained inconclusive, it is pivotal to look for novel neurotrophic strategies to protect the degenerating dopamine neurons and restore the impaired functionality of basal ganglia circuitry. There are several protein therapy related challenges that may limit the therapeutic use of NTFs for PD or other brain diseases. Firstly, proteins do not cross the BBB and therefore require surgical delivery into the brain which increases the costs and risks of the treatment. In the clinical studies with GDNF and AAV2-NTRN, the most common safety concerns have related to the intracranial drug delivery procedure and implantation of the infusion device (Lang et al. 2006; Marks et al. 2010; Whone et al. 2019). Secondly, the production of biological therapeutics for clinical use is not trivial. Special attention has to be paid to the batch-to-batch variations and stability of the recombinant protein. Additionally, the levels host cell contaminants or endotoxins (such as LPS) have to be carefully controlled. Thirdly, formation of neutralizing antibodies against the recombinant NTF treatment is a particular concern (Lang et al. 2006; Heiss et al. 2019). Anti-NTF antibodies can potentially limit the efficacy of the therapeutic protein and cross-react with the corresponding endogenous NTF causing its loss-of-function. Finally, low bioavailability and poor tissue penetration are well-known issues with the GFL-therapeutics as discussed in the previous chapter. Small molecules that retain the neurotrophic activity of NTFs by activating the same receptors and signaling pathways provide an attractive approach to circumvent these protein therapy related challenges.

Using virtual and cell-based screening methods a novel family of small molecule GFL mimetics, including BT13 and BT44, has been identified and optimized (Saarma et al. 2014; Sidorova et al. 2017). These compounds seem to diffuse readily in the brain parenchyma and penetrate the BBB. They activate GFL-GFR $\alpha$ -RET signaling complex, but the exact mechanism of action is not elucidated yet. Molecular docking simulations have identified two possible binding sites: the putative allosteric modulation site in GFR $\alpha$  or the GFR $\alpha$ -interfacing surface of RET (Ivanova et al. 2018). The latter would imply that the compounds act as a direct RET agonists. Recent results support their function as direct RET agonists because of their ability to phosphorylate RET and activate downstream signaling cascades also in the absence of GFR $\alpha$  co-receptors (Sidorova et al. 2017; Viisanen et al. 2020; Study III; Study IV). BT13 and BT44 are first-in-class molecules with drug-like properties that selectively (see below) activate RET and its downstream survival-promoting signaling pathways *in vitro*. This work demonstrated that they can markedly alleviate amphetamine-induced rotational behavior in a unilateral 6-OHDA lesion model of PD in rats. BT44 also showed potential for restoring TH-ir fibers in the striatum of hemiparkinsonian rats, but the neuroprotective effects on dopaminergic cell bodies in the SN remained unattained. BT13 and BT44 serve as promising lead compounds; with further optimization they can be developed into a novel disease-modifying therapy for PD.

The potential off-target effects of BT44 were assessed in a panel of *in vitro* assays by an external company (Eurofins CEREP SA, France, see details Study IV supplementary data). BT44 (1  $\mu$ M) did not affect the activity of selected ion channels, G-protein coupled receptors, transporters, kinases or dopamine metabolizing enzymes. In all assays, the target inhibition or stimulation was less than

25% which, according to the company's guidelines for result interpretation, reflects assay variability and indicates the lack of a significant effect of the test compound. Our data also show that BT44 only supports the survival of RET-expressing wild-type, but not RET knockout, dopamine neurons and fails to activate ERK-related signaling in TrkB-expressing murine fibroblasts lending support to the selectivity of BT44 for RET.

BT44 promoted functional recovery in 6-OHDA lesioned rats. When we analyzed the amphetamine-induced turning rate per 5 min at 12 weeks post lesion, we saw an interesting feature in BT44 0.3  $\mu$ g/24h -treated animals. They showed divergent time-dependence of the rotational asymmetry and seemed to recover faster from the amphetamine challenge when compared to other treatment groups. It can be speculated that BT44 alters dopamine dynamics. For example, reduced dopamine uptake through DAT or activity of dopamine metabolizing enzymes could account for slower clearance of dopamine from the extracellular space of the lesioned striatum, and consequently faster recovery from the amphetamine-induced turning response. Important to note that in Study III, BT13 was shown to acutely increase the level of extracellular dopamine in the dorsal striatum in mice. As mentioned earlier, RET signaling is suggested to negatively regulate DAT function and/or cell surface trafficking (Boger et al. 2007; Kopra et al. 2017; Littrell et al. 2012; Zhu et al. 2015). Moreover, it is possible that as a lipophilic compound, BT44 is able to diffuse into the non-lesioned hemisphere during the long infusion period where it could augment dopamine clearance from the extracellular space, for example by enhancing COMT-mediated degradation of dopamine as was shown to happen after GDNF administration in Study I. Further experiments assessing, for example, the effects of BT44 on DAT or COMT activity or its diffusion into the contralateral striatum are needed to explain the altered rotation rate profile.

In the *in vivo* experiments, we encountered solubility problems related BT-compounds; they were insoluble to aqueous vehicles forcing us to use PG as a vehicle. Even with PG, the solution required heating and sonication in order to get the compounds thoroughly dissolved. We cannot be sure if any precipitation occurred in the infusion pumps or when the solution got in contact with aqueous extracellular fluid. These solubility issues should be resolved before proceeding to further *in vivo* experiments. The high-resolution 3D structure of GFR $\alpha$ 1 and the model for GDNF-GFR $\alpha$ 1 interaction provide an opportunity for rational structure-based drug design (Leppänen et al. 2004). Novel molecules capable of binding to specific GFR $\alpha$  receptors with improved drug-like properties could help to overcome some of the difficulties related to the current BT compounds. After lead optimization, either BT-scaffold-based compounds or new GFL mimetics could be suitable for systemic administration. Compared to intracranial delivery it would be a less risky approach for large-scale clinical use and would help to overcome the problematic question about the most effective delivery site in the brain. However, the peripheral expression of GFR $\alpha$  and RET receptors should be considered when systemic administration of GFL mimetics is planned. Adverse effects can occur as a result of RET activation for example in the PNS, testis or thyroid gland.

Apart from direct RET agonists, an attractive therapeutic approach for harnessing RET signaling pathways would be to develop selective positive allosteric modulators (PAMs) of GFR $\alpha$ 1. The first compound possessing such biological activity, XIB4035, has been described, but not tested in

animal models of PD (Tokugawa et al. 2003; Hedstrom et al. 2014). GFR $\alpha$ 1-PAMs would potentiate the trophic effects of endogenous GDNF and provide major advantages over direct RET agonists: Firstly, they could reduce the risk of RET-induced adverse effects as compared to exogenous GDNF or direct RET agonists. The activation of RET would only occur in the presence of endogenous GDNF preserving the homeostatic regulation mechanisms of GDNF-GFR $\alpha$ 1-RET signaling, and administration of unphysiological concentrations of GDNF would not be needed. Secondly, they could provide superior selectivity as compared to direct RET agonists owing to the limited expression of particular GFR $\alpha$ -RET pairs in different brain areas and other organs (Trupp et al. 1997; Golden et al. 1998). Lastly, binding of a compound directly to RET on the surface interfacing with GFR $\alpha$ 1 can potentially disrupt the signaling of endogenous GDNF. This partial antagonism could explain the submaximal efficacy of BT44 as compared to GDNF in our neurorestoration experiment (Study IV). In addition, continuous RET activation during the treatment infusion period might have caused downregulation of the receptor. In this case, subsequent withdrawal of the RET agonist would lead to an acute depletion of trophic support contributing to the lack of neuroprotective effect on nigral neurons.

## 6.6 Methodological considerations

### 6.6.1 Considerations relating to 6-OHDA lesion models

In our neuroprotection experiment (Study III) evaluating the efficacy of BT13 in hemiparkinsonian rats, a single unilateral injection of 6-OHDA (16  $\mu$ g) into the dorsal striatum resulted in relatively mild dopaminergic degeneration: in vehicle-treated rats, the density of TH-ir fibers in the striatum was reduced approximately by 55% and the number of TH-ir neurons in the SNpc by 27%, which correspond to the estimated state of the nigrostriatal system in PD patients at the onset of the motor symptoms, i.e. at early stage of the disease (Burke and O'Malley 2013). The lesion size was similar than the “regressive” lesion described by Penttinen et al. (2016). Because the size of the lesion was at the threshold of the motor manifestations, small changes at cellular level may have caused disproportionally profound functional improvement as seen with the almost completely vanished rotational behavior in GDNF and BT13 -treated rats (Björklund and Dunnett 2019). We also saw spontaneous recovery in terms of rotational behavior in vehicle-treated animals. This phenomenon is well-documented in the literature for rats with a small (0-30% loss of the SNpc neurons) or medium (30-75% loss of the SNpc neurons) 6-OHDA lesion (Stanic et al. 2003; Stanic et al. 2003b; Penttinen et al. 2016). The clear amphetamine-induced turning bias observed after two weeks is recovered by 10-16 weeks post lesion in animals with a medium-size lesion and even faster in animals with a small lesion. This functional recovery is associated with axonal regrowth, sprouting and phenotypic recovery of the nigrostriatal dopamine neurons (Blanchard et al. 1996; Finkelstein et al. 2000; Stanic et al. 2003). Spontaneous recovery is a confounding factor in our neuroprotection experiment implicating that the functional effects of BT13 have to be interpreted with adequate caution. The lack of effect on TH-ir and DAT-ir fibers in the striatum reinforces the view that the reduction in amphetamine-induced rotational behavior is not due to neuroprotection. Perhaps, a larger 6-OHDA lesion could have produced a more stable lesion with a better correlation between the histological and functional read-outs.

In the next experiment (Study IV), we wanted to utilize neurorestoration paradigm with a much more severe 6-OHDA lesion. In the neurorestoration model, it is possible to balance animals into experimental groups according to baseline rotation scores after the lesion (Figure 4.2.). This helps to control interindividual variation and reduce the groups sizes needed to see therapeutic effects. Of note, as BT13 was speculated to regulate DAT activity (Study III), we wanted to make sure that BT44 could not directly interfere with the neurotoxic effect of 6-OHDA which is dependent on DAT-mediated uptake into dopamine neurons. Most importantly, the delivery of the experimental compounds at the time when the lesion is already established reflects much better the clinical context in PD making the neurorestoration paradigm a more relevant model as compared to the neuroprotection paradigm. In addition, we wanted to produce a bigger and more stable lesion in order to get rid of spontaneous recovery and provide a larger window for the experimental therapeutics to produce their effects.

The 6-OHDA delivery paradigm employed in our neurorestoration experiment, indeed, resulted in a much more severe lesion in agreement with the earlier results (Penttinen et al. 2016). In the lesion control group, the density of TH-ir fibers in the striatum was reduced approximately by 90% and the number of TH-ir cell bodies in the SNpc by 77% at two weeks post lesion. The number of TH-ir cells in the SNpc of PBS and PG -treated animals at 12 weeks post lesion was significantly lower as compared to the lesion control group indicating that the dopaminergic degeneration was not fully developed at the time of the treatment initiation but rather retrogradely progressing. It is well known that striatal denervation occurs rapidly after 6-OHDA injection into the striatum, but the cell bodies in the SNpc die gradually over several weeks or months (Björklund et al. 1997). The lesion size in the neurorestoration experiment reflects the state of the nigrostriatal system in advanced PD, i.e. the status of patients when intracerebral treatment procedures can be considered (Kordower et al. 2013). The neuronal damage was accompanied by persistent motor deficits as indicated by prominent amphetamine-induced rotational asymmetry and decrease in spontaneous contralateral forepaw use in the cylinder test in vehicle-treated rats. Regardless of the severe lesion, BT44 was able to induce delayed functional recovery in turning behavior and show small, but significant, restorative effect on dopaminergic processes in the striatum. Considering that the behavioral data from the cylinder test were not in line with the amphetamine-induced rotation data and no effects on dopaminergic cell bodies in the SNpc were observed, the neurorestorative potential of BT44 need to be confirmed in future experiments. It is recognized, that the aggressive lesion may have limited the neurorestorative effects of BT44 in our study.

Earlier studies have shown that striatal 6-OHDA lesion can downregulate GFR $\alpha$ 1 and RET expression in the nigrostriatal pathway up to one month post lesion (Marco et al. 2002; Kozlowski et al. 2004; Gavin et al. 2014). Due to the fact that the delivery of the RET agonists occurred during this period of time, the downregulation of RET can negatively affect their neuroprotective efficacy. The extent of the receptor downregulation in the disease model would be an important factor to clarify in the future before proceeding into large-scale efficacy studies in order to gain better understanding of the potency of the experimental compounds *in vivo*. The ability of GDNF to halt the degeneration of dopaminergic cell bodies is facilitated by its retrograde transport to the SNpc via the remaining axons while there are no data demonstrating the retrograde transport of BT-

compounds (Lapchak et al. 1997; Tomac et al. 1995). Obviously, the considerably higher potency of GDNF as compared to BT-compounds as illustrated in Table 5.1 also accounts for its better efficacy in the neuroprotection and restoration experiments.

The unilateral 6-OHDA lesion model is based upon the imbalance of the nigrostriatal system between the hemispheres. The effects of unilaterally delivered interventions are compared to the intact side of the brain. Thus, this model is not optimal for testing the effects of lipid soluble compounds that readily spread to the contralateral hemisphere after unilateral administration and potentially elicit effects there too. BT-compounds tested in our studies are very lipophilic. Their putative diffusion to the control side during the seven- or 14-day infusion may compromise the read-outs that are based on measuring the asymmetry between the hemispheres.

## **6.6.2 Relevance of the behavioral tests in 6-OHDA lesion models**

### **6.6.2.1 Correlation between amphetamine-induced turning rate and nigrostriatal integrity**

Although amphetamine-induced rotation test has become the standard tool to monitor motor impairment and functional recovery in experiments with neuroprotective interventions, there are some important pitfalls that should be carefully considered in the future use of this test like Björklund and Dunnett (2019) have described in their critical reappraisal. We also analyzed the correlation between amphetamine-induced turning rate and the striatal TH-ir fiber density and nigral TH-ir cell number in the neurorestoration experiment (Study IV). In agreement with earlier reports, we did not find good correlations between these measures (Winkler et al. 1996; Kirik et al. 1998; Kozłowski et al. 2000; Voutilainen et al. 2009; Tronci et al. 2012; Björklund and Dunnett 2019).

Other neuronal circuits than nigrostriatal dopamine pathway may be responsible for the functional recovery seen in the experiment. For example, amphetamine induces dopamine release from dendritic vesicles in the SNr where dopamine primarily acts at presynaptic D1 receptors of the striatonigral MSNs facilitating GABA release (Rommelfanger and Wichmann 2010; Timmerman and Abercrombie 1996). This reduces the overactive firing of GABAergic SNr neurons projecting to the thalamus that is implicated in the motor imbalance of the unilateral 6-OHDA model. In our neurorestoration experiment, amphetamine-induced dendritic dopamine release in the SNr may have been augmented in GDNF-treated rats due to the significant preservation of dopaminergic cell bodies in the SNpc. Indeed, *in vivo* microdialysis measurements have also demonstrated enhanced amphetamine-evoked dopamine release in the SN of GDNF-treated rats (Hoffman et al. 1997). Thus, GDNF-induced significant functional improvement in animals with advanced dopaminergic denervation can be mediated by increased dopamine release in the striatal output structures. The sparse remaining striatal innervation may have a minor contribution in this case giving an explanation for the mismatch between the turning behavior and striatal fiber preservation.

In order to avoid amphetamine-evoked dopamine release in the extra-striatal structures as a possible confounding factor, it could have been beneficial to include apomorphine-induced rotational assessment in our behavioral test battery. The denervation of the striatum was at the level that is generally considered to be sufficient to cause supersensitization of the postsynaptic dopamine receptors which is a prerequisite for apomorphine-induced rotation test (Ungerstedt 1971). This test has been reported to correlate better with the loss of both striatal TH-ir fibers and nigral TH-ir cell bodies in 6-OHDA lesioned mice as compared to amphetamine-induced rotation test (Grealish et al. 2010). Apomorphine-induced turning behavior also had a better capacity to discriminate mice according to the extent of their nigrostriatal damage.

#### 6.6.2.2 Other behavioral tests

Amphetamine-induced rotation test is a practical experiment to assess motor asymmetry and functional recovery in neuroprotection studies. PD patients, however, show a more complex repertoire of motor deficits than can be recapitulated with simple drug-induced motor tests. Therefore, behavioral tests allowing quantification of spontaneous motor and sensorimotor functions would provide more informative outcomes (Dunnett 2005). It is recommended to combine amphetamine-induced rotation test with at least one other test measuring spontaneous motor behavior (Björklund and Dunnett 2019). For 6-OHDA lesioned rats, paw-use asymmetry measured with the cylinder, stepping or staircase test, and for 6-OHDA lesioned mice e.g. sensorimotor behavior in the corridor test, would be profitable adjuncts.

For these reasons, we also analyzed spontaneous forepaw-use asymmetry using the cylinder test in our neurorestoration experiment (Study IV). Despite significant recovery in the amphetamine-induced turning behavior, neither GDNF nor BT44 were able to normalize forepaw use in the cylinder test. These data are in accordance with earlier reports where GDNF protein injection or gene therapy into 6-OHDA-lesioned striatum significantly alleviated amphetamine-induced turning behavior but showed no effects in the cylinder test (Georgievska et al. 2002; Gasmi et al. 2007b; Yue et al. 2014). Also in 6-OHDA lesioned mice, the behavioral deficits observed in the cylinder test showed poor correlation with amphetamine-induced turning behavior as well as with the integrity of the nigrostriatal pathway (Grealish et al. 2010). It should be noted, that not only the total extent of striatal dopaminergic denervation, but also its regional distribution affects the functional output and magnitude of the motor impairment (Kirik et al. 1998; Björklund and Dunnett 2019). Amphetamine-induced turning rate is known to be particularly sensitive to changes in dopaminergic innervation within the dorsomedial and dorsolateral parts of the striatum, whereas the ventrolateral parts of the striatum have more pronounced control on movement initiation, sensorimotor orientation and skilled motor behavior. Since our treatment infusions were directed into the dorsomedial striatum, they might have had limited effects on the ventrolateral striatum, and thus on spontaneous forepaw use in the cylinder test.

Notably, most of the behavioral tests used to assess the non-motor symptoms of PD are not particularly designed for parkinsonian animals (Asakawa et al. 2016). We are lacking specific tools for the assessment of non-motor symptoms especially in rodent models of PD. Development of such behavioral tests would be essential for the advancement of this important, but rather



neglected field of research. Apart from the behavioral test, also other approaches can be considered to follow longitudinal functional changes in animal models of PD. For example, *in vivo* electrophysiology offers another approach to detect functional changes in various neuronal circuits in freely-moving animals.

### 6.6.3 Validity of the disease models

The encouraging preclinical results with NTF-based therapies have failed to translate into positive outcomes in clinical trials. To enable successful clinical translation of disease-modifying treatments, we should take a critical look at the limitations of the current preclinical models. Are the results obtained with these models relevant in terms of the clinical condition? As experimental animals do not naturally develop PD, the disease has to be artificially induced by targeting one or few pathological processes of the disease.

A major obstacle in developing effective disease-modifying therapies for PD is the lack of animal models that fully recapitulate the pathological and functional disease progression and the phenotype of PD (Paul and Sullivan 2018). The current animal models typically mimic only one or few aspects of the disease. The traditional neurotoxin-based (e.g. 6-OHDA and MPTP) animal models, for example, reproduce the degeneration of nigrostriatal dopamine neurons and neuroinflammation but fail to replicate several other crucial features such as the accumulation of  $\alpha$ -syn containing protein inclusions. The mechanism of the neuronal insult is also different from the human disease. One of the important characteristics of human PD is that the pathology progresses over decades. Neurotoxins, instead, produce an acute lesion leaving little time for compensatory mechanisms to take place. The genetic animal models of PD seem to exhibit good construct validity as they are based on mutations that cause genetic forms of the human disease. However, they fail to produce a clear parkinsonian phenotype in terms of neurodegeneration and motor symptoms.

Traditionally, the neurotoxin-induced animal models are focused on insulting the nigrostriatal dopamine neurons. This, indeed, offers a good approach to model motor dysfunctions but does not take into consideration the widespread nature of the disease. PD pathology is not restricted to the nigrostriatal dopamine system but affects multiple neuronal populations and neurotransmitter systems, which also give rise to the numerous non-motor symptoms of PD and should be better considered in preclinical drug development (Olanow et al. 2015b). Overall, the neurotoxin models serve as good tools for testing symptomatic treatments but might not be as useful in the development of disease-modifying therapies.

One common issue with the rodent models of PD is the routine use of young adult animals as experimental subjects. Metabolism, plasticity and neuronal function in the young brain differ from those of the aged brain which makes the translation to PD patients even harder. Perhaps, establishment of disease models in aged animals could partly help to overcome this translatability problem. Another common problem relates to the timing of a neurotrophic therapy (Paul and Sullivan 2018). Studies where the therapy is administered after the nigrostriatal pathway has already degenerated, like in our neurorestoration experiment (Study IV), are more relevant to the

clinical scenario where a substantial nigrostriatal damage already exists at the time when the therapy can be initiated. In the majority of the preclinical studies, however, NTFs have been administered before or shortly after the lesion. Thus, there is a mismatch with clinical trials where the NTF interventions are mostly tested in patients with advanced PD.

Since most animal models tend to reproduce only few features of human PD, it is essential to test potential treatments in a combination of different disease models. In addition to neurotoxin-based models,  $\alpha$ -syn-based models should be part of the preclinical testing. Neuroinflammation, as one important aspect of the PD pathobiology, also deserves more attention. Furthermore, dopamine neurons generated from PD patient-derived induced pluripotent stem (iPS) cells should be included in the battery of preclinical PD models. They harbor the full human-specific metabolic and regulatory pathways, and thus provide a valuable tool to investigate dopamine metabolism and mechanisms of dopaminergic degeneration (Meiser et al. 2013). Other possible platforms to be used in drug development include, for example, organoids, 3D cell constructs and organ-on-a-chips. The importance of such new platforms is growing, and they will help us to replace some of the *in vivo* experiments according to 3R principles.

If neurotrophic therapies will enter the clinical use, they will most probably be used in combination with standard PD medications. Therefore, it is of outmost importance to evaluate their efficacy in animal models together with other anti-parkinsonian treatments. For example, GDNF delivery into the lateral ventricle of MPTP-lesioned marmosets was shown to reduce the severity of L-DOPA-induced dyskinesias (Iravani et al. 2001). These type of co-administration experiments are warranted to improve the translatability of preclinical NTF studies.

## 6.7 General discussion

PD is a common neurodegenerative disorder with complex pathobiology and multitude of genetic and environmental risk factors (Kalia and Lang 2015). The diagnosis is preceded by a several years or even decades long prodromal phase, during which the characteristic parkinsonian Lewy pathology progresses from the periphery into the brain. Eventually, patients become afflicted by cardinal motor symptoms that arise from the progressive loss of nigrostriatal dopamine neurons. Equally importantly, several other neuronal populations are also affected which contributes to various non-motor symptoms of PD. The current treatments mainly target the motor impairment by replenishing reduced dopaminergic neurotransmission within the striatum. None of them, however, is capable to modify the neurodegenerative process or slow down the progression of the disease. Novel disease-modifying agents are urgently needed. NTFs and their biological effects mimicking small molecule compounds hold promise for this quest, but more preclinical work is required to further characterize their mechanisms of action, confirm their neuroprotective potential and uncover their effects on other brain cells than nigrostriatal dopamine neurons. It is not clear, for example, whether exogenous NTFs directed to the nigrostriatal dopamine system have significant effects on the non-motor symptoms of PD. Systemically administered NTF mimetics with widespread target engagement in all afflicted brain areas could potentially help to tackle also these disabling features of the disease.

PD is a clinically heterogeneous disorder. There are many ways to classify various subtypes of PD, for example based on motor symptoms (e.g. tremor-dominant vs. non-tremor-dominant), cognitive features, age at onset, rate of progression or various combinations of these (Marras and Lang 2013). The different subtypes are believed to have distinct etiologies and pathogenesis. Better understanding of those would allow the identification of novel therapeutic targets and open avenues for the development of effective and safe disease-modifying therapies.

GDNF and NRTN have been considered as the most promising candidates for the first neurotrophic therapy for PD based on the solid body of preclinical evidence. Failures in the phase II clinical trials, however, have raised questions whether RET is an optimal target receptor. In PD models induced by viral vector-mediated overexpression of  $\alpha$ -syn, gene delivery of GDNF to the SNpc had no effect on the survival of dopamine neurons or motor impairment (Lo Bianco et al. 2004; Decressac et al. 2011). Importantly,  $\alpha$ -syn overexpression was shown to downregulate the transcription factor Nurr1 and its downstream target RET diminishing the intracellular response to GDNF (Decressac et al. 2012). Nurr1 and RET expression was also found to be reduced in nigral neurons in PD patients suggesting that  $\alpha$ -syn-induced downregulation of RET might account for the negative results of GDNF in the  $\alpha$ -syn overexpressing animal models and its limited efficacy in the clinical trials (Chu et al. 2006; Decressac et al. 2012). It should be noted, however, that there are also contrasting reports showing no downregulation of Nurr1 or RET in PD patients (Bäckman et al. 2006; Su et al. 2017; Walker et al. 1998). Formation of  $\alpha$ -syn aggregates may also impede the axonal transport of GDNF like it has been shown to impair the transportation of BDNF signaling endosomes (Chung et al. 2009; Volpicelli-Daley et al. 2014). This is an important factor to consider when planning the future experiments and deciding the administration sites of NTFs.

Gain-of-function mutations in *Ret* gene causing the inherited cancer syndromes MEN2 and familial medullary thyroid carcinoma have triggered concerns regarding the oncogenic potential of RET activation (Plaza-Menacho et al. 2006). However, with an intermittent dosing scheme, the activation of RET would not be continuous like in the RET-related cancers which would help to manage the risk for tumors. Important to note, that in a recent study investigating the long-term effects of GDNF overexpression from the endogenous locus, continuously increased activation of GDNF-RET signaling did not elicit adverse effects or tumors (Turconi et al. 2020). Intermittent delivery also results in smaller cumulative doses as compared to continuous dosing schemes (Whone et al. 2019). This further helps to avoid potential safety risks. Target engagement in a pulsatile fashion may also offer benefits over continuous receptor stimulation in terms of ligand-dependent receptor desensitization that potentially reduces the responsiveness of target tissues to a NTF therapy (Lohse 1993).

Due to the abovementioned issues, it is vital to continue looking also for other, RET-independent opportunities for a disease-modifying therapy. Here, unconventional NTFs CDNF and MANF have shown great promise in preclinical studies. However, our knowledge of their biological role and mechanism(s) of action is still limited (Lindahl et al. 2017). The discovery of their putative receptors would enable better understanding of their physiological functions and development of new neurotrophic candidates based on their signaling mechanisms.

### 6.7.1 Importance of biomarker development

It is imperative to develop better diagnostic tools and novel biomarkers in parallel with disease-modifying therapies to allow earlier intervention with neuroprotective agents, and consequently an improved therapeutic outcome. With the current diagnostic criteria, PD is identified only upon the manifestation of the motor symptoms which occurs several years after the neurodegenerative process has started (Miller and O'Callaghan 2015; Parnetti et al. 2019). This hinders the attempts to intervene at the earliest phase of the disease when there is still a sufficient number of surviving dopaminergic neurons and axons that are capable to respond to the therapy. A reliable prodromal biomarker could provide a prolonged time frame, during which disease-modifying candidates could be administered to halt or slow down the cell loss. Currently, this time frame is not very long from the diagnosis to virtually complete loss of dopaminergic fibers in the putamen (4 to 5 years) (Burke and O'Malley 2013; Kordower et al. 2013). The earlier the NTF-based therapy can be initiated, the better efficacy it is expected to have as was demonstrated in the clinical trial with AAV2-NRTN (Olanow et al. 2015). In an effort to enable enrolment of patients at the earliest stage of PD into clinical trials of neuroprotective therapies, the International Parkinson and Movement Disorder Society task force has developed research criteria for prodromal PD (Berg et al. 2015). Identification of individuals in the prodromal phase would also facilitate the clarification of the disease mechanisms and progression. On the other hand, there is a greater risk for misdiagnosis, particularly with atypical parkinsonian syndromes, at the early stages of the disease (Tolosa et al. 2006).

PD should not be conceptualized as a unitary disorder (Berg et al. 2014). It comprises a spectrum of subtypes with distinct etiologies, pathological processes and therapeutic needs. Validated biomarkers would help to stratify patients according to their individual form of the disease. More precise diagnosis would pave the way for better understanding of the pathological processes in each subtype and more personalized disease management. Importantly, disease subtyping would contribute to the development of more translational disease models and design of more successful clinical trials with stratified inclusion criteria. The failed attempts to find neuroprotective strategies for PD may stem from the reductionist approach in the conducted clinical trials which have paid little attention to the variability of the disease at the individual level (Espay et al. 2017).

Currently, UPDRS is used for longitudinal follow-up of the disease progression based on episodic and subjective clinical evaluations. Thus, drug development decisions in clinical trials rely on sparse datapoints contributing to frequent and costly failures. There is a need for reliable longitudinal biomarkers. Development of digital biomarkers, for example, may help to address many of the current diagnostic shortcomings in an economical fashion. They would allow an objective approach to continuously track fluctuations in motor and non-motor symptoms during patients' daily life. The resulting rich real-world datasets may prove to be highly predictive in assessing clinical improvement in PD studies and permit personalized therapeutic adaptations.

Ideally, preclinical drug development would already be accompanied with a translatable biomarker that could be followed through the whole process from animal models to clinical trials

and regular patient monitoring. This kind of translatable biomarker would help to predict the effects of an intervention in a patient population based on preclinical tests in animal models. For example, imaging modalities would be suitable for translatable biomarkers because they provide a direct approach to measure specific neurofunctional properties and the same methodology can be applied both to experimental animals and humans.

### **6.7.2 Issues with the clinical studies**

Despite promising outcomes in preclinical studies and open-label phase I trials, none of the controlled double-blind phase II trials have reported clear clinical benefit for intracranial GDNF or NRTN therapy in patients with moderate or advanced PD (Hegarty et al. 2017). Negative outcomes from these proof-of-concept studies have raised questions whether NTFs in general are translatable into an effective disease-modifying therapy or whether the trials have been conducted sub-optimally. If the NTF hypothesis is still valid, the limitations of the studies should be carefully appraised.

The importance of conducting placebo-controlled double-blind trials should be underlined. Marked placebo effect is a well-described phenomenon in PD trials, especially when neurosurgical procedures are involved, which may explain the efficacy differences between the open-label and controlled double-blind trials (Goetz et al. 2008; Alterman et al. 2011). Therefore, no conclusions should be drawn from the open-label studies that are not properly controlled for the placebo effect.

Patients enrolled in the phase II trials may have been too advanced to benefit from the NTF therapy (Olanow et al. 2015). Dopamine neurons are so profoundly degenerated and dysfunctional at the later stages of PD that they might not be capable of responding to the therapy anymore. Indeed, greater benefits have been observed in a subgroup of patients who received AAV2-NRTN therapy less than five years after the diagnosis (Bartus 2015; Olanow et al. 2015). In the future, it might be necessary to recruit patients at an earlier stage of the disease which complicates weighing the risks against the benefits.

Transport of intraputamenally delivered GDNF or NRTN to the dopaminergic cell bodies in the SNpc is thought to be necessary for their pro-survival effects. Thus, the feasibility of treating advanced PD patients with these factors is questioned because the putamenal innervation is almost completely disappeared within 4-5 years post diagnosis (Kordower et al. 2013). This denervation potentially contributes to the impaired retrograde axonal transport, as reported in AAV2-NRTN-treated patients enrolled in the clinical trial more than five years post diagnosis (Bartus et al. 2011; Bartus et al. 2015).

There are some indications of clinical benefits at the later time points of the trials suggesting delayed and sustained effects for NTFs (Love et al. 2005; Marks et al. 2010; Patel et al. 2005). Thus, a longer-term follow-up might be required to detect possible clinical improvements. Ethical aspects, however, need to be considered in order to avoid excessively long exposure to placebo in a surgical setting.

It has been suggested that technical differences in the drug delivery systems and infusion protocols partly contributed to the varying results between the phase I and phase II trials (Salvatore et al. 2006; Sherer et al. 2006; Morrison et al. 2007). Furthermore, explanations for the lack of efficacy in the phase II trials can also include insufficient dosage, heterogeneous or inadequate distribution of GDNF and NRTN in the target tissue and formation of neutralizing antibodies (Tatarewicz et al. 2007).

### **6.7.3 Future directions**

Despite holding great therapeutic potential for PD, it might well be that NTFs are not sufficient alone to combat neurodegeneration and restore neuronal function. Combination therapies may offer an opportunity to maximize the benefits of NTFs. Coadministration of different NTFs seems to be more efficient in protecting, repairing and activating dopamine neurons than the delivery of a single NTF (Voutilainen et al. 2017). NTFs could provide additive therapeutic benefit also when administered in conjunction with other treatment strategies. Usually, synergistic therapies help to reduce the dose of each individual treatment, and thus mitigate the risk for possible adverse effects.

It is becoming evident that the underlying pathological processes such as Lewy pathology or the augmented neuroinflammatory state have to be resolved in order to achieve favorable conditions for neuroregeneration. Only then, neurotrophic therapies would have a better opportunity to promote neuronal survival. Thus, it would be worthwhile to study the effects of NTFs in conjunction with, for example, experimental anti- $\alpha$ -syn antibodies that reduce the spread of  $\alpha$ -syn aggregates in the brain. Chronic neuroinflammation is one of the key elements in PD pathogenesis contributing to the neuronal dysfunction and degeneration. Reactive microglia are the primary mediators of neuroinflammation. Astrocytes, in turn, regulate the microglial responses by secreting factors that reduce their phagocytic activity and production of ROS. Therefore, combining anti-inflammatory (or immunomodulatory) treatments with trophic factors may open new avenues to induce regenerative processes in PD. Interestingly, experiments in cultured midbrain microglial cells identified GDNF as one of the major astrocyte-derived factors regulating their activation in a GFR $\alpha$ 1 receptor -dependent manner (Rocha et al. 2012). CDNF and MANF have also been shown to reduce the secretion of pro-inflammatory cytokines from glial cells suggesting an important regulatory role in neuroinflammation (Zhao et al. 2013; Zhu et al. 2016; Neves et al. 2016). Thus, they might provide synergistic benefits if administered together with other NTFs (Voutilainen et al. 2017).

The promising safety outcomes in the first-in-human study with CDNF encourage its further clinical development. It will be most interesting to see quantitative data on the clinical efficacy of intraputamenally delivered CDNF in PD patients. Our Studies I and II provide support for the clinical development of CDNF and MANF. The main outcomes in Study I suggest that NTFs have diverging effects on dopamine release and metabolizing enzymes in the normal rat brain. It was promising to see that CDNF and MANF did not decrease dopaminergic neurotransmission in the striatum. Thus, they would be most probably compatible with the current dopaminergic medications of PD. Another important finding was the COMT activity increasing effect of GDNF which calls for

attention in possible clinical trials in the future in terms of the compatibility of GDNF therapy with the existing medications of patients, in particularly with COMT-inhibitors. More extensive preclinical experiments determining the effects of NTFs on dopamine neurochemistry in a time-dependent manner should be performed. In addition, their coadministration together with standard symptomatic PD medications should be studied in animal models of PD to support the clinical development.

NTFs are protein therapeutics which complicates their clinical use and restricts them to more advanced disease conditions due to the need for intracranial administration. The initiation of the treatment at a later stage of the disease can potentially reduce their neuroprotective potential. To overcome these limitations, we have developed small molecule NTF mimetics that could be suitable for peripheral administration at an earlier phase of the disease. Indeed, BT13 and BT44 seemed to induce functional recovery in one commonly used animal model of PD when infused into the brain at the time or few weeks after the neurotoxin lesion (Studies III and IV). These RET signaling activating compounds serve as promising leads that deserve further development. First, their pharmacochemical properties should be improved to enable better solubility into aqueous vehicles and longer half-life in the brain. Secondly, their potency should be increased in order to avoid need for micromolar concentrations, and consequently reduce the risk for unspecific off-target effects. Higher efficacy would also be desirable in order to better reveal their neuroprotective potential. Lastly, the mechanism of direct RET agonism reduces their specificity and increases the risk for antagonized endogenous GFL-GFR $\alpha$  signaling. Therefore, the development of the next generation compounds towards GFR $\alpha$  PAMs would most likely improve their specificity and safety profile.

With the improved GFL-mimicking drug candidates, the neuroprotective efficacy after peripheral administration should be tested in proper dose-response studies using several well-validated PD models to gain a more comprehensive picture of their effects on PD pathology. A critical question that remains is whether they would work in  $\alpha$ -syn-based PD models where Nurr1 and RET may be downregulated (Decressac et al. 2012). As we learned from Studies III and IV, the behavioral read-outs did not correlate optimally with the histological read-outs measuring the integrity of the nigrostriatal dopamine system. Therefore, more attention should be paid to validate the animal models and behavioral assessments properly in pilot experiments before applying them into large-scale *in vivo* studies.

## 7 CONCLUSIONS

The main objective of this work was to clarify the functional effects of intracranially delivered GDNF, CDNF and MANF on the nigrostriatal dopamine system in the normal rat brain and to examine the neuroprotective potential of novel small molecule RET agonists in a 6-OHDA rat model of PD. Generally, the results support the preclinical characterization of CDNF, MANF and RET agonists as potential neurotrophic therapies for PD.

The principal findings and conclusions of this thesis are:

1. Intrastriatal injection of MANF enhances stimulus-evoked dopamine release and DOPAC/dopamine turnover in the striatum of normal freely-moving rats. GDNF-injection, instead, increases *in vivo* TH and COMT activity and decreases MAO-A activity in the striatum of normal rats. These divergent and long-lasting effects of exogenously administered NTFs on dopamine neurochemistry are noteworthy when considering their compatibility of with the existing dopaminergic medications of the patients.
2. CDNF readily diffuses in the brainstem but is not transported to distal brain areas after an injection to the SN of intact rats. Nigrally injected CDNF also colocalizes with TH-ir, but not with PV-ir, neurons in the SN. These data suggest dopaminergic selectivity of CDNF and facilitate the selection of optimal delivery site and protocol for CDNF.
3. Small molecule RET agonists BT13 and BT44 induce the phosphorylation of RET and activate the downstream pro-survival signaling pathways Akt and ERK. They promote the survival of cultured midbrain dopamine neurons in a RET-dependent manner. BT13 stimulates striatal dopamine release *in vivo* and improves functional recovery in a 6-OHDA rat model of PD. BT44 alleviates amphetamine-induced motor dysfunction in 6-OHDA lesioned rats and shows potential to restore striatal TH-ir fibers. Unlike GFLs, BT13 and BT44 are able to cross the BBB. They serve as promising lead compounds but require further drug design and development in terms of their efficacy and pharmacochemical properties.

Taken together, these results shed light on the functional effects and distribution properties of exogenously administered NTFs on the nigrostriatal dopamine system in the intact rat brain. This work also provides a proof-of-principle for small molecule RET agonists mimicking the neurotrophic effects of GFLs in an animal model of PD.



## ACKNOWLEDGEMENTS

This work was done at the Division of Pharmacology and Pharmacotherapy, Faculty of Pharmacy, University of Helsinki in collaboration with the Institute of Biotechnology, University of Helsinki during the years 2012-2020.

The work was funded by the Academy of Finland, Tekes (3iRegeneration), the Parkinson's UK and the Michael J. Fox Foundation for Parkinson's Research, each of which is sincerely acknowledged for making this research possible. I also want to acknowledge the Finnish Parkinson Foundation, Orion Research Foundation, the Finnish Pharmacological Society, the Foundation for Drug Research and the Finnish Pharmacists' Association for the personal financial support.

I wish to express my deepest gratitude to the following persons:

My supervisor professor Raimo Tuominen, the Head of the Division of Pharmacology and Pharmacotherapy, for providing invaluable guidance into the scientific world and neuropharmacological research. Thanks to his inspiring pharmacology lectures I got the spark to pursue a scientific career in the field of neuropharmacology. The freedom that he gave me in carrying out the experiments and communicating the results allowed me to mature into an independent researcher. I sincerely appreciate his openness and optimism that helped me to complete this work.

My second supervisor docent Merja Voutilainen for mentoring and encouraging me throughout these years. With her practical and scientific guidance, I was able to get started with my experiments. Her expertise in neurotrophic factor research and pharmacological *in vivo* methods had a great impact on completing the projects. I am also grateful for her comments and advices during the process of writing this thesis.

The reviewers of my thesis, professor (hc) Seppo Kaakkola and docent Sanna Janhunen, for their constructive suggestions that helped me to substantially improve the thesis manuscript, and professor Edgar Kramer for agreeing to act as my opponent in the public defense of this thesis.

All my coauthors, especially Arun Mahato, Tanel Visnapuu and Katrina Albert, for the fruitful cooperation. Without their efforts this work would have remained unrealized. I want to express my special thanks to Dr. Susanne Bäck who supervised my MSc thesis and introduced me to the world of animal experiments, and docent Petteri Piepponen who was always available to provide precious guidance on biostatistics and neurochemical research methods. I warmly thank professor Mart Saarma and docent Yulia Sidorova, from the Institute of Biotechnology, for the close collaboration and their profound knowledge of neurotrophic factor research. They were the key persons enabling the studies with the innovative RET activating compounds.

All my awesome colleagues at the Division for creating such a pleasant, supportive and well-functioning working environment. In particular, I am thankful to Jaakko Kopra and Timo Myöhänen

for their friendship and all the scientific and not-so-scientific discussions that have been enlightening. I will have fond memories of the fun times spent together.

Laboratory technicians Kati Rautio, Marjo Vaha and Liisa Konttinen for the most valuable technical assistance. Without their help and supportive attitude, the diverse challenges of laboratory work would have been difficult to overcome.

My MSc students Reinis Svarcbahts, Arttu Kontti, Charlotte Zeitler, Jette Sakki and Laura Bravo Burguillos who I was privileged to supervise. I am grateful for their contributions, as well as for the opportunity to enrich my own learning process when supervising them.

The collaborators in the projects that were beyond this thesis: docents Mikko Airavaara and Andrii Domanskyi at the Institute of Biotechnology, professor Hélder Santos at the Division of Pharmaceutical Chemistry and Technology and docent Claudio Rivera in the Neuroscience Center. These projects taught me a lot. My deep gratitude also belongs to my supervisors at Roche Innovation Center Basel, Drs. Philipp Schönenberger, Markus Britschgi and Roger Redondo, and all the other amazing Roche colleagues for acquainting me with the practices of pharmaceutical industry, mentoring me and providing me plenty of inspiration. My two internship periods in Switzerland broadened my scientific perspective and offered some of the best times of my life.

All the wonderful colleague pharmacists in the Pharmacy of Pakila for providing the most comfortable and flexible environment to keep up pharmacy work alongside my doctoral studies. Working with them was truly a pleasure.

My long-standing friends outside the lab, Teemu, Joonas, Jere, Henri, Mikko, Justus and Lassi, for all the unforgettable moments we have shared together and for counterbalancing my work-to-life ratio over these years. I am also grateful to my ice hockey friends in Tavastia hockey club for providing such great circumstances to continue with my beloved sport. Catching pucks on ice and laughing in the locker room was the perfect way to forget the hardships of the research work.

I owe my warmest gratitude to my family for their support and love throughout the long journey of education. I am grateful to my dear parents Riitta and Jaakko for raising me hard-working and teaching me the value of education. I am also thankful to my brothers Jani-Petteri and Jari-Pekka and my sister Jaana for being such good role models and showing how one can achieve anything with determination and good manners. My special thanks go to my goddaughter Fanni-Sofia who draw the excellent rat picture used in Figure 2.11.

Last but not least, I want to thank Sara for her endless love and understanding. She has always been there to encourage me over the toughest times and given me precious peer support. I also want to thank Sara's family for their hospitality, caring and joy of life.

*J-M Renko*

Sant Cugat del Vallès, November 2020

# REFERENCES

- Aarsland D, Brønne K, Larsen JP, Tysnes OB, et al., 2009. Cognitive impairment in incident, untreated Parkinson disease. *Neurology* 72, 1121–1126.
- Abdelmotilib H, Maltbie T, Delic V, Liu Z, et al., 2017.  $\alpha$ -Synuclein fibril-induced inclusion spread in rats and mice correlates with dopaminergic Neurodegeneration. *Neurobiol Dis* 105, 84–98.
- Abeliovich A, Schmitz Y, Fariñas I, Choi-Lundberg D, et al., 2000. Mice Lacking  $\alpha$ -Synuclein Display Functional Deficits in the Nigrostriatal Dopamine System. *Neuron* 25, 239–252.
- Airaksinen MS, and Saarma M, 2002. The GDNF Family: Signalling, Biological Functions and Therapeutic Value. *Nat Rev Neurosci* 3, 383–394.
- Airaksinen MS, Titievsky A, and Saarma M, 1999. GDNF Family Neurotrophic Factor Signaling: Four Masters, One Servant? *Mol Cell Neurosci* 13, 313–325.
- Airavaara M, Chiocco MJ, Howard DB, Zuchowski KL, et al., 2010. Widespread cortical expression of MANF by AAV serotype 7: Localization and protection against ischemic brain injury. *Exp Neurol* 225, 104–113.
- Airavaara M, Harvey BK, Voutilainen MH, Shen H, et al., 2012. CDNF Protects the Nigrostriatal Dopamine System and Promotes Recovery After MPTP Treatment in Mice. *Cell Transplant* 21, 1213–1223.
- Airavaara M, Mijatovic J, Vihavainen T, Piepponen TP, et al., 2006. In heterozygous GDNF knockout mice the response of striatal dopaminergic system to acute morphine is altered. *Synapse* 59, 321–329.
- Airavaara M, Shen H, Kuo C-C, Peränen J, et al., 2009. Mesencephalic astrocyte-derived neurotrophic factor reduces ischemic brain injury and promotes behavioral recovery in rats. *J Comp Neurol* 515, 116–124.
- Albert K, Renko J-M, Mätlä K, Airavaara M, et al., 2019. Cerebral Dopamine Neurotrophic Factor Diffuses Around the Brainstem and Does Not Undergo Anterograde Transport After Injection to the Substantia Nigra. *Front Neurosci* 13.
- Albin RL, Young AB, and Penney JB, 1989. The functional anatomy of basal ganglia disorders. *Trends Neurosci* 12, 366–375.
- Altermann RL, Tagliati M, and Olanow CW, 2011. Open-label surgical trials for Parkinson disease: Time for reconsideration. *Ann Neurol* 70, 5–8.
- Alvarez-Erviti L, Rodriguez-Oroz MC, Cooper JM, Caballero C, et al., 2010. Chaperone-Mediated Autophagy Markers in Parkinson Disease Brains. *Arch Neurol* 67, 1464–1472.
- Anders J, Kjær S, and Ibáñez CF, 2001. Molecular Modeling of the Extracellular Domain of the RET Receptor Tyrosine Kinase Reveals Multiple Cadherin-like Domains and a Calcium-binding Site. *J Biol Chem* 276, 35808–35817.
- Anderson DW, Neavin T, Smith JA, and Schneider JS, 2001. Neuroprotective effects of pramipexole in young and aged MPTP-treated mice. *Brain Res* 905, 44–53.
- Andres R, Forgie A, Wyatt S, Chen Q, et al., 2001. Multiple effects of artemin on sympathetic neurone generation, survival and growth. *Development* 128, 3685–3695.
- Angelova PR, Ludtmann MHR, Horrocks MH, Negoda A, et al., 2016.  $Ca^{2+}$  is a key factor in  $\alpha$ -synuclein-induced neurotoxicity. *J Cell Sci* 129.
- Aosaki T, Kiuchi K, and Kawaguchi Y, 1998. Dopamine D1-Like Receptor Activation Excites Rat Striatal Large Spiny Neurons In Vitro. *J Neurosci* 18, 5180–5190.
- Apostolou A, Shen Y, Liang Y, Luo J, et al., 2008. Armet, a UPR-upregulated protein, inhibits cell proliferation and ER stress-induced cell death. *Exp Cell Res* 314, 2454–2467.
- Appel E, Kolman O, Kazimirsky G, Blumberg PM, et al., 1997. Regulation of GDNF expression in cultured astrocytes by inflammatory stimuli. *Neuroreport* 8, 3309–3312.
- Arenas E, Trupp M, Akerud P, and Ibáñez CF, 1995. GDNF prevents degeneration and promotes the phenotype of brain noradrenergic neurons in vivo. *Neuron* 15, 1465–73.
- Arighi E, Borrello MG, and Sariola H, 2005. RET tyrosine kinase signaling in development and cancer. *Cytokine Growth Factor Rev* 16, 441–467.
- Armstrong MJ, and Okun MS, 2020. Diagnosis and Treatment of Parkinson Disease: A Review. *JAMA* 323, 548–560.
- Aron AR, and Poldrack RA, 2006. Cortical and Subcortical Contributions to Stop Signal Response Inhibition: Role of the Subthalamic Nucleus. *J Neurosci* 26, 2424–2433.
- Aron L, and Klein R, 2011. Repairing the parkinsonian brain with neurotrophic factors. *Trends Neurosci* 34, 88–100.
- Asakawa T, Fang H, Sugiyama K, Nozaki T, et al., 2016. Animal behavioral assessments in current research of Parkinson's disease. *Neurosci Biobehav Rev* 65, 63–94.
- Ascherio A, and Schwarzschild MA, 2016. The epidemiology of Parkinson's disease: risk factors and prevention. *Lancet Neurol*.
- Atik A, Stewart T, and Zhang J, 2016. Alpha-Synuclein as a Biomarker for Parkinson's Disease. *Brain Pathol* 26, 410–418.
- Back S, Peränen J, Galli E, Pulkila P, et al., 2013. Gene therapy with AAV2-CDNF provides functional benefits in a rat model of Parkinson's disease. *Brain Behav* 3.
- Baker M, Belachew SM, Gossens C, and Lindemann M, 2019. Digital biomarkers for cognition and movement diseases or disorders. U.S. Patent Application No. 16/299,869.
- Baloh RH, Gorodinsky A, Golden JP, Tansey MG, et al., 1998. GFR $\alpha$ 3 is an orphan member of the GDNF/neurturin/persephin receptor family. *Proc Natl Acad Sci U S A* 95, 5801–6.
- Baloh RH, Tansey MG, Golden JP, Creedon DJ, et al., 1997a. TrnR2, a Novel Receptor That Mediates Neurturin and GDNF Signaling through Ret. *Neuron* 18, 793–802.
- Baloh RH, Tansey MG, Lampe PA, Fahrner TJ, et al., 1998. Artemin, a novel member of the GDNF ligand family, supports peripheral and

- central neurons and signals through the GFR $\alpha$ 3-RET receptor complex. *Neuron* 21, 1291–1302.
- Bar-Am O, Weinreb O, Amit T, and Youdim MBH, 2005. Regulation of Bcl-2 family proteins, neurotrophic factors, and APP processing in the neurorescue activity of propargylamine. *FASEB J* 19, 1899–1901.
- Barraud Q, Obeid I, Aubert I, Barrière G, et al., 2010. Neuroanatomical study of the A11 diencephalospinal pathway in the non-human primate. *PLoS One* 5, e13306–e13306.
- Barrett PJ, and Timothy Greenamyre J, 2015. Post-translational modification of  $\alpha$ -synuclein in Parkinson's disease. *Brain Res* 1628, 247–253.
- Bartus RT, 2015. Gene therapy for Parkinson's disease: a decade of progress supported by posthumous contributions from volunteer subjects. *Neural Regen Res* 10, 1586–1588.
- Bartus RT, Baumann TL, Siffert J, Herzog CD, et al., 2013. Safety/feasibility of targeting the substantia nigra with AAV2-neurturin in Parkinson patients. *Neurology* 80, 1698–1701.
- Bartus RT, Herzog CD, Chu Y, Wilson A, et al., 2011. Bioactivity of AAV2-neurturin gene therapy (CERE-120): Differences between Parkinson's disease and nonhuman primate brains. *Mov Disord* 26, 27–36.
- Bartus RT, Kordower JH, Johnson EM, Brown L, et al., 2015. Post-mortem assessment of the short and long-term effects of the trophic factor neurturin in patients with  $\alpha$ -synucleinopathies. *Neurobiol Dis* 78, 162–171.
- Batchelor PE, Liberatore GT, Wong JYF, Porritt MJ, et al., 1999. Activated Macrophages and Microglia Induce Dopaminergic Sprouting in the Injured Striatum and Express Brain-Derived Neurotrophic Factor and Glial Cell Line-Derived Neurotrophic Factor. *J Neurosci* 19, 1708–1716.
- Baudet C, Mikaelis A, Westphal H, Johansen J, et al., 2000. Positive and negative interactions of GDNF, NTN and ART in developing sensory neuron subpopulations, and their collaboration with neurotrophins. *Development* 127, 4335–4344.
- Baudet C, Pozas E, Adameyko I, Andersson E, et al., 2008. Retrograde Signaling onto Ret during Motor Nerve Terminal Maturation. *J Neurosci* 28, 963–975.
- Beach TG, Adler CH, Sue LI, Vedders L, et al., 2010. Multi-organ distribution of phosphorylated  $\alpha$ -synuclein histopathology in subjects with Lewy body disorders. *Acta Neuropathol* 119, 689–702.
- Beck KD, Irwin I, Valverde J, Brennan TJ, et al., 1996. GDNF induces a dystonia-like state in neonatal rats and stimulates dopamine and serotonin synthesis. *Neuron* 16, 665–673.
- Bedford L, Hay D, Devoy A, Paine S, et al., 2008. Depletion of 26S Proteasomes in Mouse Brain Neurons Causes Neurodegeneration and Lewy-Like Inclusions Resembling Human Pale Bodies. *J Neurosci* 28, 8189–8198.
- Bellou V, Belbasis L, Tzoulaki I, Evangelou E, et al., 2016. Environmental risk factors and Parkinson's disease: An umbrella review of meta-analyses. *Parkinsonism Relat Disord* 23, 1–9.
- Benamer HTS, Patterson J, Wyper DJ, Hadley DM, et al., 2000. Correlation of Parkinson's disease severity and duration with 123I-FP-CIT SPECT striatal uptake. *Mov Disord* 15, 692–698.
- Bennett DLH, Boucher TJ, Armanini MP, Poulsen KT, et al., 2000. The Glial Cell Line-Derived Neurotrophic Factor Family Receptor Components Are Differentially Regulated within Sensory Neurons after Nerve Injury. *J Neurosci* 20, 427–437.
- Bentea E, Verbruggen L, and Massie A, 2017. The Proteasome Inhibition Model of Parkinson's Disease. *J Parkinsons Dis*.
- Benveniste H, and Hansen AJ, 1991. Practical aspects of using microdialysis for determination of brain interstitial concentrations, in: Robinson, T.E., Justice Jr., J.B. (Eds.), *Microdialysis in the Neurosciences*. Elsevier Science Publishers BV, Amsterdam, pp. 81–100.
- Berg D, Postuma RB, Adler CH, Bloem BR, et al., 2015. MDS research criteria for prodromal Parkinson's disease. *Mov Disord*.
- Berg D, Postuma RB, Bloem B, Chan P, et al., 2014. Time to redefine PD? Introductory statement of the MDS Task Force on the definition of Parkinson's disease. *Mov Disord* 29.
- Bernard V, Normand E, and Bloch B, 1992. Phenotypical characterization of the rat striatal neurons expressing muscarinic receptor genes. *J Neurosci* 12, 3591–3600.
- Bespalov Maxim M., Sidorova Y a., Tumova S, Ahonen-Bishopp A, et al., 2011a. Heparan sulfate proteoglycan syndecan-3 is a novel receptor for GDNF, neurturin, and artemin. *J Cell Biol* 192, 153–169.
- Bespalov MM, Sidorova YA, Suleymanova I, Thompson J, et al., 2016. Novel agonist of GDNF family ligand receptor RET for the treatment of experimental neuropathy. *bioRxiv*.
- Besset V, Scott RP, and Ibáñez CF, 2000. Signaling complexes and protein-protein interactions involved in the activation of the Ras and phosphatidylinositol 3-kinase pathways by the c-Ret receptor tyrosine kinase. *J Biol Chem* 275, 39159–39166.
- Betarbet R, Sherer TB, MacKenzie G, Garcia-Osuna M, et al., 2000. Chronic systemic pesticide exposure reproduces features of Parkinson's disease. *Nat Neurosci* 3, 1301–1306.
- Bevan MD, Booth PAC, Eaton SA, and Bolam JP, 1998. Selective Innervation of Neostriatal Interneurons by a Subclass of Neuron in the Globus Pallidus of the Rat. *J Neurosci* 18, 9438–9452.
- Bezard E, and Przedborski S, 2011. A tale on animal models of Parkinson's disease. *Mov Disord* 26, 993–1002.
- Bhattacharyya A, Watson FL, Pomeroy SL, Zhang YZ, et al., 2002. High-resolution imaging demonstrates dynein-based vesicular transport of activated Trk receptors. *J Neurobiol* 51, 302–312.
- Bilak MM, Shifrin DA, Corse AM, Bilak SR, et al., 1999. Neuroprotective Utility and Neurotrophic Action of Neurturin in Postnatal Motor Neurons: Comparison with GDNF and Persephin. *Mol Cell Neurosci* 13, 326–336.
- Birkmayer W, and Hornykiewicz O, 1961. Der L-Dioxyphenylalanin-Effekt bei der Parkinson-Kinese. *Wien Klin Wochenschr* 73, 787–788.
- Bizon JL, Lauterborn JC, and Gall CM, 1999. Subpopulations of striatal interneurons can be distinguished on the basis of neurotrophic factor expression. *J Comp Neurol* 408, 283–298.
- Björklund A, and Dunnett SB, 2019. The Amphetamine Induced Rotation Test: A Re-Assessment of Its Use as a Tool to Monitor Motor Impairment and Functional Recovery in Rodent Models of Parkinson's Disease. *J Parkinsons Dis* 9, 17–29.
- Björklund A, and Dunnett SB, 2007. Dopamine neuron systems in the brain: an update. *Trends Neurosci* 30, 194–202.
- Björklund A, and Nolin A, 1973. Fluorescence histochemical and microspectrofluorometric mapping of dopamine and noradrenaline cell

- groups in the rat diencephalon. *Brain Res* 51, 193–205.
- Björklund A, Rosenblad C, Winkler C, and Kirik D, 1997. Studies on Neuroprotective and Regenerative Effects of GDNF in a Partial Lesion Model of Parkinson's Disease. *Neurobiol Dis* 4, 186–200.
- Blanchard V, Anglade P, Dziewczapolski G, Savasta M, et al., 1996. Dopaminergic sprouting in the rat striatum after partial lesion of the substantia nigra. *Brain Res* 709, 319–325.
- Blandini F, Armentero MT, Fancello R, Blaugrund E, et al., 2004. Neuroprotective effect of rasagiline in a rodent model of Parkinson's disease. *Exp Neurol* 187, 455–459.
- Blauwendraat C, Nalls MA, and Singleton AB, 2020. The genetic architecture of Parkinson's disease. *Lancet Neurol* 19, 170–178.
- Blesa J, and Przedborski S, 2014. Parkinson's disease: animal models and dopaminergic cell vulnerability. *Front Neuroanat* 8, 1–12.
- Bloch A, Probst A, Bissig H, Adams H, et al., 2006.  $\alpha$ -Synuclein pathology of the spinal and peripheral autonomic nervous system in neurologically unimpaired elderly subjects. *Neuropathol Appl Neurobiol* 32, 284–295.
- Block ML, and Hong J-S, 2005. Microglia and inflammation-mediated neurodegeneration: Multiple triggers with a common mechanism. *Prog Neurobiol* 76, 77–98.
- Bobo RH, Laske DW, Akbasak A, Morrison PF, et al., 1994. Convection-enhanced delivery of macromolecules in the brain. *Proc Natl Acad Sci* 91, 2076–2080.
- Boger H a, Middaugh LD, Patrick KS, Ramamoorthy S, et al., 2007. Long-term consequences of methamphetamine exposure in young adults are exacerbated in glial cell line-derived neurotrophic factor heterozygous mice. *J Neurosci* 27, 8816–8825.
- Bootcov MR, Bauskin AR, Valenzuela SM, Moore AG, et al., 1997. MIC-1, a novel macrophage inhibitory cytokine, is a divergent member of the TGF- $\beta$  superfamily. *Proc Natl Acad Sci* 94, 11514–11519.
- Bothwell M, 2014. NGF, BDNF, NT3, and NT4. Springer, Berlin, Heidelberg, pp. 3–15.
- Böttner M, Suter-Crazzolara C, Schober A, and Unsicker K, 1999. Expression of a novel member of the TGF- $\beta$  superfamily, growth/differentiation factor-15/macrophage-inhibiting cytokine-1 (GDF-15/MIC-1) in adult rat tissues. *Cell Tissue Res* 297, 103–110.
- Bourque MJ, and Trudeau LE, 2000. GDNF enhances the synaptic efficacy of dopaminergic neurons in culture. *Eur J Neurosci* 12, 3172–3180.
- Bowenkamp KE, Hoffman AF, Gerhardt GA, Henry MA, et al., 1995. Glial cell line-derived neurotrophic factor supports survival of injured midbrain dopaminergic neurons. *J Comp Neurol* 355, 479–489.
- Braak H, de Vos RAI, Bohl J, and Del Tredici K, 2006. Gastric alpha-synuclein immunoreactive inclusions in Meissner's and Auerbach's plexuses in cases staged for Parkinson's disease-related brain pathology. *Neurosci Lett* 396, 67–72.
- Braak H, Ghebremedhin E, Rüb U, Bratzke H, et al., 2004. Stages in the development of Parkinson's disease-related pathology. *Cell Tissue Res* 318, 121–134.
- Braak H, Rüb U, Gai WP, and Del Tredici K, 2003a. Idiopathic Parkinson's disease: possible routes by which vulnerable neuronal types may be subject to neuroinvasion by an unknown pathogen. *J Neural Transm* 110, 517–536.
- Braak H, Tredici K Del, Rüb U, de Vos RA., et al., 2003b. Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol Aging* 24, 197–211.
- Bracci E, Centonze D, Bernardi G, and Calabresi P, 2002. Dopamine Excites Fast-Spiking Interneurons in the Striatum. *J Neurophysiol* 87, 2190–2194.
- Bresjanac M, and Antauer G, 2000. Reactive Astrocytes of the Quinolinic Acid-Lesioned Rat Striatum Express GFR $\alpha$ 1 as Well as GDNF in Vivo. *Exp Neurol* 164, 53–59.
- Brimblecombe KR, and Cragg SJ, 2017. The Striosome and Matrix Compartments of the Striatum: A Path through the Labyrinth from Neurochemistry toward Function. *ACS Chem Neurosci* 8, 235–242.
- Brizard M, Carcenac C, Bemelmans A-P, Feuerstein C, et al., 2006. Functional reinnervation from remaining DA terminals induced by GDNF lentivirus in a rat model of early Parkinson's disease. *Neurobiol Dis* 21, 90–101.
- Bromek E, Haduch A, Golembiowska K, and Daniel WA, 2011. Cytochrome P450 mediates dopamine formation in the brain in vivo. *J Neurochem* 118, 806–815.
- Bronfman FC, Escudero CA, Weis J, and Kruttgen A, 2007. Endosomal transport of neurotrophins: Roles in signaling and neurodegenerative diseases. *Dev Neurobiol* 67, 1183–1203.
- Brown RSE, Herbison AE, and Grattan DR, 2015. Effects of Prolactin and Lactation on A15 Dopamine Neurons in the Rostral Preoptic Area of Female Mice. *J Neuroendocrinol* 27, 708–717.
- Brück D, Wenning GK, Stefanova N, and Fellner L, 2016. Glia and alpha-synuclein in neurodegeneration: A complex interaction. *Neurobiol Dis* 85, 262–274.
- Brücke T, Asenbaum S, Pirker W, Djamshidian S, et al., 1997. Measurement of the dopaminergic degeneration in Parkinson's disease with [123I] $\beta$ -CIT and SPECT BT - Advances in Research on Neurodegeneration, in: Riederer, P., Calne, D.B., Horowski, R., Mizuno, Y., Poewe, W., Youdim, M.B.H. (Eds.), . Springer Vienna, Vienna, pp. 9–24.
- Bruhn H, 2005. A short guided tour through functional and structural features of saposin-like proteins, *Biochem. J.*
- Brundin P, Ma J, and Kordower JH, 2016. How strong is the evidence that Parkinson's disease is a prion disorder? *Curr Opin Neurol.*
- Buj-Bello A, Adu J, Piñón LGP, Horton A, et al., 1997. Neurturin responsiveness requires a GPI-linked receptor and the Ret receptor tyrosine kinase. *Nature* 387, 721–724.
- Buj-Bello A, Buchman VL, Horton A, Rosenthal A, et al., 1995a. GDNF is an age-specific survival factor for sensory and autonomic neurons. *Neuron* 15, 821–828.
- Burke RE, Antonelli M, and Sulzer D, 1998. Glial Cell Line-Derived Neurotrophic Growth Factor Inhibits Apoptotic Death of Postnatal Substantia Nigra Dopamine Neurons in Primary Culture. *J Neurochem* 71, 517–525.
- Burke RE, and O'Malley K, 2013. Axon degeneration in Parkinson's disease. *Exp Neurol* 246, 72–83.
- Burns RS, Chiueh CC, Markey SP, Ebert MH, et al., 1983. A primate model of parkinsonism: selective destruction of dopaminergic neurons in the pars compacta of the substantia nigra by N-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. *Proc Natl Acad Sci* 80, 4546–4550.

- Burré J, Sharma M, Tsetsenis T, Buchman V, et al., 2010.  $\alpha$ -Synuclein Promotes SNARE-Complex Assembly in Vivo and in Vitro. *Science* (80-) 329, 1663–1667.
- Burré J, Vivona S, Diao J, Sharma M, et al., 2013. Properties of native brain  $\alpha$ -synuclein. *Nature* 498, E4–E6.
- Butowt R, and von Bartheld CS, 2007. Conventional kinesin-I motors participate in the anterograde axonal transport of neurotrophins in the visual system. *J Neurosci Res* 85, 2546–2556.
- Butowt R, and von Bartheld CS, 2001. Sorting of Internalized Neurotrophins into an Endocytic Transcytosis Pathway via the Golgi System: Ultrastructural Analysis in Retinal Ganglion Cells. *J Neurosci* 21, 8915 LP – 8930.
- Buu NT, Duhaime J, Savard C, Truong L, et al., 1981. Presence of Conjugated Catecholamines in Rat Brain: A New Method of Analysis of Catecholamine Sulfates. *J Neurochem* 36, 769–772.
- Bäckman CM, Shan L, Zhang YJ, Hoffer BJ, et al., 2006. Gene expression patterns for GDNF and its receptors in the human putamen affected by Parkinson's disease: A real-time PCR study. *Mol Cell Endocrinol* 252, 160–166.
- Cabin DE, Shimazu K, Murphy D, Cole NB, et al., 2002. Synaptic Vesicle Depletion Correlates with Attenuated Synaptic Responses to Prolonged Repetitive Stimulation in Mice Lacking  $\alpha$ -Synuclein. *J Neurosci* 22, 8797–8807.
- Cacalano G, Fariñas I, Wang L-C, Hagler K, et al., 1998. GFR $\alpha$ 1 Is an Essential Receptor Component for GDNF in the Developing Nervous System and Kidney. *Neuron* 21, 53–62.
- Cachope R, Mateo Y, Mathur BN, Irving J, et al., 2012. Selective Activation of Cholinergic Interneurons Enhances Accumbal Phasic Dopamine Release: Setting the Tone for Reward Processing. *Cell Rep* 2, 33–41.
- Cai Y, and Ford CP, 2018. Dopamine Cells Differentially Regulate Striatal Cholinergic Transmission across Regions through Corelease of Dopamine and Glutamate. *Cell Rep* 25, 3148-3157.e3.
- Caillé I, Dumartin B, and Bloch B, 1996. Ultrastructural localization of D1 dopamine receptor immunoreactivity in rat striatonigral neurons and its relation with dopaminergic innervation. *Brain Res* 730, 17–31.
- Calabresi P, Centonze D, Pisani A, Sancesario G, et al., 1998. Muscarinic IPSPs in rat striatal cholinergic interneurons. *J Physiol* 510, 421–427.
- Calabresi P, Picconi B, Tozzi A, and Filippo M Di, 2007. Dopamine-mediated regulation of corticostriatal synaptic plasticity. *Trends Neurosci* 30, 211–219.
- Cao J-P, Wang H-J, Yu J-K, Yang H, et al., 2008. Involvement of NCAM in the effects of GDNF on the neurite outgrowth in the dopamine neurons. *Neurosci Res* 61, 390–397.
- Cao SS, Luo KL, and Shi L, 2016. Endoplasmic Reticulum Stress Interacts With Inflammation in Human Diseases. *J Cell Physiol* 231, 288–294.
- Cardozo DL, and Bean BP, 1995. Voltage-dependent calcium channels in rat midbrain dopamine neurons: modulation by dopamine and GABAB receptors. *J Neurophysiol* 74, 1137–1148.
- Carlsson A, 2001. A Half-Century of Neurotransmitter Research: Impact on Neurology and Psychiatry (Nobel Lecture). *ChemBioChem* 2, 484–493.
- Carlsson A, 1959. The occurrence, distribution and physiological role of catecholamines in the nervous system. *Pharmacol Rev* 11, 490–493.
- Carlsson A, Davis JN, Kehr W, Lindqvist M, et al., 1972. Simultaneous measurement of tyrosine and tryptophan hydroxylase activities in brain in Vivo using an inhibitor of the aromatic amino acid decarboxylase. *Naunyn Schmiedeberg's Arch Pharmacol* 275, 153–168.
- Carlsson A, Lindqvist M, and Magnusson T, 1957. 3,4-Dihydroxyphenylalanine and 5-Hydroxytryptophan as Reserpine Antagonists. *Nature* 180, 1200.
- Carlsson A, Lindqvist M, Magnusson T, and Waldeck B, 1958. On the Presence of 3-Hydroxytyramine in Brain. *Science* (80-) 127, 471.
- Carniti C, Perego C, Mondellini P, Pierotti MA, et al., 2003. PP1 Inhibitor Induces Degradation of RETMEN2A and RETMEN2B Oncoproteins through Proteosomal Targeting. *Cancer Res* 63, 2234–2243.
- Carter MT, Yome JL, Marcil MN, Martin CA, et al., 2001. Conservation of RET proto-oncogene splicing variants and implications for RET isoform function. *Cytogenet Genome Res* 95, 169–176.
- Cass WA, and Peters LE, 2010. Neurturin effects on nigrostriatal dopamine release and content: Comparison with GDNF. *Neurochem Res* 35, 727–734.
- Cass WA, Walker DJ, and Manning MW, 1999. Augmented methamphetamine-induced overflow of striatal dopamine 1 day after GDNF administration. *Brain Res* 827, 104–112.
- Castellanos G, Fernández-Seara MA, Lorenzo-Betancor O, Ortega-Cubero S, et al., 2015. Automated Neuromelanin Imaging as a Diagnostic Biomarker for Parkinson's Disease. *Mov Disord* 30, 945–952.
- Castrén E, and Hen R, 2013. Neuronal plasticity and antidepressant actions. *Trends Neurosci* 36, 259–267.
- Caviness JN, Hentz JG, Belden CM, Shill HA, et al., 2015. Longitudinal EEG Changes Correlate with Cognitive Measure Deterioration in Parkinson's Disease. *J Parkinson's Dis* 5, 117–124.
- Centonze D, Bracci E, Pisani A, Gubellini P, et al., 2002. Activation of dopamine D1-like receptors excites LTS interneurons of the striatum. *Eur J Neurosci* 15, 2049–2052.
- Centonze D, Grande C, Usiello A, Gubellini P, et al., 2003. Receptor Subtypes Involved in the Presynaptic and Postsynaptic Actions of Dopamine on Striatal Interneurons. *J Neurosci* 23, 6245–6254.
- Cerchia L, 2006. An Autocrine Loop Involving Ret and Glial Cell-Derived Neurotrophic Factor Mediates Retinoic Acid-Induced Neuroblastoma Cell Differentiation. *Mol Cancer Res* 4, 481–488.
- Cersosimo MG, and Benarroch EE, 2012. Autonomic involvement in Parkinson's disease: Pathology, pathophysiology, clinical features and possible peripheral biomarkers. *J Neurol Sci* 313, 57–63.
- Chahine LM, and Stern MB, 2017. Parkinson's Disease Biomarkers: Where Are We and Where Do We Go Next? *Mov Disord Clin Pract* 4, 796–805.
- Chandra S, Gallardo G, Fernández-Chacón R, Schlüter OM, et al., 2005.  $\alpha$ -Synuclein Cooperates with CSP $\alpha$  in Preventing Neurodegeneration. *Cell* 123, 383–396.
- Chao CC, Ma YL, Chu KY, and Lee EH., 2003. Integrin  $\alpha$ v and NCAM mediate the effects of GDNF on DA neuron survival, outgrowth, DA

- turnover and motor activity in rats. *Neurobiol Aging* 24, 105–116.
- Charara A, Heilman C, Levey AI, and Smith Y, 1999. Pre- and postsynaptic localization of GABAB receptors in the basal ganglia in monkeys. *Neuroscience* 95, 127–140.
- Charbit AR, Akerman S, and Goadsby PJ, 2010. Dopamine: what's new in migraine? *Curr Opin Neurol* 23, 275–281.
- Charbit AR, Akerman S, Holland PR, and Goadsby PJ, 2009. Neurons of the Dopaminergic/Calcitonin Gene-Related Peptide A11 Cell Group Modulate Neuronal Firing in the Trigemino-cervical Complex: An Electrophysiological and Immunohistochemical Study. *J Neurosci* 29, 12532–12541.
- Chase TN, and Oh JD, 2000. Striatal mechanisms and pathogenesis of Parkinsonian signs and motor complications. *Ann Neurol* 47.
- Chaudhry FA, Edwards RH, and Fonnum F, 2008. Vesicular Neurotransmitter Transporters as Targets for Endogenous and Exogenous Toxic Substances. *Annu Rev Pharmacol Toxicol* 48, 277–301.
- Chaudhuri KR, and Schapira AH V, 2009. Non-motor symptoms of Parkinson's disease: dopaminergic pathophysiology and treatment. *Lancet Neurol* 8, 464–474.
- Chauhan NB, Siegel GJ, and Lee JM, 2001. Depletion of glial cell line-derived neurotrophic factor in substantia nigra neurons of Parkinson's disease brain. *J Chem Neuroanat* 21, 277–288.
- Chavarría C, Rodríguez-Bottero S, Quijano C, Cassina P, et al., 2018. Impact of monomeric, oligomeric and fibrillar alpha-synuclein on astrocyte reactivity and toxicity to neurons. *Biochem J* 475, 3153–3169.
- Chen L-W, Zhang J-P, Kwok-Yan Shum D, and Chan Y-S, 2006. Localization of nerve growth factor, neurotrophin-3, and glial cell line-derived neurotrophic factor in nestin-expressing reactive astrocytes in the caudate-putamen of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-treated C57/Bl mice. *J Comp Neurol* 497, 898–909.
- Chen L, Feng L, Wang X, Du J, et al., 2015. Mesencephalic Astrocyte-Derived Neurotrophic Factor Is Involved in Inflammation by Negatively Regulating the NF- $\kappa$ B Pathway. *Sci Rep* 5, 8133.
- Chen Y-C, Sundvik M, Rozov S, Priyadarshini M, et al., 2012. MANF regulates dopaminergic neuron development in larval zebrafish. *Dev Biol* 370, 237–249.
- Cheng H-C, Ulane CM, and Burke RE, 2010. Clinical progression in Parkinson disease and the neurobiology of axons. *Ann Neurol* 67, 715–725.
- Cheng L, Liu Y, Zhao H, Zhang W, et al., 2013a. Lentiviral-mediated transfer of CDNF promotes nerve regeneration and functional recovery after sciatic nerve injury in adult rats. *Biochem Biophys Res Commun* 440.
- Cheng L, Zhao H, Zhang W, Liu B, et al., 2013b. Overexpression of conserved dopamine neurotrophic factor (CDNF) in astrocytes alleviates endoplasmic reticulum stress-induced cell damage and inflammatory cytokine secretion. *Biochem Biophys Res Commun* 435.
- Cheng Q, Wang X, Shen Yujun, Shen Yuxian, et al., 2014. Deficiency of IRE1 and PERK Signal Pathways in Systemic Lupus Erythematosus. *Am J Med Sci* 348, 465–473.
- Cho J, Kholodilov NG, and Burke RE, 2004a. Patterns of developmental mRNA expression of neurturin and GFR $\alpha$ 2 in the rat striatum and substantia nigra do not suggest a role in the regulation of natural cell death in dopamine neurons. *Dev Brain Res* 148, 143–149.
- Cho J, Yarygina O, Oo TF, Kholodilov NG, et al., 2004b. Glial cell line-derived neurotrophic factor receptor GFR $\alpha$ 1 is expressed in the rat striatum during postnatal development. *Mol Brain Res* 127, 96–104.
- Choi-Lundberg DL, and Bohn MC, 1995. Ontogeny and distribution of glial cell line-derived neurotrophic factor (GDNF) mRNA in rat. *Dev Brain Res* 85, 80–88.
- Chou KL, Stacy M, Simuni T, Miyasaki J, et al., 2018. The spectrum of “off” in Parkinson's disease: What have we learned over 40 years? *Parkinsonism Relat Disord* 51, 9–16.
- Chu Y, Bartus RT, Manfredsson FP, Olanow CW, et al., 2020. Long-term post-mortem studies following neurturin gene therapy in patients with advanced Parkinson's disease. *Brain* 143, 960–975.
- Chu Y, Dodiya H, Aebischer P, Olanow CW, et al., 2009. Alterations in lysosomal and proteasomal markers in Parkinson's disease: Relationship to alpha-synuclein inclusions. *Neurobiol Dis* 35, 385–398.
- Chu Y, and Kordower JH, 2010. Lewy body pathology in fetal grafts. *Ann N Y Acad Sci* 1184, 55–67.
- Chu Y, Le W, Kompoliti K, Jankovic J, et al., 2006. Nurr1 in Parkinson's disease and related disorders. *J Comp Neurol* 494, 495–514.
- Chu Y, Morfini GA, Langhamer LB, He Y, et al., 2012. Alterations in axonal transport motor proteins in sporadic and experimental Parkinson's disease. *Brain* 135, 2058–2073.
- Chu Y, Muller S, Tavares A, Barret O, et al., 2019. Intrastriatal alpha-synuclein fibrils in monkeys: spreading, imaging and neuropathological changes. *Brain* 142, 3565–3579.
- Chuhma N, Mingote S, Moore H, and Rayport S, 2014. Dopamine Neurons Control Striatal Cholinergic Neurons via Regionally Heterogeneous Dopamine and Glutamate Signaling. *Neuron* 81, 901–912.
- Chuhma N, Tanaka KF, and Nagai T, 2019. The physiology and pathophysiology of basal ganglia: From signal transduction to circuits. *Neurochem Int* 131, 104544.
- Chung CY, Koprich JB, Siddiqi H, and Isacson O, 2009. Dynamic Changes in Presynaptic and Axonal Transport Proteins Combined with Striatal Neuroinflammation Precede Dopaminergic Neuronal Loss in a Rat Model of AAV  $\alpha$ -Synucleinopathy. *J Neurosci* 29, 3365–3373.
- Cilia R, Tunesi S, Marotta G, Cereda E, et al., 2016. Survival and dementia in GBA-associated Parkinson's disease: The mutation matters. *Ann Neurol* 80, 662–673.
- Clarke CE, and Guttman M, 2002. Dopamine agonist monotherapy in Parkinson's disease. *Lancet* 360, 1767–1769.
- Clarkson J, and Herbison AE, 2011. Dual Phenotype Kisspeptin-Dopamine Neurons of the Rostral Periventricular Area of the Third Ventricle Project to Gonadotrophin-Releasing Hormone Neurons. *J Neuroendocrinol* 23, 293–301.
- Clemens S, Rye D, and Hochman S, 2006. Restless legs syndrome. *Neurology* 67, 125–130.
- Cochemé HM, and Murphy MP, 2008. Complex I is the Major Site of Mitochondrial Superoxide Production by Paraquat. *J Biol Chem* 283, 1786–1798.
- Cohen AD, Tillerson JL, Smith AD, Schallert T, et al., 2003. Neuroprotective effects of prior limb use in 6-hydroxydopamine-treated rats:

- possible role of GDNF. *J Neurochem* 85, 299–305.
- Compta Y, Parkkinen L, O'Sullivan SS, Vandrovcova J, et al., 2011. Lewy- and Alzheimer-type pathologies in Parkinson's disease dementia: which is more important? *Brain* 134, 1493–1505.
- Connolly BS, and Lang AE, 2014. Pharmacological treatment of Parkinson disease: A review. *JAMA - J Am Med Assoc*.
- Constantinescu R, and Mondello S, 2013. Cerebrospinal Fluid Biomarker Candidates for Parkinsonian Disorders. *Front Neurol* 3.
- Cordero-Llana Ó, Houghton BC, Rinaldi F, Taylor H, et al., 2015a. Enhanced efficacy of the CDNF/MANF family by combined intranigral overexpression in the 6-OHDA rat model of Parkinson's disease. *Mol Ther* 23.
- Cotzias GC, Papavasiliou PS, and Gellene R, 1969. Modification of Parkinsonism — Chronic Treatment with L-Dopa. *N Engl J Med* 280, 337–345.
- Coulpier M, Anders J, and Ibáñez CF, 2002. Coordinated activation of autophosphorylation sites in the RET receptor tyrosine kinase: importance of tyrosine 1062 for GDNF mediated neuronal differentiation and survival. *J Biol Chem* 277, 1991–1999.
- Coulpier M, and Ibáñez CF, 2004. Retrograde propagation of GDNF-mediated signals in sympathetic neurons. *Mol Cell Neurosci* 27, 132–139.
- Cragg SJ, and Rice ME, 2004. DANCing past the DAT at a DA synapse. *Trends Neurosci* 27, 270–277.
- Cramer SC, Sur M, Dobkin BH, O'Brien C, et al., 2011. Harnessing neuroplasticity for clinical applications. *Brain* 134, 1591–1609.
- Crittenden JR, and Graybiel AM, 2016. Disease-Associated Changes in the Striosome and Matrix Compartments of the Dorsal Striatum, in: Steiner, H., Tseng, K.Y.B.T.-H. of B.N. (Eds.), *Handbook of Basal Ganglia Structure and Function*, Second Edition. Elsevier, pp. 783–802.
- Crupi MJF, Yoganathan P, Bone LN, Lian E, et al., 2015. Distinct Temporal Regulation of RET Isoform Internalization: Roles of Clathrin and AP2. *Traffic* 16, 1155–1173.
- da Silva JA, Tecuapetla F, Paixão V, and Costa RM, 2018. Dopamine neuron activity before action initiation gates and invigorates future movements. *Nature* 554, 244–248.
- Dahlström A, and Fuxe K, 1964. Evidence For The Existence Of Monoamine-Containing Neurons In The Central Nervous System. I. Demonstration Of Monoamines In The Cell Bodies Of Brain Stem Neurons. *Acta Physiol Scand Suppl* 232:1-55.
- Danilova T, and Lindahl M, 2018. Emerging Roles for Mesencephalic Astrocyte-Derived Neurotrophic Factor (MANF) in Pancreatic Beta Cells and Diabetes. *Front Physiol* 9, 1457.
- Danzer KM, Haasen D, Karow AR, Moussaud S, et al., 2007. Different Species of  $\alpha$ -Synuclein Oligomers Induce Calcium Influx and Seeding. *J Neurosci* 27, 9220–9232.
- Daubner SC, Le T, and Wang S, 2011. Tyrosine hydroxylase and regulation of dopamine synthesis. *Arch Biochem Biophys* 508, 1–12.
- Dauer W, and Przedborski S, 2003. Parkinson's disease: Mechanisms and models. *Neuron* 39, 889–909.
- Davies AM, 1996. The neurotrophic hypothesis: where does it stand? *Philos Trans R Soc Lond B Biol Sci* 351, 389–94.
- Dawson TM, Ko HS, and Dawson VL, 2010. Genetic Animal Models of Parkinson's Disease. *Neuron* 66, 646–661.
- de Lau LM, and Breteler MM, 2006. Epidemiology of Parkinson's disease. *Lancet Neurol* 5, 525–535.
- De Vos KJ, Grierson AJ, Ackerley S, and Miller CCJ, 2008. Role of Axonal Transport in Neurodegenerative Diseases. *Annu Rev Neurosci* 31, 151–173.
- Decressac M, Kadkhodaei B, Mattsson B, Laguna A, et al., 2012.  $\alpha$ -Synuclein-Induced Down-Regulation of Nurr1 Disrupts GDNF Signaling in Nigral Dopamine Neurons. *Sci Transl Med* 4, 163ra156.
- Decressac M, Ulusoy A, Mattsson B, Georgievska B, et al., 2011. GDNF fails to exert neuroprotection in a rat  $\alpha$ -synuclein model of Parkinson's disease. *Brain* 134, 2302–2311.
- Deng H, Wang P, and Jankovic J, 2018. The genetics of Parkinson disease. *Ageing Res Rev*.
- Deumens R, Blokland A, and Prickaerts J, 2002. Modeling Parkinson's Disease in Rats: An Evaluation of 6-OHDA Lesions of the Nigrostriatal Pathway. *Exp Neurol* 175, 303–317.
- Devi L, Raghavendran V, Prabhu BM, Avadhani NG, et al., 2008. Mitochondrial Import and Accumulation of  $\alpha$ -Synuclein Impair Complex I in Human Dopaminergic Neuronal Cultures and Parkinson Disease Brain. *J Biol Chem* 283, 9089–9100.
- Di Maio R, Barrett PJ, Hoffman EK, Barrett CW, et al., 2016.  $\alpha$ -synuclein binds to TOM20 and inhibits mitochondrial protein import in Parkinson's disease. *Sci Transl Med*.
- Di Maio R, Hoffman EK, Rocha EM, Keeney MT, et al., 2018. LRRK2 activation in idiopathic Parkinson's disease. *Sci Transl Med* 10, eaar5429.
- Dickson DW, 2012. Parkinson's Disease and Parkinsonism: Neuropathology. *Cold Spring Harb Perspect Med* 2, 1–15.
- Dickson DW, Uchikado H, Fujishiro H, and Tsuboi Y, 2010. Evidence in favor of Braak staging of Parkinson's disease. *Mov Disord* 25, S78–S82.
- Doherty KM, Silveira-Moriyama L, Parkkinen L, Healy DG, et al., 2013. Parkin Disease: A Clinicopathologic Entity? *JAMA Neurol* 70, 571–579.
- Doppler K, Volkman J, and Sommer C, 2016. Skin biopsies in the differential diagnosis of parkinsonism: are we ready for simplified protocols? *Brain* 139, e5.
- Dorsey ER, Elbaz A, Nichols E, Abd-Allah F, et al., 2018. Global, regional, and national burden of Parkinson's disease, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol* 17, 939–953.
- Døskeland AP, and Flatmark T, 2002. Ubiquitination of soluble and membrane-bound tyrosine hydroxylase and degradation of the soluble form. *Eur J Biochem* 269, 1561–1569.
- Driver JA, Logroschino G, Gaziano JM, and Kurth T, 2009. Incidence and remaining lifetime risk of Parkinson disease in advanced age. *Neurology* 72, 432–438.
- Dufey E, Sepúlveda D, Rojas-Rivera D, and Hetz C, 2014. Cellular Mechanisms of Endoplasmic Reticulum Stress Signaling in Health and Disease. 1. An overview. *Am J Physiol Physiol* 307, C582–C594.
- Duffy MF, Collier TJ, Patterson JR, Kemp CJ, et al., 2018. Lewy body-like alpha-synuclein inclusions trigger reactive microgliosis prior to nigral degeneration. *J Neuroinflammation* 15, 1–18.
- Dunkley PR, Bobrovskaya L, Graham ME, Von Nagy-Felsobuki EI, et al., 2004. Tyrosine hydroxylase phosphorylation: regulation and



- consequences. *J Neurochem* 91, 1025–1043.
- Dunnett SB, and Lelos M, 2010. Behavioral analysis of motor and non-motor symptoms in rodent models of Parkinson's disease, in: Björklund, A., Cenci, M.A. (Eds.), *Recent Advances in Parkinson's Disease*, Progress in Brain Research. Elsevier, pp. 35–51.
- Dunnett Stephen B, 2005. Chapter V Motor function(s) of the nigrostriatal dopamine system: Studies of lesions and behavior, in: Dunnett, S B, Bentivoglio, M., Björklund, A., Hökfelt, T. (Eds.), *Dopamine, Handbook of Chemical Neuroanatomy*. Elsevier, pp. 237–301.
- Durbec P, Marcos-Gutierrez C V, Kilkeny C, Grigoriou M, et al., 1996. GDNF signalling through the Ret receptor tyrosine kinase. *Nature*.
- Durbec PL, Larsson-Blomberg LB, Schuchardt A, Costantini F, et al., 1996. Common origin and developmental dependence on c-ret of subsets of enteric and sympathetic neuroblasts. *Development* 122, 349–358.
- Dzamko N, and Halliday GM, 2012. An emerging role for LRRK2 in the immune system. *Biochem Soc Trans* 40, 1134–1139.
- Ebendal T, Tomac A, Hoffer BJ, and Olson L, 1995. Glial cell line-derived neurotrophic factor stimulates fiber formation and survival in cultured neurons from peripheral autonomic ganglia. *J Neurosci Res* 40, 276–284.
- Eberling JL, Dave KD, and Frasier MA, 2013.  $\alpha$ -synuclein Imaging: A Critical Need for Parkinson's Disease Research. *J Parkinsons Dis* 3, 565–567.
- Ebrahimi-Fakhari D, Wahlster L, and McLean PJ, 2012. Protein degradation pathways in Parkinson's disease: curse or blessing. *Acta Neuropathol* 124, 153–172.
- Edison P, Ahmed I, Fan Z, Hinz R, et al., 2013. Microglia, Amyloid, and Glucose Metabolism in Parkinson's Disease with and without Dementia. *Neuropsychopharmacology* 38, 938–949.
- Eggert K, Schlegel J, Oertel W, Würz C, et al., 1999. Glial cell line-derived neurotrophic factor protects dopaminergic neurons from 6-hydroxydopamine toxicity in vitro. *Neurosci Lett* 269, 178–182.
- Ehlers MD, Kaplan DR, Price DL, and Koliatsos VE, 1995. NGF-stimulated retrograde transport of TrkA in the mammalian nervous system. *J Cell Biol* 130, 149–156.
- Ehringer H, and Hornykiewicz O, 1960. Verteilung Von Noradrenalin Und Dopamin (3-Hydroxytyramin) Im Gehirn Des Menschen Und Ihr Verhalten Bei Erkrankungen Des Extrapyramidalen Systems. *Klin Wochenschr* 38, 1236–1239.
- Eisenhofer G, 2001. The role of neuronal and extraneuronal plasma membrane transporters in the inactivation of peripheral catecholamines. *Pharmacol Ther* 91, 35–62.
- Eisenhofer G, Åneman A, Friberg P, Hooper D, et al., 1997. Substantial Production of Dopamine in the Human Gastrointestinal Tract. *J Clin Endocrinol Metab* 82, 3864–3871.
- Eisenhofer G, Kopin IJ, and Goldstein DS, 2004. Catecholamine metabolism: a contemporary view with implications for physiology and medicine. *Pharmacol Rev* 56, 331–349.
- Ekstrand MI, Terzioglu M, Galter D, Zhu S, et al., 2007. Progressive parkinsonism in mice with respiratory-chain-deficient dopamine neurons. *Proc Natl Acad Sci* 104, 1325–1330.
- Emborg ME, 2007. Nonhuman Primate Models of Parkinson's Disease. *ILAR J* 48, 339–355.
- Emborg ME, Moirano J, Raschke J, Bondarenko V, et al., 2009. Response of aged parkinsonian monkeys to in vivo gene transfer of GDNF. *Neurobiol Dis* 36, 303–311.
- Emmerson PJ, Wang F, Du Y, Liu Q, et al., 2017. The metabolic effects of GDF15 are mediated by the orphan receptor GFRAL. *Nat Med* 23, 1215–1219.
- Enokido Y, de Sauvage F, Hongo J-A, Ninkina N, et al., 1998. GFR $\alpha$ -4 and the tyrosine kinase Ret form a functional receptor complex for persephin. *Curr Biol* 8, 1019–1022.
- Enomoto H, Araki T, Jackman A, Heuckeroth RO, et al., 1998. GFR  $\alpha$ 1-deficient mice have deficits in the enteric nervous system and kidneys. *Neuron* 21, 317–24.
- Enomoto H, Crawford PA, Gorodinsky A, Heuckeroth RO, et al., 2001. RET signaling is essential for migration, axonal growth and axon guidance of developing sympathetic neurons. *Development* 128, 3963–3974.
- Enomoto H, Heuckeroth RO, Golden JP, Johnson EM, et al., 2000. Development of cranial parasympathetic ganglia requires sequential actions of GDNF and neurturin. *Development* 127, 4877–4889.
- Enterría-Morales D, López-López I, López-Barneo J, and d'Anglemont de Tassigny X, 2020. Role of glial cell line-derived neurotrophic factor in the maintenance of adult mesencephalic catecholaminergic neurons. *Mov Disord* n/a.
- Eriksson PS, Perfilieva E, Björk-Eriksson T, Alborn A-M, et al., 1998. Neurogenesis in the adult human hippocampus. *Nat Med* 4, 1313–1317.
- Ernst A, Alkass K, Bernard S, Salehpour M, et al., 2014. Neurogenesis in the striatum of the adult human brain 156, 1072–1083.
- Eslamboli A, Cummings RM, Ridley RM, Baker HF, et al., 2003. Recombinant adeno-associated viral vector (rAAV) delivery of GDNF provides protection against 6-OHDA lesion in the common marmoset monkey (*Callithrix jacchus*). *Exp Neurol* 184, 536–548.
- Espay AJ, Bonato P, Nahab FB, Maetzler W, et al., 2016. Technology in Parkinson's disease: Challenges and opportunities. *Mov Disord* 31, 1272–1282.
- Espay AJ, Brundin P, and Lang AE, 2017. Precision medicine for disease modification in Parkinson disease. *Nat Rev Neurol*.
- Espay AJ, Morgante F, Merola A, Fasano A, et al., 2018. Levodopa-induced dyskinesia in Parkinson disease: Current and evolving concepts. *Ann Neurol* 84, 797–811.
- Eusebi P, Giannandrea D, Biscetti L, Abraha I, et al., 2017. Diagnostic utility of cerebrospinal fluid  $\alpha$ -synuclein in Parkinson's disease: A systematic review and meta-analysis. *Mov Disord* 32, 1389–1400.
- Fallon JH, and Moore RY, 1978. Catecholamine innervation of the basal forebrain IV. Topography of the dopamine projection to the basal forebrain and neostriatum. *J Comp Neurol* 180, 545–579.
- Feng L, Wang CY, Jiang H, Oho C, et al., 1999. Differential effects of GDNF and BDNF on cultured ventral mesencephalic neurons. *Mol Brain Res* 66, 62–70.
- Finkelstein DI, Stanic D, Parish CL, Tomas D, et al., 2000. Axonal sprouting following lesions of the rat substantia nigra. *Neuroscience* 97, 99–112.
- Fleetwood-Walker SM, Hope PJ, and Mitchell R, 1988. Antinociceptive actions of descending dopaminergic tracts on cat and rat dorsal horn

- somatosensory neurones. *J Physiol* 399, 335–348.
- Forgie A, Doxakis E, Buj-Bello A, Wyatt S, et al., 1999. Differences and Developmental Changes in the Responsiveness of PNS Neurons to GDNF and Neurturin. *Mol Cell Neurosci* 13, 430–440.
- Fougère M, Flaive A, Frigon A, and Ryczko D, 2019. Descending dopaminergic control of brainstem locomotor circuits. *Curr Opin Physiol* 8, 30–35.
- Fox SH, Brochie JM, and Lang AE, 2008. Non-dopaminergic treatments in development for Parkinson's disease. *Lancet Neurol* 7, 927–938.
- Fox SH, Katzenschlager R, Lim SY, Barton B, et al., 2018. International Parkinson and movement disorder society evidence-based medicine review: Update on treatments for the motor symptoms of Parkinson's disease. *Mov Disord*.
- François C, Yelnik J, Tandé D, Agid Y, et al., 1999. Dopaminergic cell group A8 in the monkey: Anatomical organization and projections to the striatum. *J Comp Neurol* 414, 334–347.
- Fundin BT, Mikaelis A, Westphal H, and Ernfors P, 1999. A rapid and dynamic regulation of GDNF-family ligands and receptors correlate with the developmental dependency of cutaneous sensory innervation. *Development* 126, 2597–2610.
- Fuxe K, Agnati LF, Marcoli M, and Borroto-Escuela DO, 2015. Volume Transmission in Central Dopamine and Noradrenaline Neurons and Its Astroglial Targets. *Neurochem Res* 40, 2600–2614.
- Gagne JJ, and Power MC, 2010. Anti-inflammatory drugs and risk of Parkinson disease. *Neurology* 74, 995–1002.
- Galli E, Härkönen T, Sainio MT, Ustav M, et al., 2016. Increased circulating concentrations of mesencephalic astrocyte-derived neurotrophic factor in children with type 1 diabetes. *Sci Rep* 6, 29058.
- Galter D, Pernold K, Yoshitake T, Lindqvist E, et al., 2010. MitoPark mice mirror the slow progression of key symptoms and L-DOPA response in Parkinson's disease. *Genes, Brain Behav* 9, 173–181.
- Gan-Or Z, Dion PA, and Rouleau GA, 2015. Genetic perspective on the role of the autophagy-lysosome pathway in Parkinson disease. *Autophagy* 11, 1443–1457.
- Gao F-J, Wu J-H, Li T-T, Du S-S, et al., 2017a. Identification of Mesencephalic Astrocyte-Derived Neurotrophic Factor as a Novel Neuroprotective Factor for Retinal Ganglion Cells. *Front Mol Neurosci* 10, 76.
- Gao F-J, Zhang S-H, Li T-T, Wu J-H, et al., 2017b. Expression and Distribution of Mesencephalic Astrocyte-Derived Neurotrophic Factor in the Retina and Optic Nerve. *Front Hum Neurosci* 10, 686.
- Gao X, Chen H, Schwarzschild MA, and Ascherio A, 2011. Use of ibuprofen and risk of Parkinson disease. *Neurology* 76, 863–869.
- Garcès A, Haase G, Airaksinen MS, Livet J, et al., 2000. GFR $\alpha$ 1 Is Required for Development of Distinct Subpopulations of Motoneuron. *J Neurosci* 20, 4992 LP – 5000.
- Garcès A, Livet J, Grillet N, Henderson CE, et al., 2001. Responsiveness to neurturin of subpopulations of embryonic rat spinal motoneuron does not correlate with expression of GFR $\alpha$ 1 or GFR $\alpha$ 2. *Dev Dyn* 220, 189–197.
- Garea-Rodríguez E, Eesmaa A, Lindholm P, Schlumbohm C, et al., 2016. Comparative Analysis of the Effects of Neurotrophic Factors CDNF and GDNF in a Nonhuman Primate Model of Parkinson's Disease. *PLoS One* 11, e0149776.
- Gash DM, Zhang Z, Ai Y, Grondin R, et al., 2005. Trophic factor distribution predicts functional recovery in parkinsonian monkeys. *Ann Neurol* 58, 224–233.
- Gash DM, Zhang Z, Cass WA, Ovadia A, et al., 1995. Morphological and functional effects of intranigally administered GDNF in normal rhesus monkeys. *J Comp Neurol* 363, 345–358.
- Gash DM, Zhang Z, Ovadia A, Cass WA, et al., 1996. Functional recovery in parkinsonian monkeys treated with GDNF. *Nature* 380, 252–255.
- Gasmi M, Brandon EP, Herzog CD, Wilson A, et al., 2007a. AAV2-mediated delivery of human neurturin to the rat nigrostriatal system: Long-term efficacy and tolerability of CERE-120 for Parkinson's disease. *Neurobiol Dis* 27, 67–76.
- Gasmi M, Herzog CD, Brandon EP, Cunningham JJ, et al., 2007c. Striatal Delivery of Neurturin by CERE-120, an AAV2 Vector for the Treatment of Dopaminergic Neuron Degeneration in Parkinson's Disease. *Mol Ther* 15, 62–68.
- Gasser PJ, 2019. Roles for the uptake2 transporter OCT3 in regulation of dopaminergic neurotransmission and behavior. *Neurochem Int* 123, 46–49.
- Gauthier J, Parent M, Lévesque M, and Parent A, 1999. The axonal arborization of single nigrostriatal neurons in rats. *Brain Res* 834, 228–232.
- Gavin AM, Walsh S, Wyatt S, O'Keefe GW, et al., 2014. 6-Hydroxydopamine induces distinct alterations in GDF5 and GDNF mRNA expression in the rat nigrostriatal system in vivo. *Neurosci Lett* 561, 176–181.
- Georgievska B, Kirik D, and Björklund A, 2004. Overexpression of glial cell line-derived neurotrophic factor using a lentiviral vector induces time- and dose-dependent downregulation of tyrosine hydroxylase in the intact nigrostriatal dopamine system. *J Neurosci* 24, 6437–6445.
- Georgievska B, Kirik D, and Björklund A, 2002. Aberrant sprouting and downregulation of tyrosine hydroxylase in lesioned nigrostriatal dopamine neurons induced by long-lasting overexpression of glial cell line derived neurotrophic factor in the striatum by lentiviral gene transfer. *Exp Neurol* 177, 461–474.
- Gerdeman GL, Ronesi J, and Lovinger DM, 2002. Postsynaptic endocannabinoid release is critical to long-term depression in the striatum. *Nat Neurosci* 5, 446–451.
- Gerfen CR, and Surmeier DJ, 2011. Modulation of striatal projection systems by dopamine. *Annu Rev Neurosci* 34, 441–466.
- Gerhard A, Pavese N, Hotton G, Turkheimer F, et al., 2006. In vivo imaging of microglial activation with [ $^{11}$ C](R)-PK11195 PET in idiopathic Parkinson's disease. *Neurobiol Dis* 21, 404–412.
- Gerhardt GA, Cass WA, Huettl P, Brock S, et al., 1999. GDNF improves dopamine function in the substantia nigra but not the putamen of unilateral MPTP-lesioned rhesus monkeys. *Brain Res* 817, 163–171.
- Gesi M, Soldani P, Giorgi FS, Santinami A, et al., 2000. The role of the locus coeruleus in the development of Parkinson's disease. *Neurosci Biobehav Rev* 24, 655–668.
- Giguère N, Burke Nanni S, and Trudeau L-E, 2018. On Cell Loss and Selective Vulnerability of Neuronal Populations in Parkinson's Disease. *Front Neurol*.

- Gill SS, Patel NK, Hotton GR, O'Sullivan K, et al., 2003. Direct brain infusion of glial cell line-derived neurotrophic factor in Parkinson disease. *Nat Med* 9, 589–595.
- Glazner GW, Mu X, and Springer JE, 1998. Localization of glial cell line-derived neurotrophic factor receptor alpha and c-ret mRNA in rat central nervous system. *J Comp Neurol* 391, 42–49.
- Glembotski CC, Thuermer DJ, Huang C, Vekich JA, et al., 2012a. Mesencephalic astrocyte-derived neurotrophic factor protects the heart from ischemic damage and is selectively secreted upon sarco/endoplasmic reticulum calcium depletion. *J Biol Chem* 287, 25893–25904.
- Goedert M, Spillantini MG, Del Tredici K, and Braak H, 2013. 100 years of Lewy pathology. *Nat Rev Neurol*.
- Goetz CG, Wu J, McDermott MP, Adler CH, et al., 2008. Placebo response in Parkinson's disease: Comparisons among 11 trials covering medical and surgical interventions. *Mov Disord* 23, 690–699.
- Golden JP, Baloh RH, Kotzbauer PT, Lampe PA, et al., 1998. Expression of neurturin, GDNF, and their receptors in the adult mouse CNS. *J Comp Neurol* 398, 139–150.
- Golden JP, DeMaro JA, Osborne PA, Milbrandt J, et al., 1999. Expression of Neurturin, GDNF, and GDNF Family-Receptor mRNA in the Developing and Mature Mouse. *Exp Neurol* 158, 504–528.
- Golden JP, Milbrandt J, and Johnson EM, 2003. Neurturin and persephin promote the survival of embryonic basal forebrain cholinergic neurons in vitro. *Exp Neurol* 184, 447–455.
- Goldsmith PC, Thind KK, Song T, Kim EJ, et al., 1990. Location of the Neuroendocrine Dopamine Neurons in the Monkey Hypothalamus by Retrograde Tracing and Immunostaining\*, \*\*. *J Neuroendocrinol* 2, 169–179.
- Gomes CAR V, Vaz SH, Ribeiro JA, and Sebastião AM, 2006. Glial cell line-derived neurotrophic factor (GDNF) enhances dopamine release from striatal nerve endings in an adenosine A2A receptor-dependent manner. *Brain Res* 1113, 129–136.
- Gonon FG, and Buda MJ, 1985. Regulation of dopamine release by impulse flow and by autoreceptors as studied by in vivo voltammetry in the rat striatum. *Neuroscience* 14, 765–774.
- Good CH, Hoffman AF, Hoffer BJ, Chefer VI, et al., 2011. Impaired nigrostriatal function precedes behavioral deficits in a genetic mitochondrial model of Parkinson's disease. *FASEB J* 25, 1333–1344.
- Grace AA, and Bunney BS, 1984. The control of firing pattern in nigral dopamine neurons: single spike firing. *J Neurosci* 4, 2877–2890.
- Granhall A-C, Reyland M, Albeck D, Sanders L, et al., 2000. Glial Cell Line-Derived Neurotrophic Factor Is Essential for Postnatal Survival of Midbrain Dopamine Neurons. *J Neurosci* 20, 3182–3190.
- Grattan DR, 2015. The hypothalamo-prolactin axis. *J Endocrinol* 226, T101–T122.
- Graybiel AM, and Ragsdale CW, 1978. Histochemically distinct compartments in the striatum of human, monkeys, and cat demonstrated by acetylthiocholinesterase staining. *Proc Natl Acad Sci* 75, 5723–5726.
- Grealish S, Mattsson B, Draxler P, and Björklund A, 2010. Characterisation of behavioural and neurodegenerative changes induced by intranigral 6-hydroxydopamine lesions in a mouse model of Parkinson's disease. *Eur J Neurosci* 31, 2266–2278.
- Green DR, 2000. Apoptotic pathways: paper wraps stone blunts scissors. *Cell* 102, 1–4.
- Greenamyre JT, and Hastings TG, 2004. Parkinson's -Divergent Causes, Convergent Mechanisms. *Science (80- )* 304, 1120–1122.
- Grimes ML, Zhou J, Beattie EC, Yuen EC, et al., 1996. Endocytosis of Activated TrkA: Evidence that Nerve Growth Factor Induces Formation of Signaling Endosomes. *J Neurosci* 16, 7950–7964.
- Grondin R, Cass WA, Zhang Z, Stanford JA, et al., 2003. Glial cell line-derived neurotrophic factor increases stimulus-evoked dopamine release and motor speed in aged rhesus monkeys. *J Neurosci* 23, 1974–80.
- Grondin R, Littrell OM, Zhang Z, Ai Y, et al., 2019. GDNF revisited: A novel mammalian cell-derived variant form of GDNF increases dopamine turnover and improves brain biodistribution. *Neuropharmacology* 147, 28–36.
- Grondin R, Zhang Z, Ai Y, Ding F, et al., 2008. Intraputamenal infusion of exogenous neurturin protein restores motor and dopaminergic function in the globus pallidus of MPTP-lesioned rhesus monkeys. *Cell Transplant* 17, 373–381.
- Grondin R, Zhang Z, Yi A, Cass WA, et al., 2002. Chronic, controlled GDNF infusion promotes structural and functional recovery in advanced Parkinsonian monkeys. *Brain*.
- Grothe C, and Timmer M, 2007. The physiological and pharmacological role of basic fibroblast growth factor in the dopaminergic nigrostriatal system. *Brain Res Rev* 54, 80–91.
- Gubellini P, and Kachidian P, 2015. Animal models of Parkinson's disease: An updated overview. *Rev Neurol (Paris)*.
- Gulberg HC, and Marsden CA, 1975. Catechol-O-Methyl Transferase: Pharmacological Aspects and Physiological Role. *Pharmacol Rev* 27, 135–206.
- Gulley JM, and Zahniser NR, 2003. Rapid regulation of dopamine transporter function by substrates, blockers and presynaptic receptor ligands. *Eur J Pharmacol* 479, 139–152.
- Guo H, Tang Z, Yu Y, Xu L, et al., 2002. Apomorphine induces trophic factors that support fetal rat mesencephalic dopaminergic neurons in cultures. *Eur J Neurosci* 16, 1861–1870.
- Hadaczek P, Johnston L, Forsayeth J, and Bankiewicz KS, 2010a. Pharmacokinetics and bioactivity of glial cell line-derived factor (GDNF) and neurturin (NTN) infused into the rat brain. *Neuropharmacology* 58, 1114–1121.
- Hallett PJ, Spoelgen R, Hyman BT, Standaert DG, et al., 2006. Dopamine D1 Activation Potentiates Striatal NMDA Receptors by Tyrosine Phosphorylation-Dependent Subunit Trafficking. *J Neurosci* 26, 4690–4700.
- Halliday GM, Blumbergs PC, Cotton RGH, Blessing WW, et al., 1990. Loss of brainstem serotonin- and substance P-containing neurons in Parkinson's disease. *Brain Res* 510, 104–107.
- Halliday GM, and McCann H, 2010. The progression of pathology in Parkinson's disease. *Ann N Y Acad Sci* 1184, 188–195.
- Hamburger V, 1939. Motor and Sensory Hyperplasia following Limb-Bud Transplantations in Chick Embryos. *Physiol Zool* 12, 268–284.
- Hamburger V, 1934. The effects of wing bud extirpation on the development of the central nervous system in chick embryos. *J Exp Zool* 68, 449–494.
- Hamburger V, and Levi-Montalcini R, 1949. Proliferation, differentiation and degeneration in the spinal ganglia of the chick embryo under

- normal and experimental conditions. *J Exp Zool* 111, 457–501.
- Hamilton J F, Morrison PF, Chen MY, Harvey-White J, et al., 2001. Heparin coinfusion during convection-enhanced delivery (CED) increases the distribution of the glial-derived neurotrophic factor (GDNF) ligand family in rat striatum and enhances the pharmacological activity of neurturin. *Exp Neurol* 168, 155–161.
- Hamza TH, Zabetian CP, Tenesa A, Laederach A, et al., 2010. Common genetic variation in the HLA region is associated with late-onset sporadic Parkinson's disease. *Nat Genet* 42, 781–785.
- Hansen C, Sanchez-Ferro A, and Maetzler W, 2018. How Mobile Health Technology and Electronic Health Records Will Change Care of Patients with Parkinson's Disease. *J Parkinsons Dis* 8, S41–S45.
- Hao F, Yang C, Chen SS, Wang YY, et al., 2017. Long-term protective effects of AAV9-mesencephalic astrocyte-derived neurotrophic factor gene transfer in parkinsonian rats. *Exp Neurol* 291, 120–133.
- Hartley CL, Edwards S, Mullan L, Bell PA, et al., 2013. Armet/Manf and Creld2 are components of a specialized ER stress response provoked by inappropriate formation of disulphide bonds: implications for genetic skeletal diseases. *Hum Mol Genet* 22, S262–S275.
- Hassani O-K, François C, Yelnik J, and Féger J, 1997. Evidence for a dopaminergic innervation of the subthalamic nucleus in the rat. *Brain Res* 749, 88–94.
- Hauser RA, Lew MF, Hurtig HI, Ondo WG, et al., 2009. Long-term outcome of early versus delayed rasagiline treatment in early Parkinson's disease. *Mov Disord* 24, 564–573.
- Hawkes CH, Del Tredici K, and Braak H, 2009. Parkinson's Disease. *Ann N Y Acad Sci* 1170, 615–622.
- Hawkes CH, Del Tredici K, and Braak H, 2007. Parkinson's disease: a dual-hit hypothesis. *Neuropathol Appl Neurobiol* 33, 599–614.
- Hayashi H, Ichihara M, Iwashita T, Murakami H, et al., 2000. Characterization of intracellular signals via tyrosine 1062 in RET activated by glial cell line-derived neurotrophic factor. *Oncogene* 19, 4469–4475.
- He Y, Appel S, and Le W, 2001. Minocycline inhibits microglial activation and protects nigral cells after 6-hydroxydopamine injection into mouse striatum. *Brain Res* 909, 187–193.
- Hebert LE, Weuve J, Scherr PA, and Evans DA, 2013. Alzheimer disease in the United States (2010–2050) estimated using the 2010 census. *Neurology* 80, 1778–1783.
- Hebert MA, and Gerhardt GA, 1997. Behavioral and neurochemical effects of intranigral administration of glial cell line-derived neurotrophic factor on aged Fischer 344 rats. *J Pharmacol Exp Ther* 282, 760–8.
- Hebert MA, Van Horne CG, Hoffer BJ, and Gerhardt GA, 1996. Functional effects of GDNF in normal rat striatum: presynaptic studies using in vivo electrochemistry and microdialysis. *J Pharmacol Exp Ther* 279, 1181–90.
- Hedstrom KL, Murtie JC, Albers K, Calcutt NA, et al., 2014. Treating small fiber neuropathy by topical application of a small molecule modulator of ligand-induced GFR $\alpha$ /RET receptor signaling. *Proc Natl Acad Sci* 111, 2325–2330.
- Hegarty S V., Lee DJ, O'Keeffe GW, and Sullivan AM, 2017. Effects of intracerebral neurotrophic factor application on motor symptoms in Parkinson's disease: A systematic review and meta-analysis. *Parkinsonism Relat Disord* 38, 19–25.
- Heikkilä RE, Nicklas WJ, Vyas I, and Duvoisin RC, 1985. Dopaminergic toxicity of rotenone and the 1-methyl-4-phenylpyridinium ion after their stereotaxic administration to rats: Implication for the mechanism of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine toxicity. *Neurosci Lett* 62, 389–394.
- Heiss JD, Lungu C, Hammoud DA, Herscovitch P, et al., 2019. Trial of magnetic resonance-guided putaminal gene therapy for advanced Parkinson's disease. *Mov Disord* mds27724.
- Helkamaa T, Reenilä I, Tuominen RK, Soinila S, et al., 2007. Increased catechol-O-methyltransferase activity and protein expression in OX-42-positive cells in the substantia nigra after lipopolysaccharide microinfusion. *Neurochem Int* 51, 412–423.
- Hellman M, Arumäe U, Yu LY, Lindholm P, et al., 2011. Mesencephalic astrocyte-derived neurotrophic factor (MANF) has a unique mechanism to rescue apoptotic neurons. *J Biol Chem* 286, 2675–2680.
- Hellmich HL, Kos L, Cho ES, Mahon KA, et al., 1996. Embryonic expression of glial cell-line derived neurotrophic factor (GDNF) suggests multiple developmental roles in neural differentiation and epithelial-mesenchymal interactions. *Mech Dev* 54, 95–105.
- Heman-Ackah SM, Manzano R, Hoozemans JJM, Scheper W, et al., 2017. Alpha-synuclein induces the unfolded protein response in Parkinson's disease SNCA triplication iPSC-derived neurons. *Hum Mol Genet* 26, 4441–4450.
- Henderson CE, Phillips HS, Pollock RA, Davies AM, et al., 1994. GDNF: a potent survival factor for motoneurons present in peripheral nerve and muscle. *Science* 266, 1062–4.
- Henderson MJ, Richie CT, Airavaara M, Wang Y, et al., 2013. Mesencephalic astrocyte-derived neurotrophic factor (MANF) secretion and cell surface binding are modulated by KDEL receptors. *J Biol Chem* 288.
- Henderson MX, Cornblath EJ, Darwich A, Zhang B, et al., 2019. Spread of  $\alpha$ -synuclein pathology through the brain connectome is modulated by selective vulnerability and predicted by network analysis. *Nat Neurosci* 22.
- Hendry IA, Stöckel K, Thoenen H, and Iversen LL, 1974. The retrograde axonal transport of nerve growth factor. *Brain Res* 68, 103–121.
- Hernádi I, Grabenhorst F, and Schultz W, 2015. Planning activity for internally generated reward goals in monkey amygdala neurons. *Nat Neurosci* 18, 461–469.
- Hernández-López S, Tkatch T, Perez-Garci E, Galarraga E, et al., 2000. D2 Dopamine Receptors in Striatal Medium Spiny Neurons Reduce L-Type Ca $^{2+}$  Currents and Excitability via a Novel PLC $\beta$ 1–IP3–Calcineurin-Signaling Cascade. *J Neurosci* 20, 8987–8995.
- Hersch SM, Ciliax BJ, Gutekunst C a, Rees HD, et al., 1995. Electron microscopic analysis of D1 and D2 dopamine receptor proteins in the dorsal striatum and their synaptic relationships with motor corticostriatal afferents. *J Neurosci* 15, 5222–5237.
- Hersch SM, Gutekunst CA, Rees HD, Heilman CJ, et al., 1994. Distribution of m1-m4 muscarinic receptor proteins in the rat striatum: light and electron microscopic immunocytochemistry using subtype-specific antibodies. *J Neurosci* 14, 3351–3363.
- Hersch SM, Yi H, Heilman CJ, Edwards RH, et al., 1997. Subcellular localization and molecular topology of the dopamine transporter in the striatum and substantia nigra. *J Comp Neurol* 388, 211–227.
- Herzog CD, Dass B, Gasmi M, Bakay R, et al., 2008. Transgene Expression, Bioactivity, and Safety of CERE-120 (AAV2-Neurturin) Following Delivery to the Monkey Striatum. *Mol Ther* 16, 1737–1744.

- Herzog CD, Dass B, Holden JE, Stansell III J, et al., 2007. Striatal delivery of CERE-120, an AAV2 vector encoding human neurturin, enhances activity of the dopaminergic nigrostriatal system in aged monkeys. *Mov Disord* 22, 1124–1132.
- Hetz C, and Saxena S, 2017. ER stress and the unfolded protein response in neurodegeneration. *Nat Rev Neurol* 13, 477–491.
- Heuckeroth RO, Enomoto H, Grider JR, Golden JP, et al., 1999. Gene Targeting Reveals a Critical Role for Neurturin in the Development and Maintenance of Enteric, Sensory, and Parasympathetic Neurons. *Neuron* 22, 253–263.
- Heuckeroth RO, Lampe PA, Johnson EM, and Milbrandt J, 1998. Neurturin and GDNF Promote Proliferation and Survival of Enteric Neuron and Glial Progenitors in Vitro. *Dev Biol* 200, 116–129.
- Hidalgo-Figueroa M, Bonilla S, Gutierrez F, Pascual a., et al., 2012. GDNF Is Predominantly Expressed in the PV+ Neostriatal Interneuron Ensemble in Normal Mouse and after Injury of the Nigrostriatal Pathway. *J Neurosci* 32, 864–872.
- Hiltunen JO, Laurikainen A, Airaksinen MS, and Saarma M, 2000. GDNF family receptors in the embryonic and postnatal rat heart and reduced cholinergic innervation in mice hearts lacking Ret or GFR $\alpha$ 2. *Dev Dyn* 219, 28–39.
- Hiroi T, Imaoka S, and Funae Y, 1998. Dopamine Formation from Tyramine by CYP2D6. *Biochem Biophys Res Commun* 249, 838–843.
- Hirsch EC, and Hunot S, 2009. Neuroinflammation in Parkinson's disease: a target for neuroprotection? *Lancet Neurol* 8, 382–397.
- Hirsch EC, Orieux G, Muriel MP, Francois C, et al., 2003. Nondopaminergic neurons in Parkinson's disease. *Adv Neurol* 91, 29–37.
- Hnasko TS, Chuhma N, Zhang H, Goh GY, et al., 2010. Vesicular Glutamate Transport Promotes Dopamine Storage and Glutamate Corelease In Vivo. *Neuron* 65, 643–656.
- Hoffer BJ, Hoffman A, Bowenkamp K, Huettl P, et al., 1994. Glial cell line-derived neurotrophic factor reverses toxin-induced injury to midbrain dopaminergic neurons in vivo. *Neurosci Lett* 182, 107–111.
- Hoffman AF, van Horne CG, Eken S, Hoffer BJ, et al., 1997. In Vivo Microdialysis Studies of Somatodendritic Dopamine Release in the Rat Substantia Nigra: Effects of Unilateral 6-OHDA Lesions and GDNF. *Exp Neurol* 147, 130–141.
- Honma Y, Araki T, Gianino S, Bruce A, et al., 2002. Artemin Is a Vascular-Derived Neurotrophic Factor for Developing Sympathetic Neurons. *Neuron* 35, 267–282.
- Hooks BM, Papale AE, Paletzki RF, Feroze MW, et al., 2018. Topographic precision in sensory and motor corticostriatal projections varies across cell type and cortical area. *Nat Commun* 9, 3549.
- Hoozemans JJM, Haastert ES [van, Eikelenboom P, Vos] RAI [de, et al., 2007. Activation of the unfolded protein response in Parkinson's disease. *Biochem Biophys Res Commun* 354, 707–711.
- Horger BA, Nishimura MC, Armanini MP, Wang LC, et al., 1998. Neurturin exerts potent actions on survival and function of midbrain dopaminergic neurons. *J Neurosci* 18, 4929–37.
- Hornykiewicz O, 2002. Dopamine miracle: From brain homogenate to dopamine replacement. *Mov Disord* 17, 501–508.
- Hornykiewicz O, 1978. Psychopharmacological implications of dopamine and dopamine antagonists: a critical evaluation of current evidence. *Neuroscience* 3, 773–783.
- Hoseki J, Sasakawa H, Yamaguchi Y, Maeda M, et al., 2010. Solution structure and dynamics of mouse ARMET. *FEBS Lett* 584, 1536–1542.
- Hou J-GG, Lin L-FH, and Mytilineou C, 1996. Glial Cell Line-Derived Neurotrophic Factor Exerts Neurotrophic Effects on Dopaminergic Neurons In Vitro and Promotes Their Survival and Regrowth After Damage by 1-Methyl-4-Phenylpyridinium. *J Neurochem* 66, 74–82.
- Howe CL, Valletta JS, Rusnak AS, and Mobley WC, 2001. NGF signaling from clathrin-coated vesicles: Evidence that signaling endosomes serve as a platform for the Ras-MAPK pathway. *Neuron* 32, 801–814.
- Howe MW, and Dombeck DA, 2016. Rapid signalling in distinct dopaminergic axons during locomotion and reward. *Nature* 535, 505–510.
- Howell MJ, and Schenck CH, 2015. Rapid Eye Movement Sleep Behavior Disorder and Neurodegenerative Disease. *JAMA Neurol* 72, 707–712.
- Howells DW, Porritt MJ, Wong JYF, Batchelor PE, et al., 2000. Reduced BDNF mRNA Expression in the Parkinson's Disease Substantia Nigra. *Exp Neurol* 166, 127–135.
- Hsiao EC, Koniaris LG, Zimmers-Koniaris T, Sebald SM, et al., 2000. Characterization of Growth-Differentiation Factor 15, a Transforming Growth Factor  $\beta$  Superfamily Member Induced following Liver Injury. *Mol Cell Biol* 20, 3742–3751.
- Hsu J-Y, Crawley S, Chen M, Ayupova DA, et al., 2017. Non-homeostatic body weight regulation through a brainstem-restricted receptor for GDF15. *Nature* 550, 255–259.
- Huang EJ, and Reichardt LF, 2001. Neurotrophins: Roles in Neuronal Development and Function. *Annu Rev Neurosci* 24, 677–736.
- Hudson J, Granholm A-C, Gerhardt GA, Henry MA, et al., 1995. Glial cell line-derived neurotrophic factor augments midbrain dopaminergic circuits in vivo. *Brain Res Bull* 36, 425–432.
- Hunot S, Dugas N, Faucheux B, Hartmann A, et al., 1999. Fc $\epsilon$ R2/CD23 Is Expressed in Parkinson's Disease and Induces, In Vitro, Production of Nitric Oxide and Tumor Necrosis Factor- $\alpha$  in Glial Cells. *J Neurosci* 19, 3440–3447.
- Huotari A, Penttinen A-M, Bäck S, Voutilainen MH, et al., 2018. Combination of CDNF and Deep Brain Stimulation Decreases Neurological Deficits in Late-stage Model Parkinson's Disease. *Neuroscience* 374, 250–263.
- Huttunen HJ, and Saarma M, 2019. CDNF Protein Therapy in Parkinson's Disease. *Cell Transplant*.
- Hyland BI, Reynolds JNJ, Hay J, Perk CG, et al., 2002. Firing modes of midbrain dopamine cells in the freely moving rat. *Neuroscience* 114, 475–492.
- Ibáñez CF, 1998. Emerging themes in structural biology of neurotrophic factors. *Trends Neurosci* 21, 438–444.
- Ibáñez CF, and Andressoo J-O, 2017. Biology of GDNF and its receptors — Relevance for disorders of the central nervous system. *Neurobiol Dis* 97, 80–89.
- Iida M, Miyazaki I, Tanaka K, Kabuto H, et al., 1999. Dopamine D2 receptor-mediated antioxidant and neuroprotective effects of ropinirole, a dopamine agonist. *Brain Res* 838, 51–59.
- Ilieva M, Nielsen J, Korshunova I, Gotfryd K, et al., 2019. Artemin and an artemin-derived peptide, artefin, induce neuronal survival, and differentiation through ret and NCAM. *Front Mol Neurosci* 12.
- Iravani MM, Costa S, Jackson MJ, Tel BC, et al., 2001. GDNF reverses priming for dyskinesia in MPTP-treated, L-DOPA-primed common marmosets. *Eur J Neurosci* 13, 597–608.

- Irwin DJ, Lee VM-Y, and Trojanowski JQ, 2013. Parkinson's disease dementia: convergence of  $\alpha$ -synuclein, tau and amyloid- $\beta$  pathologies. *Nat Rev Neurosci* 14, 626–636.
- Ishida H, Saba R, Kokkinopoulos I, Hashimoto M, et al., 2016. GFRA2 Identifies Cardiac Progenitors and Mediates Cardiomyocyte Differentiation in a RET-Independent Signaling Pathway. *Cell Rep* 16, 1026–1038.
- Ito K, and Enomoto H, 2016. Retrograde transport of neurotrophic factor signaling: implications in neuronal development and pathogenesis. *J Biochem*.
- Ivanova L, Tammiku-Taul J, García-Sosa AT, Sidorova Y, et al., 2018. Molecular Dynamics Simulations of the Interactions between Glial Cell Line-Derived Neurotrophic Factor Family Receptor GFR $\alpha$ 1 and Small-Molecule Ligands. *ACS Omega* 3, 11407–11414.
- Iwase T, Jung CG, Bae H, Zhang M, et al., 2005. Glial cell line-derived neurotrophic factor-induced signaling in Schwann cells. *J Neurochem* 94, 1488–1499.
- Jackson-Lewis V, and Przedborski S, 2007. Protocol for the MPTP mouse model of Parkinson's disease. *Nat Protoc* 2, 141–151.
- Jain S, Naughton CK, Yang M, Strickland A, et al., 2004. Mice expressing a dominant-negative Ret mutation phenocopy human Hirschsprung disease and delineate a direct role of Ret in spermatogenesis. *Development* 131, 5503–5513.
- Janezic S, Threlfell S, Dodson PD, Dowie MJ, et al., 2013. Deficits in dopaminergic transmission precede neuron loss and dysfunction in a new Parkinson model. *Proc Natl Acad Sci U S A* 110, E4016–E4025.
- Jankovic J, Chen S, and Le WD, 2005. The role of Nurr1 in the development of dopaminergic neurons and Parkinson's disease. *Prog Neurobiol* 77, 128–138.
- Jaszai J, Farkas L, Galter D, Reuss B, et al., 1998. GDNF-related factor persephin is widely distributed throughout the nervous system. *J Neurosci Res* 53, 494–501.
- Jellinger KA, 2012. Interaction between pathogenic proteins in neurodegenerative disorders. *J Cell Mol Med* 16, 1166–1183.
- Jellinger KA, 2009. A critical evaluation of current staging of  $\alpha$ -synuclein pathology in Lewy body disorders. *Biochim Biophys Acta - Mol Basis Dis* 1792, 730–740.
- Jia Y, Gall CM, and Lynch G, 2010. Presynaptic BDNF Promotes Postsynaptic Long-Term Potentiation in the Dorsal Striatum. *J Neurosci* 30, 14440–14445.
- Jijiwa M, Kawai K, Fukihara J, Nakamura A, et al., 2008. GDNF-mediated signaling via RET tyrosine 1062 is essential for maintenance of spermatogonial stem cells. *Genes to Cells* 13, 365–374.
- Jin X, and Costa RM, 2010. Start/stop signals emerge in nigrostriatal circuits during sequence learning. *Nature* 466, 457–462.
- Jing S, Wen D, Yu Y, Holst PL, et al., 1996. GDNF-induced activation of the ret protein tyrosine kinase is mediated by GDNFR- $\alpha$ , a novel receptor for GDNF. *Cell* 85, 1113–24.
- Jing S, Yu Y, Fang M, Hu Z, et al., 1997. GFR $\alpha$ -2 and GFR $\alpha$ -3 are two new receptors for ligands of the GDNF family. *J Biol Chem* 272, 33111–33117.
- Jomary C, Thomas M, Grist J, Milbrandt J, et al., 1999. Expression patterns of neurturin and its receptor components in developing and degenerative mouse retina. *Invest Ophthalmol Vis Sci* 40, 568–574.
- Jones IW, Bolam JP, and Wonnacott S, 2001. Presynaptic localisation of the nicotinic acetylcholine receptor  $\beta$ 2 subunit immunoreactivity in rat nigrostriatal dopaminergic neurones. *J Comp Neurol* 439, 235–247.
- Joyce JN, Woolsey C, Ryoo H, Borwege S, et al., 2004. Low dose pramipexole is neuroprotective in the MPTP mouse model of Parkinson's disease, and downregulates the dopamine transporter via the D3 receptor. *BMC Biol* 2, 22.
- Julku UH, Panhelainen AE, Tiilikainen SE, Svarcbahts R, et al., 2016. Prolyl Oligopeptidase Regulates Dopamine Transporter Phosphorylation in the Nigrostriatal Pathway of Mouse. *Mol Neurobiol* 1–13.
- Käenmäki M, Tammimäki A, Myöhänen T, Pakarinen K, et al., 2010. Quantitative role of COMT in dopamine clearance in the prefrontal cortex of freely moving mice. *J Neurochem* 114, 1745–1755.
- Kalaitzakis ME, Graeber MB, Gentleman SM, and Pearce RKB, 2008. The dorsal motor nucleus of the vagus is not an obligatory trigger site of Parkinson's disease: a critical analysis of  $\alpha$ -synuclein staging. *Neuropathol Appl Neurobiol* 34, 284–295.
- Kalia LV, 2018. Biomarkers for cognitive dysfunction in Parkinson's disease. *Parkinsonism Relat Disord* 46, S19–S23.
- Kalia LV, and Lang AE, 2015. Parkinson's disease. *Lancet* 386, 896–912.
- Kalia LV, Kalia SK, McLean PJ, Lozano AM, et al., 2013.  $\alpha$ -Synuclein oligomers and clinical implications for Parkinson disease. *Ann Neurol* 73, 155–169.
- Kalia LV, Lang AE, Hazrati L-N, Fujioka S, et al., 2015. Clinical Correlations With Lewy Body Pathology in LRRK2-Related Parkinson Disease. *JAMA Neurol* 72, 100–105.
- Kalinderi K, Bostantjopoulou S, and Fidani L, 2016. The genetic background of Parkinson's disease: current progress and future prospects. *Acta Neurol Scand*.
- Karampetsou M, Ardah MT, Semitekolou M, Polissidis A, et al., 2017. Phosphorylated exogenous  $\alpha$ -synuclein fibrils exacerbate pathology and induce neuronal dysfunction in mice. *Sci Rep* 7, 1–18.
- Karpinar DP, Balija MBG, Kügler S, Opazo F, et al., 2009. Pre-fibrillar  $\alpha$ -synuclein variants with impaired  $\beta$ -structure increase neurotoxicity in Parkinson's disease models. *EMBO J* 28, 3256–3268.
- Karpowicz RJ, Trojanowski JQ, and Lee VMY, 2019. Transmission of  $\alpha$ -synuclein seeds in neurodegenerative disease: recent developments. *Lab Invest*.
- Kawaguchi Y, Wilson CJ, Augood SJ, and Emson PC, 1995. Striatal interneurons: chemical, physiological and morphological characterization. *Trends Neurosci* 18, 527–535.
- Kelley AE, Domesick VB, and Nauta WJH, 1982. The amygdalostratial projection in the rat—an anatomical study by anterograde and retrograde tracing methods. *Neuroscience* 7, 615–630.
- Kerr JND, and Wickens JR, 2001. Dopamine D-1/D-5 Receptor Activation Is Required for Long-Term Potentiation in the Rat Neostriatum In Vitro. *J Neurophysiol* 85, 117–124.
- Kettenmann H, Kirchhoff F, and Verkhratsky A, 2013. Microglia: New Roles for the Synaptic Stripper. *Neuron* 77, 10–18.

- Killinger BA, and Kordower JH, 2019. Spreading of alpha-synuclein – relevant or epiphenomenon? *J Neurochem* 150, 605–611.
- Kim Y, Park S-J, and Chen YM, 2017. Mesencephalic astrocyte-derived neurotrophic factor (MANF), a new player in endoplasmic reticulum diseases: structure, biology, and therapeutic roles. *Transl Res* 188, 1–9.
- Kim YS, and Joh TH, 2006. Microglia, major player in the brain inflammation: their roles in the pathogenesis of Parkinson's disease. *Exp Mol Med* 38, 333–347.
- Kirik D, Annett LE, Burger C, Muzyczka N, et al., 2003. Nigrostriatal  $\alpha$ -synucleinopathy induced by viral vector-mediated overexpression of human  $\alpha$ -synuclein: A new primate model of Parkinson's disease. *Proc Natl Acad Sci* 100, 2884–2889.
- Kirik D, Georgievska B, and Björklund A, 2004. Localized striatal delivery of GDNF as a treatment for Parkinson disease. *Nat Neurosci*.
- Kirik D, Rosenblad C, and Björklund A, 2000. Preservation of a functional nigrostriatal dopamine pathway by GDNF in the intrastriatal 6-OHDA lesion model depends on the site of administration of the trophic factor. *Eur J Neurosci* 12, 3871–3882.
- Kirik D, Rosenblad C, and Björklund A, 1998a. Characterization of Behavioral and Neurodegenerative Changes Following Partial Lesions of the Nigrostriatal Dopamine System Induced by Intrastriatal 6-Hydroxydopamine in the Rat. *Exp Neurol* 152, 259–277.
- Kirik D, Rosenblad C, Björklund A, and Mandel RJ, 2000. Long-term rAAV-mediated gene transfer of GDNF in the rat Parkinson's model: intrastriatal but not intranigral transduction promotes functional regeneration in the lesioned nigrostriatal system. *J Neurosci* 20, 4686–700.
- Kirik D, Rosenblad C, Burger C, Lundberg C, et al., 2002. Parkinson-Like Neurodegeneration Induced by Targeted Overexpression of  $\alpha$ -Synuclein in the Nigrostriatal System. *J Neurosci* 22, 2780–2791.
- Kish SJ, Tong J, Hornykiewicz O, Rajput A, et al., 2008. Preferential loss of serotonin markers in caudate versus putamen in Parkinson's disease. *Brain* 131, 120–131.
- Kitada T, Asakawa S, Hattori N, Matsumine H, et al., 1998. Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism. *Nature* 392, 605–608.
- Kitada T, Tong Y, Gautier CA, and Shen J, 2009. Absence of nigral degeneration in aged parkin/DJ-1/PINK1 triple knockout mice. *J Neurochem* 111, 696–702.
- Klassen BT, Hentz JG, Shill HA, Driver-Dunckley E, et al., 2011. Quantitative EEG as a predictive biomarker for Parkinson disease dementia. *Neurology* 77, 118–124.
- Klaus A, da Silva J, and Costa RM, 2019. What, If, and When to Move: Basal Ganglia Circuits and Self-Paced Action Initiation. *Annu Rev Neurosci* 42, 459–483.
- Klaus A, Martins GJ, Paixao VB, Zhou P, et al., 2017. The Spatiotemporal Organization of the Striatum Encodes Action Space. *Neuron* 95, 1171–1180.e7.
- Klegeris A, Pelech S, Giasson BI, Maguire J, et al., 2008.  $\alpha$ -Synuclein activates stress signaling protein kinases in THP-1 cells and microglia. *Neurobiol Aging* 29, 739–752.
- Klein RD, Sherman D, Ho W-H, Stone D, et al., 1997. A GPI-linked protein that interacts with Ret to form a candidate neurturin receptor. *Nature* 387, 717–721.
- Knowles PP, Murray-Rust J, Kjaer S, Scott RP, et al., 2006. Structure and chemical inhibition of the RET tyrosine kinase domain. *J Biol Chem* 281, 33577–33587.
- Kobayashi S, Ögren SO, Hoffer BJ, and Olson L, 1998. Dopamine D1 and D2 Receptor-Mediated Acute and Long-Lasting Behavioral Effects of Glial Cell Line-Derived Neurotrophic Factor Administered into the Striatum. *Exp Neurol* 154, 302–314.
- Kobori N, Waymire JC, Haycock JW, Clifton GL, et al., 2004. Enhancement of tyrosine hydroxylase phosphorylation and activity by glial cell line-derived neurotrophic factor. *J Biol Chem* 279, 2182–91.
- Kopra J, Vilenius C, Grealish S, Härma M-A, et al., 2015. GDNF is not required for catecholaminergic neuron survival in vivo. *Nat Neurosci* 18, 319–322.
- Kopra J, Panhelainen A, af Björkén S, Porokukka LL, et al., 2017. Dampened Amphetamine-Stimulated Behavior and Altered Dopamine Transporter Function in the Absence of Brain GDNF. *J Neurosci* 37, 1581–1590.
- Kordower JH, Chu Y, Hauser RA, Freeman TB, et al., 2008. Lewy body-like pathology in long-term embryonic nigral transplants in Parkinson's disease. *Nat Med* 14, 504–506.
- Kordower JH, Emborg ME, Bloch J, Ma SY, et al., 2000. Neurodegeneration prevented by lentiviral vector delivery of GDNF in primate models of Parkinson's disease. *Science* 290, 767–73.
- Kordower JH, Herzog CD, Dass B, Bakay RAE, et al., 2006. Delivery of neurturin by AAV2 (CERE-120)-mediated gene transfer provides structural and functional neuroprotection and neurorestoration in MPTP-treated monkeys. *Ann Neurol* 60, 706–715.
- Kordower JH, Olanow CW, Dodiya HB, Chu Y, et al., 2013a. Disease duration and the integrity of the nigrostriatal system in Parkinson's disease. *Brain* 136, 2419–2431.
- Kordower JH, Palfi S, Chen E-Y, Ma SY, et al., 1999. Clinicopathological findings following intraventricular glial-derived neurotrophic factor treatment in a patient with Parkinson's disease. *Ann Neurol* 46, 419–424.
- Kotzbauer PT, Lampe PA, Heuckeroth RO, Golden JP, et al., 1996. Neurturin, a relative of glial-cell-line-derived neurotrophic factor. *Nature* 384, 467–470.
- Kourtis LC, Regele OB, Wright JM, and Jones GB, 2019. Digital biomarkers for Alzheimer's disease: the mobile/wearable devices opportunity. *npj Digit Med* 2, 9.
- Koyama Y, Tsujikawa K, Matsuda T, and Baba A, 2003a. Endothelin-1 stimulates glial cell line-derived neurotrophic factor expression in cultured rat astrocytes. *Biochem Biophys Res Commun* 303, 1101–1105.
- Koyama Y, Tsujikawa K, Matsuda T, and Baba A, 2003b. Intracerebroventricular administration of an endothelin ETB receptor agonist increases expressions of GDNF and BDNF in rat brain. *Eur J Neurosci* 18, 887–894.
- Kozlowski DA, Miljan E, Bremer E, Harrod C, et al., 2004. Quantitative analyses of GFR $\alpha$ -1 and GFR $\alpha$ -2 mRNAs and tyrosine hydroxylase protein in the nigrostriatal system reveal bilateral compensatory changes following unilateral 6-OHDA lesions in the rat. *Brain Res* 1016, 170–181.
- Kozlowski DA, Connor B, Tillerson JL, Schallert T, et al., 2000. Delivery of a GDNF Gene into the Substantia Nigra after a Progressive 6-OHDA



- Lesion Maintains Functional Nigrostriatal Connections. *Exp Neurol* 166, 1–15.
- Kramer BC, Goldman AD, and Mytilineou C, 1999. Glial cell line derived neurotrophic factor promotes the recovery of dopamine neurons damaged by 6-hydroxydopamine in vitro. *Brain Res* 851, 221–227.
- Kramer ER, Aron L, Ramakers GMJ, Seitz S, et al., 2007. Absence of Ret signaling in mice causes progressive and late degeneration of the nigrostriatal system. *PLoS Biol* 5, 0616–0628.
- Kramer ER, Knott L, Su F, Dessaud E, et al., 2006. Cooperation between GDNF/Ret and ephrinA/EphA4 Signals for Motor-Axon Pathway Selection in the Limb. *Neuron* 50, 35–47.
- Kramer ER, and Liss B, 2015. GDNF–Ret signaling in midbrain dopaminergic neurons and its implication for Parkinson disease. *FEBS Lett* 589, 3760–3772.
- Kreitzer AC, 2009. Physiology and Pharmacology of Striatal Neurons. *Annu Rev Neurosci* 32, 127–147.
- Kumar A, Kopra J, Varendi K, Porokuokka LL, et al., 2015. GDNF Overexpression from the Native Locus Reveals its Role in the Nigrostriatal Dopaminergic System Function. *PLOS Genet* 11, e1005710.
- Kumer SC, and Vrana KE, 1996. Intricate Regulation of Tyrosine Hydroxylase Activity and Gene Expression. *J Neurochem* 67, 443–462.
- Kuno R, Yoshida Y, Nitta A, Nabeshima T, et al., 2006. The role of TNF-alpha and its receptors in the production of NGF and GDNF by astrocytes. *Brain Res* 1116, 12–18.
- Kupchik YM, Brown RM, Heinsbroek JA, Lobo MK, et al., 2015. Coding the direct/indirect pathways by D1 and D2 receptors is not valid for accumbens projections. *Nat Neurosci* 18, 1230–1232.
- Lang AE, Gill S, Patel NK, Lozano A, et al., 2006. Randomized controlled trial of intraputamenal glial cell line-derived neurotrophic factor infusion in Parkinson disease. *Ann Neurol* 59, 459–466.
- Langston JW, Ballard P, Tetrud JW, and Irwin I, 1983. Chronic Parkinsonism in humans due to a product of meperidine-analog synthesis. *Science* 219, 979–980.
- Langston JW, Forno LS, Tetrud J, Reeves AG, et al., 1999. Evidence of active nerve cell degeneration in the substantia nigra of humans years after 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine exposure. *Ann Neurol* 46, 598–605.
- Lanza G, Bachmann CG, Ghorayeb I, Wang Y, et al., 2017. Central and peripheral nervous system excitability in restless legs syndrome. *Sleep Med* 31, 49–60.
- Lapchak PA, Araujo DM, Hilt DC, Jiao S, et al., 1998. Topographical distribution of [125I]-glial cell line-derived neurotrophic factor in unlesioned and MPTP-lesioned rhesus monkey brain following a bolus intraventricular injection. *Brain Res* 789, 9–22.
- Lapchak PA, Jiao S, Collins F, and Miller PJ, 1997. Glial cell line-derived neurotrophic factor: distribution and pharmacology in the rat following a bolus intraventricular injection. *Brain Res* 747, 92–102.
- Larsen KE, Schmitz Y, Troyer MD, Mosharov E, et al., 2006.  $\alpha$ -Synuclein Overexpression in PC12 and Chromaffin Cells Impairs Catecholamine Release by Interfering with a Late Step in Exocytosis. *J Neurosci* 26, 11915–11922.
- Latgé C, Cabral KMS, Almeida MS, and Foguel D, 2013. <sup>1</sup>H-, <sup>13</sup>C- and <sup>15</sup>N-NMR assignment of the N-terminal domain of human cerebral dopamine neurotrophic factor (CDNF). *Biomol NMR Assign* 7, 101–103.
- Latge C, Cabral KMS, de Oliveira GAP, Raymundo DP, et al., 2015a. The Solution Structure and Dynamics of Full-length Human Cerebral Dopamine Neurotrophic Factor and Its Neuroprotective Role against  $\alpha$ -Synuclein Oligomers. *J Biol Chem* 290, 20527–20540.
- Latreille V, Carrier J, Gaudet-Fex B, Rodrigues-Brazete J, et al., 2016. Electroencephalographic prodromal markers of dementia across conscious states in Parkinson's disease. *Brain* 139, 1189–1199.
- Laurikainen A, Hiltunen JO, Thomas-Crusells J, Vanhatalo S, et al., 2000. Neurturin is a neurotrophic factor for penile parasympathetic neurons in adult rat. *J Neurobiol* 43, 198–205.
- Lavoie B, Smith Y, and Parent A, 1989. Dopaminergic innervation of the basal ganglia in the squirrel monkey as revealed by tyrosine hydroxylase immunohistochemistry. *J Comp Neurol* 289, 36–52.
- Ledda F, Paratcha G, and Ibáñez CF, 2002. Target-Derived GFR $\alpha$ 1 as an Attractive Guidance Signal for Developing Sensory and Sympathetic Axons via Activation of Cdk5. *Neuron* 36, 387–401.
- Ledda F, Paratcha G, Sandoval-Guzmán T, and Ibáñez CF, 2007. GDNF and GFR $\alpha$ 1 promote formation of neuronal synapses by ligand-induced cell adhesion. *Nat Neurosci* 10, 293–300.
- Lee A-H, Iwakoshi NN, and Glimcher LH, 2003. XBP-1 regulates a subset of endoplasmic reticulum resident chaperone genes in the unfolded protein response. *Mol Cell Biol* 23, 7448–7459.
- Lee CS, Samii A, Sossi V, Ruth TJ, et al., 2000. In vivo positron emission tomographic evidence for compensatory changes in presynaptic dopaminergic nerve terminals in Parkinson's disease. *Ann Neurol* 47, 493–503.
- Lee E, Hwang I, Park S, Hong S, et al., 2019. MPTP-driven NLRP3 inflammasome activation in microglia plays a central role in dopaminergic neurodegeneration. *Cell Death Differ* 26, 213–228.
- Lee JM, Derkinderen P, Kordower JH, Freeman R, et al., 2017. The Search for a Peripheral Biopsy Indicator of  $\alpha$ -Synuclein Pathology for Parkinson Disease. *J Neuropathol Exp Neurol* 76, 2–15.
- Lehtonen Š, Sonninen TM, Wojciechowski S, Goldsteins G, et al., 2019. Dysfunction of cellular proteostasis in Parkinson's disease. *Front Neurosci*.
- Leitner ML, Molliver DC, Osborne PA, Vejsada R, et al., 1999. Analysis of the Retrograde Transport of Glial Cell Line-Derived Neurotrophic Factor (GDNF), Neurturin, and Persephin Suggests That In Vivo Signaling for the GDNF Family is GFR $\alpha$  Coreceptor-Specific. *J Neurosci* 19, 9322–9331.
- Leppänen V-M, Bessalov MM, Runeberg-Roos P, Puurand Ü, et al., 2004. The structure of GFR $\alpha$ 1 domain 3 reveals new insights into GDNF binding and RET activation. *EMBO J* 23, 1452–1462.
- Lerner TN, Shilyansky C, Davidson TJ, Evans KE, et al., 2015. Intact-Brain Analyses Reveal Distinct Information Carried by SNc Dopamine Subcircuits. *Cell* 162, 635–647.
- Leroy E, Boyer R, Auburger G, Leube B, et al., 1998. The ubiquitin pathway in Parkinson's disease. *Nature* 395, 451–452.
- Levy MJF, Bouille F, Steinbusch HW, van den Hove DLA, et al., 2018. Neurotrophic factors and neuroplasticity pathways in the



- pathophysiology and treatment of depression. *Psychopharmacology (Berl)* 235, 2195–2220.
- LeWitt PA, Li J, Lu M, Beach TG, et al., 2013. 3-hydroxykynurenine and other Parkinson's disease biomarkers discovered by metabolomic analysis. *Mov Disord* 28, 1653–1660.
- Li J-Y, Englund E, Holton JL, Soulet D, et al., 2008. Lewy bodies in grafted neurons in subjects with Parkinson's disease suggest host-to-graft disease propagation. *Nat Med* 14, 501–503.
- Li L, Wu W, Lin LF, Lei M, et al., 1995. Rescue of adult mouse motoneurons from injury-induced cell death by glial cell line-derived neurotrophic factor. *Proc Natl Acad Sci U S A* 92, 9771–9775.
- Liberatore GT, Wong JYF, Porritt MJ, Donnan GA, et al., 1997. Expression of glial cell line-derived neurotrophic factor (GDNF) mRNA following mechanical injury to mouse striatum. *Neuroreport* 8, 3097–3101.
- Lin L-FH, Doherty DH, Lile JD, Bektesh S, et al., 1993. GDNF: a glial cell line-derived neurotrophic factor for midbrain dopaminergic neurons. *Science* 260, 1130–1132.
- Lin L-FH, Zhang TJ, Collins F, and Armes LG, 1994. Purification and Initial Characterization of Rat B49 Glial Cell Line-Derived Neurotrophic Factor. *J Neurochem* 63, 758–768.
- Lindahl M, Chalazonitis A, Palm E, Pakarinen E, et al., 2020. Cerebral dopamine neurotrophic factor–deficiency leads to degeneration of enteric neurons and altered brain dopamine neuronal function in mice. *Neurobiol Dis* 134.
- Lindahl M, Danilova T, Palm E, Lindholm P, et al., 2014a. MANF Is Indispensable for the Proliferation and Survival of Pancreatic  $\beta$  Cells. *Cell Rep* 7, 366–375.
- Lindahl M, Saarma M, and Lindholm P, 2017. Unconventional neurotrophic factors CDNF and MANF: Structure, physiological functions and therapeutic potential. *Neurobiol Dis* 97, 90–102.
- Lindahl M, Timmusk T, Rossi J, Saarma M, et al., 2000. Expression and Alternative Splicing of Mouse Gfra4 Suggest Roles in Endocrine Cell Development. *Mol Cell Neurosci* 15, 522–533.
- Lindfors PH, Lindahl M, Rossi J, Saarma M, et al., 2006. Ablation of Persephin Receptor Glial Cell Line-Derived Neurotrophic Factor Family Receptor  $\alpha 4$  Impairs Thyroid Calcitonin Production in Young Mice. *Endocrinology* 147, 2237–2244.
- Lindholm P, Peränen J, Andressoo J-O, Kalkkinen N, et al., 2008. MANF is widely expressed in mammalian tissues and differently regulated after ischemic and epileptic insults in rodent brain. *Mol Cell Neurosci* 39, 356–371.
- Lindholm P, and Saarma M, 2010. Novel CDNF/MANF family of neurotrophic factors. *Dev Neurobiol* 70, 360–371.
- Lindholm P, Voutilainen MH, Laurén J, Peränen J, et al., 2007. Novel neurotrophic factor CDNF protects and rescues midbrain dopamine neurons in vivo. *Nature* 448, 73–77.
- Lindvall O, and Björklund A, 1979. Dopaminergic innervation of the globus pallidus by collaterals from the nigrostriatal pathway. *Brain Res* 172, 169–173.
- Littrell OM, Pomerleau F, Huettl P, Surgener S, et al., 2012. Enhanced dopamine transporter activity in middle-aged Gdnf heterozygous mice. *Neurobiol Aging* 33, 427.e1–427.e14.
- Liu H, Yu C, Yu H, Zhong L, et al., 2018. Cerebral dopamine neurotrophic factor protects H9c2 cardiomyocytes from apoptosis. *Herz* 43, 346–351.
- Liu J, Zhou C, Tao X, Feng L, et al., 2015. ER stress-inducible protein MANF selectively expresses in human spleen. *Hum Immunol* 76, 823–830.
- Liu X, and Herbison AE, 2013. Dopamine Regulation of Gonadotropin-Releasing Hormone Neuron Excitability in Male and Female Mice. *Endocrinology* 154, 340–350.
- Lo Bianco C, Déglon N, Pralong W, and Aebischer P, 2004a. Lentiviral nigral delivery of GDNF does not prevent neurodegeneration in a genetic rat model of Parkinson's disease. *Neurobiol Dis* 17, 283–289.
- Lo Bianco C, Ridet J-L, Schneider BL, Déglon N, et al., 2002.  $\alpha$ -Synucleinopathy and selective dopaminergic neuron loss in a rat lentiviral-based model of Parkinson's disease. *Proc Natl Acad Sci* 99, 10813–10818.
- Loeffler DA, Camp DM, and Conant SB, 2006. Complement activation in the Parkinson's disease substantia nigra: an immunocytochemical study. *J Neuroinflammation* 3, 29.
- Lohse MJ, 1993. Molecular mechanisms of membrane receptor desensitization. *Biochim Biophys Acta - Mol Cell Res* 1179, 171–188.
- Lonka-Nevalaita L, Lume M, Leppänen S, Jokitalo E, et al., 2010. Characterization of the intracellular localization, processing, and secretion of two glial cell line-derived neurotrophic factor splice isoforms. *J Neurosci* 30, 11403–11413.
- Loughlin SE, and Fallon JH, 1984. Substantia nigra and ventral tegmental area projections to cortex: Topography and collateralization. *Neuroscience* 11, 425–435.
- Love S, Plaha P, Patel NK, Hutton GR, et al., 2005. Glial cell line–derived neurotrophic factor induces neuronal sprouting in human brain. *Nat Med* 11, 703–704.
- Lu B, Nagappan G, and Lu Y, 2014. BDNF and Synaptic Plasticity, Cognitive Function, and Dysfunction. Springer, Berlin, Heidelberg, pp. 223–250.
- Lu J, Luo L, Huang D, Liu X, et al., 2018. Photoreceptor Protection by Mesencephalic Astrocyte-Derived Neurotrophic Factor (MANF). *eNeuro* 5.
- Lu X-H, Fleming SM, Meurers B, Ackerson LC, et al., 2009. Bacterial Artificial Chromosome Transgenic Mice Expressing a Truncated Mutant Parkin Exhibit Age-Dependent Hypokinetic Motor Deficits, Dopaminergic Neuron Degeneration, and Accumulation of Proteinase K-Resistant  $\alpha$ -Synuclein. *J Neurosci* 29, 1962–1976.
- Luk KC, Kehm VM, Carroll J, Zhang B, et al., 2012a. Pathological  $\alpha$ -Synuclein Transmission Initiates Parkinson-like Neurodegeneration in Nontransgenic Mice. *Science* 338, 949–953.
- Luk KC, Kehm VM, Zhang B, O'Brien P, et al., 2012b. Intracerebral inoculation of pathological  $\alpha$ -synuclein initiates a rapidly progressive neurodegenerative  $\alpha$ -synucleinopathy in mice. *J Exp Med* 209, 975–986.
- Luth ES, Stavrovskaya IG, Bartels T, Kristal BS, et al., 2014. Soluble, prefibrillar  $\alpha$ -synuclein oligomers promote complex I-dependent,  $\text{Ca}^{2+}$ -induced mitochondrial dysfunction. *J Biol Chem* 289, 21490–21507.

- Lwin A, Orvisky E, Goker-Alpan O, LaMarca ME, et al., 2004. Glucocerebrosidase mutations in subjects with parkinsonism. *Mol Genet Metab* 81, 70–73.
- Mabe AM, Hoard JL, Duffourc MM, and Hoover DB, 2006. Localization of cholinergic innervation and neurturin receptors in adult mouse heart and expression of the neurturin gene. *Cell Tissue Res* 326, 57–67.
- Macpherson T, and Hikida T, 2019. Role of basal ganglia neurocircuitry in the pathology of psychiatric disorders. *Psychiatry Clin Neurosci* 73, 289–301.
- Maeda K, Kunimatsu J, and Hikosaka O, 2018. Amygdala activity for the modulation of goal-directed behavior in emotional contexts. *PLoS Biol* 16, e2005339.
- Mahadevan N, Demanuele C, Zhang H, Volfson D, et al., 2020. Development of digital biomarkers for resting tremor and bradykinesia using a wrist-worn wearable device. *npj Digit Med* 3, 5.
- Mahato AK, Kopra J, Renko J-M, Visnapuu T, et al., 2020. Glial Cell Line-Derived Neurotrophic Factor Receptor Rearranged During Transfection Agonist Supports Dopamine Neurons In Vitro and Enhances Dopamine Release In Vivo. *Mov Disord* 35, 245–255.
- Mahul-Mellier A-L, Bartscher J, Maharjan N, Weerens L, et al., 2020. The process of Lewy body formation, rather than simply  $\alpha$ -synuclein fibrillization, is one of the major drivers of neurodegeneration. *Proc Natl Acad Sci* 117, 4971–4982.
- Mak MK, Wong-Yu IS, Shen X, and Chung CL, 2017. Long-term effects of exercise and physical therapy in people with Parkinson disease. *Nat Rev Neurol* 13, 689–703.
- Mallet N, Ballion B, Le Moine C, and Gonon F, 2006. Cortical inputs and GABA interneurons imbalance projection neurons in the striatum of parkinsonian rats. *J Neurosci* 26, 3875–3884.
- Mallet N, Le Moine C, Charpier S, and Gonon F, 2005. Feedforward Inhibition of Projection Neurons by Fast-Spiking GABA Interneurons in the Rat Striatum In Vivo. *J Neurosci* 25, 3857–3869.
- Mallet N, Micklem BR, Henny P, Brown MT, et al., 2012. Dichotomous Organization of the External Globus Pallidus. *Neuron* 74, 1075–1086.
- Mallet N, Schmidt R, Leventhal D, Chen F, et al., 2016. Arky pallidal Cells Send a Stop Signal to Striatum. *Neuron* 89, 308–316.
- Mamaligas AA, Cai Y, and Ford CP, 2016. Nicotinic and opioid receptor regulation of striatal dopamine D2-receptor mediated transmission. *Sci Rep* 6, 37834.
- Manning-Bog AB, McCormack AL, Li J, Uversky VN, et al., 2002. The Herbicide Paraquat Causes Up-regulation and Aggregation of  $\alpha$ -Synuclein in Mice: Paraquat and  $\alpha$ -Synuclein. *J Biol Chem* 277, 1641–1644.
- Marco S, Saura J, Pérez-Navarro E, José Martí M, et al., 2002. Regulation of c-Ret, GFR $\alpha$ 1, and GFR $\alpha$ 2 in the substantia nigra Pars compacta in a rat model of Parkinson's disease. *J Neurobiol* 52, 343–351.
- Marinelli M, and McCutcheon JE, 2014. Heterogeneity of dopamine neuron activity across traits and states. *Neuroscience* 282, 176–197.
- Markowitz JE, Gillis WF, Beron CC, Neufeld SQ, et al., 2018. The Striatum Organizes 3D Behavior via Moment-to-Moment Action Selection. *Cell* 174, 44–58.e17.
- Marks WJ, Bartus RT, Siffert J, Davis CS, et al., 2010. Gene delivery of AAV2-neurturin for Parkinson's disease: a double-blind, randomised, controlled trial. *Lancet Neurol* 9, 1164–1172.
- Marks WJ, Ostrem JL, Verhagen L, Starr PA, et al., 2008. Safety and tolerability of intraputamenal delivery of CERE-120 (adeno-associated virus serotype 2-neurturin) to patients with idiopathic Parkinson's disease: an open-label, phase I trial. *Lancet Neurol* 7, 400–408.
- Marras C, Beck JC, Bower JH, Roberts E, et al., 2018. Prevalence of Parkinson's disease across North America. *npj Park Dis* 4, 21.
- Marras C, and Lang A, 2013. Parkinson's disease subtypes: lost in translation? *J Neurol Neurosurg Psychiatry* 84, 409–415.
- Marsden CD, and Parkes JD, 1976. "On-Off" Effects in Patients with Parkinson's Disease on Chronic Levodopa Therapy. *Lancet* 307, 292–296.
- Martin D, Miller G, Cullen T, Fischer N, et al., 1996. Intranigral or intrastriatal injections of GDNF: Effects on monoamine levels and behavior in rats. *Eur J Pharmacol* 317, 247–256.
- Martin I, Kim JW, Dawson VL, and Dawson TM, 2014. LRRK2 pathobiology in Parkinson's disease. *J Neurochem* 131, 554–565.
- Martinez-Martin P, Reddy P, Katzenschlager R, Antonini A, et al., 2015. EuroInf: A Multicenter Comparative Observational Study of Apomorphine and Levodopa Infusion in Parkinson's Disease. *Mov Disord* 30, 510–516.
- Maruyama W, Nitta A, Shamoto-Nagai M, Hirata Y, et al., 2004. N-Propargyl-1 (R)-aminoindan, rasagiline, increases glial cell line-derived neurotrophic factor (GDNF) in neuroblastoma SH-SY5Y cells through activation of NF- $\kappa$ B transcription factor. *Neurochem Int* 44, 393–400.
- Masure S, Cik M, Hoefnagel E, Nosrat CA, et al., 2000. Mammalian GFR $\alpha$ 4, a divergent member of the GFR $\alpha$  family of coreceptors for glial cell line-derived neurotrophic factor family ligands, is a receptor for the neurotrophic factor persephin. *J Biol Chem* 275, 39427–39434.
- Masure S, Cik M, Pangalos MN, Bonaventure P, et al., 1998. Molecular cloning, expression and tissue distribution of glial-cell-line-derived neurotrophic factor family receptor  $\alpha$ -3 (GFR $\alpha$ -3). *Eur J Biochem* 251, 622–630.
- Matsumoto M, and Hikosaka O, 2009. Two types of dopamine neuron distinctly convey positive and negative motivational signals. *Nature* 459, 837–841.
- Maurice N, Mercer J, Chan CS, Hernandez-Lopez S, et al., 2004. D2 Dopamine Receptor-Mediated Modulation of Voltage-Dependent Na<sup>+</sup> Channels Reduces Autonomous Activity in Striatal Cholinergic Interneurons. *J Neurosci* 24, 10289–10301.
- McCormack AL, Thiruchelvam M, Manning-Bog AB, Thiffault C, et al., 2002. Environmental Risk Factors and Parkinson's Disease: Selective Degeneration of Nigral Dopaminergic Neurons Caused by the Herbicide Paraquat. *Neurobiol Dis* 10, 119–127.
- McCoy MK, and Cookson MR, 2012. Mitochondrial Quality Control and Dynamics in Parkinson's Disease. *Antioxid Redox Signal* 16, 869–882.
- McGeer PL, Itagaki S, Boyes BE, and McGeer EG, 1988. Reactive microglia are positive for HLA-DR in the substantia nigra of Parkinson's and Alzheimer's disease brains. *Neurology* 38, 1285.
- McGeorge AJ, and Faull RLM, 1989. The organization of the projection from the cerebral cortex to the striatum in the rat. *Neuroscience* 29, 503–537.
- McGregor MM, and Nelson AB, 2019. Circuit Mechanisms of Parkinson's Disease. *Neuron* 101, 1042–1056.

- McIlvried LA, Albers K, and Gold MS, 2010. Distribution of Artemin and GFR $\alpha$ 3 Labeled Nerve Fibers in the Dura Mater of Rat. *Headache J Head Face Pain* 50, 442–450.
- McNaught K, Belizaire R, Isacson O, Jenner P, et al., 2003. Altered Proteasomal Function in Sporadic Parkinson's Disease. *Exp Neurol* 179, 38–46.
- McNaught K, Belizaire R, Jenner P, Olanow CW, et al., 2002. Selective loss of 20S proteasome  $\alpha$ -subunits in the substantia nigra pars compacta in Parkinson's disease. *Neurosci Lett* 326, 155–158.
- McNaught KS, and Jenner P, 2001. Proteasomal function is impaired in substantia nigra in Parkinson's disease. *Neurosci Lett* 297, 191–194.
- McNaught KS, and Jenner P, 2000. Dysfunction of rat forebrain astrocytes in culture alters cytokine and neurotrophic factor release. *Neurosci Lett* 285, 61–65.
- McNaught KSP, Björklund LM, Belizaire R, Isacson O, et al., 2002. Proteasome inhibition causes nigral degeneration with inclusion bodies in rats. *Neuroreport* 13, 1437–1441.
- Mei J, and Niu C, 2014. Effects of CDNF on 6-OHDA-induced apoptosis in PC12 cells via modulation of Bcl-2/Bax and caspase-3 activation. *Neurol Sci* 35, 1275–1280.
- Meiser J, Weindl D, and Hiller K, 2013. Complexity of dopamine metabolism. *Cell Commun Signal* 11, 34.
- Melillo RM, Santoro M, Ong S-H, Billaud M, et al., 2001. Docking protein FRS2 links the protein tyrosine kinase RET and its oncogenic forms with the mitogen-activated protein kinase signaling cascade. *Mol Cell Biol* 21, 4177–4187.
- Meng X, Lindahl M, Hyvönen ME, Parvinen M, et al., 2000. Regulation of cell fate decision of undifferentiated spermatogonia by GDNF. *Science* 287, 1489–1493.
- Mercado G, Castillo V, Soto P, López N, et al., 2018. Targeting PERK signaling with the small molecule GSK2606414 prevents neurodegeneration in a model of Parkinson's disease. *Neurobiol Dis* 112, 136–148.
- Mercado G, Valdés P, and Hetz C, 2013. An ERcentric view of Parkinson's disease. *Trends Mol Med* 19, 165–175.
- Mijatovic J, Airavaara M, Planken A, Auvinen P, et al., 2007. Constitutive Ret activity in knock-in multiple endocrine neoplasia type B mice induces profound elevation of brain dopamine concentration via enhanced synthesis and increases the number of TH-positive cells in the substantia nigra. *J Neurosci* 27, 4799–4809.
- Mijatovic J, Patrikainen O, Yavich L, Airavaara M, et al., 2008. Characterization of the striatal dopaminergic neurotransmission in MEN2B mice with elevated cerebral tissue dopamine. *J Neurochem* 105, 1716–1725.
- Mikaels Å, Livet J, Westphal H, De Lapeyrière O, et al., 2000. A dynamic regulation of GDNF-family receptors correlates with a specific trophic dependency of cranial motor neuron subpopulations during development. *Eur J Neurosci* 12, 446–456.
- Milbrandt J, de Sauvage FJ, Fahrner TJ, Baloh RH, et al., 1998a. Persephin, a Novel Neurotrophic Factor Related to GDNF and Neurturin. *Neuron* 20, 245–253.
- Miller DB, and O'Callaghan JP, 2015. Biomarkers of Parkinson's disease: Present and future. *Metabolism* 64.
- Mizobuchi N, Hoseki J, Kubota H, Toyokuni S, et al., 2007. ARMET is a soluble ER protein induced by the unfolded protein response via ERSE-II element. *Cell Struct Funct* 32, 41–50.
- Mogi M, Togari A, Kondo T, Mizuno Y, et al., 2001. Glial cell line-derived neurotrophic factor in the substantia nigra from control and parkinsonian brains. *Neurosci Lett* 300, 179–181.
- Mogi M, Togari A, Kondo T, Mizuno Y, et al., 1999. Brain-derived growth factor and nerve growth factor concentrations are decreased in the substantia nigra in Parkinson's disease. *Neurosci Lett* 270, 45–48.
- Moon HE, and Paek SH, 2015. Mitochondrial Dysfunction in Parkinson's Disease. *Exp Neurobiol* 24, 103–116.
- Moore MW, Klein RD, Fariñas I, Sauer H, et al., 1996a. Renal and neuronal abnormalities in mice lacking GDNF. *Nature* 382, 76–79.
- Morrison PF, Lonser RR, and Oldfield EH, 2007. Convective delivery of glial cell line-derived neurotrophic factor in the human putamen. *J Neurosurg* 107, 74–83.
- Mount HT, Dean DO, Alberch J, Dreyfus CF, et al., 1995. Glial cell line-derived neurotrophic factor promotes the survival and morphologic differentiation of Purkinje cells. *Proc Natl Acad Sci U S A* 92, 9092–6.
- Mufson EJ, Kroin JS, Sendera TJ, and Sobriela T, 1999. Distribution and retrograde transport of trophic factors in the central nervous system: functional implications for the treatment of neurodegenerative diseases. *Prog Neurobiol* 57, 451–484.
- Müller CP, Pum ME, Amato D, Schüttler J, et al., 2011. The in vivo neurochemistry of the brain during general anesthesia. *J Neurochem* 119, 419–446.
- Mullican SE, Lin-Schmidt X, Chin C-N, Chavez JA, et al., 2017. GFRAL is the receptor for GDF15 and the ligand promotes weight loss in mice and nonhuman primates. *Nat Med* 23, 1150–1157.
- Murphy DD, Rueter SM, Trojanowski JQ, and Lee VM-Y, 2000. Synucleins Are Developmentally Expressed, and  $\alpha$ -Synuclein Regulates the Size of the Presynaptic Vesicular Pool in Primary Hippocampal Neurons. *J Neurosci* 20, 3214–3220.
- Myers SM, Eng C, Ponder BAJ, and Mulligan LM, 1995. Characterization of RET proto-oncogene 3' splicing variants and polyadenylation sites: A novel C-terminus for RET. *Oncogene* 11, 2039–2045.
- Myöhänen TT, Schendzielorz N, and Männistö PT, 2010. Distribution of catechol-O-methyltransferase (COMT) proteins and enzymatic activities in wild-type and soluble COMT deficient mice. *J Neurochem* 113, 1632–1643.
- Männistö PT, and Kaakkola S, 1999. Catechol-O-methyltransferase (COMT): Biochemistry, Molecular Biology, Pharmacology, and Clinical Efficacy of the New Selective COMT Inhibitors. *Pharmacol Rev* 51, 593–628.
- Mätlik K, Vihinen H, Bienemann A, Palgi J, et al., 2017. Intrastrially Infused Exogenous CDNF Is Endocytosed and Retrogradely Transported to Substantia Nigra. *eNeuro* 4.
- Mätlik K, Yu L, Eesmaa A, Hellman M, et al., 2015. Role of two sequence motifs of mesencephalic astrocyte-derived neurotrophic factor in its survival-promoting activity. *Cell Death Dis* 6, e2032.
- Nadella R, Voutilainen MH, Saarma M, Gonzalez-Barrios JA, et al., 2014. Transient transfection of human CDNF gene reduces the 6-hydroxydopamine-induced neuroinflammation in the rat substantia nigra. *J Neuroinflammation* 11.
- Nakagawa T, and Schwartz JP, 2004. Gene Expression Profiles of Reactive Astrocytes in Dopamine-depleted Striatum. *Brain Pathol* 14, 275–

- 280.
- Nakagawa T, Yabe T, and Schwartz JP, 2005. Gene expression profiles of reactive astrocytes cultured from dopamine-depleted striatum. *Neurobiol Dis* 20, 275–282.
- Nalls MA, Pankratz N, Lill CM, Do CB, et al., 2014. Large-scale meta-analysis of genome-wide association data identifies six new risk loci for Parkinson's disease. *Nat Genet* 46, 989–993.
- Nambu A, Tokuno H, and Takada M, 2002. Functional significance of the cortico–subthalamo–pallidal 'hyperdirect' pathway. *Neurosci Res* 43, 111–117.
- Naughton CK, Jain S, Strickland AM, Gupta A, et al., 2006. Glial Cell-Line Derived Neurotrophic Factor-Mediated RET Signaling Regulates Spermatogonial Stem Cell Fate. *Biol Reprod* 74, 314–321.
- Naveilhan P, Baudet C, Mikaelis A, Shen L, et al., 1998. Expression and regulation of GFRalpha3, a glial cell line-derived neurotrophic factor family receptor. *Proc Natl Acad Sci U S A* 95, 1295–1300.
- Nemani VM, Lu W, Berge V, Nakamura K, et al., 2010. Increased Expression of  $\alpha$ -Synuclein Reduces Neurotransmitter Release by Inhibiting Synaptic Vesicle Reclustering after Endocytosis. *Neuron* 65, 66–79.
- Neve KA, Seamans JK, and Trantham-Davidson H, 2004. Dopamine Receptor Signaling. *J Recept Signal Transduct* 24, 165–205.
- Neves J, Zhu J, Sousa-Victor P, Konjikusic M, et al., 2016. Immune modulation by MANF promotes tissue repair and regenerative success in the retina. *Science* (80- ) 353.
- Nicklas WJ, Vyas I, and Heikkilä RE, 1985. Inhibition of NADH-linked oxidation in brain mitochondria by 1-methyl-4-phenyl-pyridine, a metabolite of the neurotoxin, 1-methyl-4-phenyl-1,2,5,6-tetrahydropyridine. *Life Sci* 36, 2503–2508.
- Nielsen J, Gotfryd K, Li S, Kulahin N, et al., 2009. Role of Glial Cell Line-Derived Neurotrophic Factor (GDNF)–Neural Cell Adhesion Molecule (NCAM) Interactions in Induction of Neurite Outgrowth and Identification of a Binding Site for NCAM in the Heel Region of GDNF. *J Neurosci* 29, 11360–11376.
- Nirenberg MJ, Vaughan R a, Uhl GR, Kuhar MJ, et al., 1996. The dopamine transporter is localized to dendritic and axonal plasma membranes of nigrostriatal dopaminergic neurons. *J Neurosci* 16, 436–447.
- Nishikawa K, Li H, Kawamura R, Osaka H, et al., 2003. Alterations of structure and hydrolase activity of parkinsonism-associated human ubiquitin carboxyl-terminal hydrolase L1 variants. *Biochem Biophys Res Commun* 304, 176–183.
- Nishino J, Mochida K, Ohfuji Y, Shimazaki T, et al., 1999. GFR $\alpha$ 3, a Component of the Artemin Receptor, Is Required for Migration and Survival of the Superior Cervical Ganglion. *Neuron* 23, 725–736.
- Nivlet L, Herrmann J, Martin DE, Meunier A, et al., 2016. Expression and functional studies of the GDNF family receptor alpha 3 in the pancreas. *J Mol Endocrinol* 56, 77–90.
- Nomoto S, Ito S, Yang L-X, and Kiuchi K, 1998. Molecular Cloning and Expression Analysis of GFR $\alpha$ -3, a Novel cDNA Related to GDNFR $\alpha$  and NTN $\alpha$ . *Biochem Biophys Res Commun* 244, 849–853.
- Norisada J, Hirata Y, Amaya F, Kiuchi K, et al., 2016. A comparative analysis of the molecular features of MANF and CDNF. *PLoS One* 11.
- Nosrat CA, Tomac A, Hoffer BJ, and Olson L, 1997. Cellular and developmental patterns of expression of Ret and glial cell line-derived neurotrophic factor receptor alpha mRNAs. *Exp Brain Res* 115, 410–422.
- Nosrat CA, Tomac A, Lindqvist E, Lindskog S, et al., 1996. Cellular expression of GDNF mRNA suggests multiple functions inside and outside the nervous system. *Cell Tissue Res* 286, 191–207.
- Noyce AJ, Bestwick JP, Silveira-Moriyama L, Hawkes CH, et al., 2012. Meta-analysis of early nonmotor features and risk factors for Parkinson disease. *Ann Neurol* 72, 893–901.
- Nutt JG, Burchiel KJ, Comella CL, Jankovic J, et al., 2003. Randomized, double-blind trial of glial cell line-derived neurotrophic factor (GDNF) in PD. *Neurology* 60, 69–73.
- O'Callaghan C, and Lewis SJG, 2017. Cognition in Parkinson's Disease, in: International Review of Neurobiology.
- O'Keefe GW, and Sullivan AM, 2018. Evidence for dopaminergic axonal degeneration as an early pathological process in Parkinson's disease. *Parkinsonism Relat Disord* 56, 9–15.
- Oh-Hashi K, Hirata Y, and Kiuchi K, 2013. Transcriptional regulation of mouse mesencephalic astrocyte-derived neurotrophic factor in Neuro2a cells. *Cell Mol Biol Lett* 18, 398–415.
- Oh-hashish K, Tanaka K, Koga H, Hirata Y, et al., 2012a. Intracellular trafficking and secretion of mouse mesencephalic astrocyte-derived neurotrophic factor. *Mol Cell Biochem* 363, 35–41.
- Ohta K, Fujinami A, Kuno S, Sakakimoto A, et al., 2004. Cabergoline Stimulates Synthesis and Secretion of Nerve Growth Factor, Brain-Derived Neurotrophic Factor and Glial Cell Line-Derived Neurotrophic Factor by Mouse Astrocytes in Primary Culture. *Pharmacology* 71, 162–168.
- Ohta M, Mizuta I, Ohta K, Nishimura M, et al., 2000. Apomorphine Up-Regulates NGF and GDNF Synthesis in Cultured Mouse Astrocytes. *Biochem Biophys Res Commun* 272, 18–22.
- Oiwa Y, Yoshimura R, Nakai K, and Itakura T, 2002. Dopaminergic neuroprotection and regeneration by neurturin assessed by using behavioral, biochemical and histochemical measurements in a model of progressive Parkinson's disease. *Brain Res* 947, 271–283.
- Okuzumi A, Kurosawa M, Hatano T, Takanashi M, et al., 2018. Rapid dissemination of alpha-synuclein seeds through neural circuits in an in-vivo prion-like seeding experiment. *Acta Neuropathol Commun* 6, 96.
- Olanow CW, Bartus RT, Volpicelli-daley LA, and Kordower JH, 2015. Trophic Factors for Parkinson's Disease : To Live or Let Die. *Mov Disord* 30, 1715–1723.
- Olanow CW, and McNaught KSP, 2006. Ubiquitin–proteasome system and Parkinson's disease. *Mov Disord* 21, 1806–1823.
- Olanow CW, Rascol O, Hauser R, Feigin PD, et al., 2009. A Double-Blind, Delayed-Start Trial of Rasagiline in Parkinson's Disease. *N Engl J Med* 361, 1268–1278.
- Omiatsek DM, Bressler AJ, Cans A-S, Andrews AM, et al., 2013. The real catecholamine content of secretory vesicles in the CNS revealed by electrochemical cytometry. *Sci Rep* 3, 1447.
- Omodaka K, Kurimoto T, Nakamura O, Sato K, et al., 2014. Artemin augments survival and axon regeneration in axotomized retinal ganglion

- cells. *J Neurosci Res* 92, 1637–1646.
- Oo TF, Ries V, Cho J, Kholodilov N, et al., 2005. Anatomical basis of glial cell line-derived neurotrophic factor expression in the striatum and related basal ganglia during postnatal development of the rat. *J Comp Neurol* 484, 57–67.
- Oppenheim RW, 1991. Cell Death During Development of the Nervous System. *Annu Rev Neurosci* 14, 453–501.
- Oppenheim RW, Houenou LJ, Johnson JE, Lin LF, et al., 1995. Developing motor neurons rescued from programmed and axotomy-induced cell death by GDNF. *Nature* 373, 344–6.
- Orozco OE, Walus L, Sah DWY, Pepinsky RB, et al., 2001. GFR $\alpha$ 3 is expressed predominantly in nociceptive sensory neurons. *Eur J Neurosci* 13, 2177–2182.
- Ortega-de San Luis C, and Pascual A, 2016. Simultaneous Detection of Both GDNF and GFR $\alpha$ 1 Expression Patterns in the Mouse Central Nervous System. *Front Neuroanat* 10.
- Ossowska K, Wardas J, Śmiałowska M, Kuter K, et al., 2005. A slowly developing dysfunction of dopaminergic nigrostriatal neurons induced by long-term paraquat administration in rats: an animal model of preclinical stages of Parkinson's disease? *Eur J Neurosci* 22, 1294–1304.
- Ouchi Y, Yoshikawa E, Sekine Y, Futatsubashi M, et al., 2005. Microglial activation and dopamine terminal loss in early Parkinson's disease. *Ann Neurol* 57, 168–175.
- Ovallath S, and Deepa P, 2013. The history of parkinsonism: Descriptions in ancient Indian medical literature. *Mov Disord* 28, 566–568.
- Pachnis V, Mankoo B, and Costantini F, 1993. Expression of the c-ret proto-oncogene during mouse embryogenesis. *Development* 119.
- Paisán-Ruiz C, Jain S, Evans EW, Gilks WP, et al., 2004. Cloning of the Gene Containing Mutations that Cause PARK8-Linked Parkinson's Disease. *Neuron* 44, 595–600.
- Palgi M, Lindström R, Peränen J, Piepponen TP, et al., 2009. Evidence that DmMANF is an invertebrate neurotrophic factor supporting dopaminergic neurons. *Proc Natl Acad Sci U S A* 106, 2429–2434.
- Palmatier MA, Hartman BK, and Johnson EM, 1984. Demonstration of retrogradely transported endogenous nerve growth factor in axons of sympathetic neurons. *J Neurosci* 4, 751 LP – 756.
- Pan T, Kondo S, Le W, and Jankovic J, 2008. The role of autophagy-lysosome pathway in neurodegeneration associated with Parkinson's disease. *Brain* 131, 1969–1978.
- Pan W, Mao T, and Dudman J, 2010. Inputs to the Dorsal Striatum of the Mouse Reflect the Parallel Circuit Architecture of the Forebrain. *Front Neuroanat* 4, 147.
- Pang SYY, Ho PWL, Liu HF, Leung CT, et al., 2019. The interplay of aging, genetics and environmental factors in the pathogenesis of Parkinson's disease. *Transl Neurodegener*.
- Parain K, Murer M, Yan Q, Faucheux B, et al., 1999. Reduced expression of brain-derived neurotrophic factor protein in Parkinson's disease substantia nigra. *Neuroreport* 10, 557–561.
- Paratcha G, and Ibáñez CF, 2002. Lipid rafts and the control of neurotrophic factor signaling in the nervous system: variations on a theme. *Curr Opin Neurobiol* 12, 542–549.
- Paratcha G, Ibáñez CF, and Ledda F, 2006. GDNF is a chemoattractant factor for neuronal precursor cells in the rostral migratory stream. *Mol Cell Neurosci* 31, 505–514.
- Paratcha G, and Ledda F, 2008a. GDNF and GFR $\alpha$ : a versatile molecular complex for developing neurons. *Trends Neurosci* 31, 384–391.
- Paratcha G, Ledda F, Baars L, Couplier M, et al., 2001. Released GFR $\alpha$ 1 Potentiates Downstream Signaling, Neuronal Survival, and Differentiation via a Novel Mechanism of Recruitment of c-Ret to Lipid Rafts. *Neuron* 29, 171–184.
- Paratcha G, Ledda F, and Ibáñez CF, 2003a. The neural cell adhesion molecule NCAM is an alternative signaling receptor for GDNF family ligands. *Cell* 113, 867–879.
- Parkash V, Lindholm P, Peränen J, Kalkkinen N, et al., 2009. The structure of the conserved neurotrophic factors MANF and CDNF explains why they are bifunctional. *Protein Eng Des Sel* 22, 233–241.
- Parker JG, Marshall JD, Ahanonu B, Wu Y-W, et al., 2018. Diametric neural ensemble dynamics in parkinsonian and dyskinetic states. *Nature* 557, 177–182.
- Parker NF, Cameron CM, Taliaferro JP, Lee J, et al., 2016. Reward and choice encoding in terminals of midbrain dopamine neurons depends on striatal target. *Nat Neurosci* 19, 845–854.
- Parkinson J, 2002. An essay on the shaking palsy. *J Neuropsychiatry Clin Neurosci* 14, 223–236.
- Parkinson Study Group, 2004. A Controlled, Randomized, Delayed-Start Study of Rasagiline in Early Parkinson Disease. *Arch Neurol* 61, 561–566.
- Parkinson Study Group, 2002. Dopamine Transporter Brain Imaging to Assess the Effects of Pramipexole vs Levodopa on Parkinson Disease Progression. *JAMA* 287, 1653–1661.
- Parkkinen L, Pirttilä T, and Alafuzoff I, 2008. Applicability of current staging/categorization of  $\alpha$ -synuclein pathology and their clinical relevance. *Acta Neuropathol* 115, 399–407.
- Parnetti L, Gaetani L, Eusebi P, Paciotti S, et al., 2019. CSF and blood biomarkers for Parkinson's disease. *Lancet Neurol* 18, 573–586.
- Pascual A, Hidalgo-Figueroa M, Piruat JJ, Pintado CO, et al., 2008. Absolute requirement of GDNF for adult catecholaminergic neuron survival. *Nat Neurosci* 11, 755–761.
- Pascual A, and López-Barneo J, 2015. Reply to “GDNF is not required for catecholaminergic neuron survival in vivo.” *Nat Neurosci* 18, 322–323.
- Patel NK, Bunnage M, Plaha P, Svendsen CN, et al., 2005. Intrapataminal infusion of glial cell line-derived neurotrophic factor in PD: A two-year outcome study. *Ann Neurol* 57, 298–302.
- Patel NK, Pavese N, Hotton GR, et al., 2013. Benefits of putaminal GDNF infusion in Parkinson disease are maintained after GDNF cessation. *Neurology* 81, 1176–1178.
- Paul G, and Sullivan AM, 2018. Trophic factors for Parkinson's disease: Where are we and where do we go from here? *Eur J Neurosci*.
- Paul G, Zachrisson O, Varrone A, Almqvist P, et al., 2015. Safety and tolerability of intracerebroventricular PDGF-BB in Parkinson's disease

- patients. *J Clin Invest* 125, 1339–1346.
- Paxinos G, and Watson C, 1998. The rat brain in stereotaxic coordinates, 4th editio. ed. Academic Press, Inc., San Diego.
- Penttinen A-M, Suleymanova I, Albert K, Anttila J, et al., 2016. Characterization of a new low-dose 6-hydroxydopamine model of Parkinson's disease in rat. *J Neurosci Res* 94, 318–328.
- Penttinen A-M, Parkkinen I, Blom S, Kopra J, et al., 2018. Implementation of deep neural networks to count dopamine neurons in substantia nigra. *Eur J Neurosci* 48.
- Perese DA, Ulman J, Viola J, Ewing SE, et al., 1989. A 6-hydroxydopamine-induced selective parkinsonian rat model. *Brain Res* 494, 285–293.
- Perez-Pardo P, Kliest T, Dodiya HB, Broersen LM, et al., 2017. The gut-brain axis in Parkinson's disease: Possibilities for food-based therapies. *Eur J Pharmacol* 817, 86–95.
- Petrova PS, Raibekas A, Pevsner J, Vigo N, et al., 2003. MANF: A new mesencephalic, astrocyte-derived neurotrophic factor with selectivity for dopaminergic neurons. *J Mol Neurosci* 20, 173–187.
- Pezzoli G, and Cereda E, 2013. Exposure to pesticides or solvents and risk of Parkinson disease. *Neurology* 80, 2035–2041.
- Piccinini E, Kalkkinen N, Saarma M, and Runeberg-Roos P, 2013. Glial cell line-derived neurotrophic factor: Characterization of mammalian posttranslational modifications. *Ann Med* 45, 66–73.
- Pichel JG, Shen L, Sheng HZ, Granholm AC, et al., 1996. Defects in enteric innervation and kidney development in mice lacking GDNF. *Nature* 382, 73–6.
- Pickel VM, Nirenberg MJ, and Milner T a, 1996. Ultrastructural view of central catecholaminergic transmission: immunocytochemical localization of synthesizing enzymes, transporters and receptors. *J Neurocytol* 25, 843–856.
- Pickrell AM, and Youle RJ, 2015. The roles of PINK1, Parkin, and mitochondrial fidelity in Parkinson's disease. *Neuron*.
- Piepponen TP, and Skujins A, 2001. Rapid and sensitive step gradient assays of glutamate, glycine, taurine and  $\gamma$ -aminobutyric acid by high-performance liquid chromatography-fluorescence detection with o-phthalaldehyde-mercaptoethanol derivatization with an emphasis on microdialysis sample. *J Chromatogr B Biomed Sci Appl* 757, 277–283.
- Pignatelli A, and Belluzzi O, 2017. Dopaminergic Neurons in the Main Olfactory Bulb: An Overview from an Electrophysiological Perspective. *Front Neuroanat* 11, 7.
- Piltonen M, Beshpalov MM, Ervasti D, Matilainen T, et al., 2009. Heparin-binding determinants of GDNF reduce its tissue distribution but are beneficial for the protection of nigral dopaminergic neurons. *Exp Neurol* 219, 499–506.
- Plaza-Menacho I, Burzynski GM, Groot JW de, Eggen BJL, et al., 2006. Current concepts in RET-related genetics, signaling and therapeutics. *Trends Genet* 22, 627–636.
- Politis M, 2014. Neuroimaging in Parkinson disease: from research setting to clinical practice. *Nat Rev Neurol* 10, 708–722.
- Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, et al., 1997. Mutation in the  $\alpha$ -Synuclein Gene Identified in Families with Parkinson's Disease. *Science (80- )* 276, 2045–2047.
- Porras G, Li Q, and Bezard E, 2012. Modeling Parkinson's Disease in Primates: The MPTP Model. *Cold Spring Harb Perspect Med* 2, 1–10.
- Postuma RB, Aarsland D, Barone P, Burn DJ, et al., 2012. Identifying prodromal Parkinson's disease: Pre-Motor disorders in Parkinson's disease. *Mov Disord* 27, 617–626.
- Postuma RB, Berg D, Stern M, Poewe W, et al., 2015. MDS clinical diagnostic criteria for Parkinson's disease. *Mov Disord*.
- Pothos EN, Davila V, and Sulzer D, 1998. Presynaptic recording of quanta from midbrain dopamine neurons and modulation of the quantal size. *J Neurosci* 18, 4106–4118.
- Pouloupoulos M, Levy OA, and Alcalay RN, 2012. The neuropathology of genetic Parkinson's disease. *Mov Disord* 27, 831–842.
- Prediger RDS, Aguiar AS, Moreira ELG, Matheus FC, et al., 2011. The Intranasal Administration of 1-Methyl-4-Phenyl-1,2,3,6-Tetrahydropyridine (MPTP): A New Rodent Model to Test Palliative and Neuroprotective Agents for Parkinsons disease. *Curr Pharm Des*.
- Prediger RDS, Aguiar AS, Rojas-Mayorquin AE, Figueiredo CP, et al., 2010. Single Intranasal Administration of 1-Methyl-4-Phenyl-1,2,3,6-Tetrahydropyridine in C57BL/6 Mice Models Early Preclinical Phase of Parkinson's Disease. *Neurotox Res* 17, 114–129.
- Prediger RDS, Batista LC, Medeiros R, Pandolfo P, et al., 2006. The risk is in the air: Intranasal administration of MPTP to rats reproducing clinical features of Parkinson's disease. *Exp Neurol* 202, 391–403.
- Prensa L, and Parent A, 2001. The Nigrostriatal Pathway in the Rat: A Single-Axon Study of the Relationship between Dorsal and Ventral Tier Nigral Neurons and the Striosome/Matrix Striatal Compartments. *J Neurosci* 21, 7247–7260.
- Pringsheim T, Jette N, Frolkis A, and Steeves TDL, 2014. The prevalence of Parkinson's disease: A systematic review and meta-analysis. *Mov Disord*.
- Puopolo M, 2019. The hypothalamic-spinal dopaminergic system: a target for pain modulation. *Neural Regen Res* 14, 925–930.
- Qu S, Ondo WG, Zhang X, Xie WJ, et al., 2006. Projections of diencephalic dopamine neurons into the spinal cord in mice. *Exp Brain Res* 168, 152–156.
- Ramirez AD, Hao J, Kethchum E, Kwok K, et al., 2004. 536. Design and production of an AAV2 vector encoding a chimeric cDNA allowing efficient secretion of functional human neurturin from human cells. *Mol Ther* 9, S202–S203.
- Ramirez AD, Wong SK-F, and Menniti FS, 2003. Pramipexole inhibits MPTP toxicity in mice by dopamine D3 receptor dependent and independent mechanisms. *Eur J Pharmacol* 475, 29–35.
- Rannikko EH, Weber SS, and Kahle PJ, 2015. Exogenous  $\alpha$ -synuclein induces toll-like receptor 4 dependent inflammatory responses in astrocytes. *BMC Neurosci* 16, 57.
- Rappold PM, Cui M, Chesser AS, Tibbett J, et al., 2011. Paraquat neurotoxicity is mediated by the dopamine transporter and organic cation transporter-3. *Proc Natl Acad Sci* 108, 20766–20771.
- Raykhel I, Alanen H, Salo K, Jurvansuu J, et al., 2007. A molecular specificity code for the three mammalian KDEL receptors. *J Cell Biol* 179, 1193–204.
- Recasens A, Dehay B, Bové J, Carballo-Carbajal I, et al., 2014. Lewy body extracts from Parkinson disease brains trigger  $\alpha$ -synuclein

- pathology and neurodegeneration in mice and monkeys. *Ann Neurol* 75, 351–362.
- Ren X, Zhang T, Gong X, Hu G, et al., 2013. AAV2-mediated striatum delivery of human CDNF prevents the deterioration of midbrain dopamine neurons in a 6-hydroxydopamine induced parkinsonian rat model. *Exp Neurol* 248.
- Renko J-M, Bäck S, Voutilainen MH, Piepponen TP, et al., 2018. Mesencephalic Astrocyte-Derived Neurotrophic Factor (MANF) Elevates Stimulus-Evoked Release of Dopamine in Freely-Moving Rats. *Mol Neurobiol*.
- Rey NL, Steiner JA, Maroof N, Luk KC, et al., 2016. Widespread transneuronal propagation of  $\alpha$ -synucleinopathy triggered in olfactory bulb mimics prodromal Parkinson's disease. *J Exp Med* 213, 1759–1778.
- Riccio A, Pierchala BA, Ciarallo CL, and Ginty DD, 1997. An NGF-TrkA-Mediated Retrograde Signal to Transcription Factor CREB in Sympathetic Neurons. *Science (80- )* 277, 1097–1100.
- Rice ME, and Cragg SJ, 2008. Dopamine spillover after quantal release: Rethinking dopamine transmission in the nigrostriatal pathway. *Brain Res Rev* 58, 303–313.
- Richardson DS, Lai AZ, and Mulligan LM, 2006. RET ligand-induced internalization and its consequences for downstream signaling. *Oncogene* 25, 3206–3211.
- Richardson DS, Rodrigues DM, Hyndman BD, Crupi MJF, et al., 2012. Alternative splicing results in RET isoforms with distinct trafficking properties. *Mol Biol Cell* 23, 3838–3850.
- Richardson JR, Quan Y, Sherer TB, Greenamyre JT, et al., 2005. Paraquat Neurotoxicity is Distinct from that of MPTP and Rotenone. *Toxicol Sci* 88, 193–201.
- Richman C, Rashid S, Prashar S, Mishra R, et al., 2018. C. elegans MANF Homolog Is Necessary for the Protection of Dopaminergic Neurons and ER Unfolded Protein Response. *Front Neurosci* 12, 1–11.
- Rickard SM, Mummery RS, Mulloy B, and Rider CC, 2003. The binding of human glial cell line-derived neurotrophic factor to heparin and heparan sulfate: importance of 2-O-sulfate groups and effect on its interaction with its receptor, GFR $\alpha$ 1. *Glycobiology* 13, 419–426.
- Robertson HA, 1992. Dopamine receptor interactions: some implications for the treatment of Parkinson's disease. *Trends Neurosci* 15, 201–206.
- Rocha EM, De Miranda B, and Sanders LH, 2018. Alpha-synuclein: Pathology, mitochondrial dysfunction and neuroinflammation in Parkinson's disease. *Neurobiol Dis* 109, 249–257.
- Rocha SM, Cristovão AC, Campos FL, Fonseca CP, et al., 2012. Astrocyte-derived GDNF is a potent inhibitor of microglial activation. *Neurobiol Dis* 47, 407–415.
- Rolinski M, Griffanti L, Piccini P, Roussakis AA, et al., 2016. Basal ganglia dysfunction in idiopathic REM sleep behaviour disorder parallels that in early Parkinson's disease. *Brain* 139, 2224–2234.
- Rolinski M, Griffanti L, Szeewczyk-Krolkowski K, Menke RAL, et al., 2015. Aberrant functional connectivity within the basal ganglia of patients with Parkinson's disease. *NeuroImage Clin* 8, 126–132.
- Rommelfanger KS, and Wichmann T, 2010. Extrastriatal dopaminergic circuits of the basal ganglia. *Front Neuroanat* 4, 139.
- Ron D, and Walter P, 2007. Signal integration in the endoplasmic reticulum unfolded protein response. *Nat Rev Mol Cell Biol* 8, 519–529.
- Rosenblad C, Georgievska B, and Kirik D, 2003. Long-term striatal overexpression of GDNF selectively downregulates tyrosine hydroxylase in the intact nigrostriatal dopamine system. *Eur J Neurosci* 17, 260–270.
- Rosenblad C, Grønborg M, Hansen C, Blom N, et al., 2000. In Vivo Protection of Nigral Dopamine Neurons by Lentiviral Gene Transfer of the Novel GDNF-Family Member Neublastin/Artemin. *Mol Cell Neurosci* 15, 199–214.
- Rosenblad C, Kirik D, Devaux B, Moffat B, et al., 1999. Protection and regeneration of nigral dopaminergic neurons by neurturin or GDNF in a partial lesion model of Parkinson's disease after administration into the striatum or the lateral ventricle. *Eur J Neurosci* 11, 1554–1566.
- Rossi J, Herzog K-H, Vöikar V, Hiltunen PH, et al., 2003. Alimentary tract innervation deficits and dysfunction in mice lacking GDNF family receptor  $\alpha$ 2. *J Clin Invest* 112, 707–716.
- Rossi J, Luukko K, Poteryaev D, Laurikainen A, et al., 1999. Retarded Growth and Deficits in the Enteric and Parasympathetic Nervous System in Mice Lacking GFR $\alpha$ 2, a Functional Neurturin Receptor. *Neuron* 22, 243–252.
- Rossi J, Tomac A, Saarma M, and Airaksinen MS, 2000. Distinct roles for GFR $\alpha$ 1 and GFR $\alpha$ 2 signalling in different cranial parasympathetic ganglia in vivo. *Eur J Neurosci* 12, 3944–3952.
- Rubinsztein DC, 2006. The roles of intracellular protein-degradation pathways in neurodegeneration. *Nature* 443, 780–786.
- Runeberg-Roos P, Piccinini E, Penttinen A-M, Mätlik K, et al., 2016. Developing therapeutically more efficient Neurturin variants for treatment of Parkinson's disease. *Neurobiol Dis* 96, 335–345.
- Ryan BJ, Hoek S, Fon EA, and Wade-Martins R, 2015. Mitochondrial dysfunction and mitophagy in Parkinson's: From familial to sporadic disease. *Trends Biochem Sci*.
- Saarma M, Mati K, Mehias P, and Michailovich BM, 2014. Methods of facilitating neural cell survival using GDNF family ligand (GFL) mimetics or RET signalling pathway activators. 8,901,129.
- Saavedra A, Baltazar G, and Duarte EP, 2008. Driving GDNF expression: The green and the red traffic lights. *Prog Neurobiol* 86, 186–215.
- Saavedra A, Baltazar G, and Duarte EP, 2007. Interleukin-1 $\beta$  mediates GDNF up-regulation upon dopaminergic injury in ventral midbrain cell cultures. *Neurobiol Dis* 25, 92–104.
- Saeed U, Compagnone J, Aviv RI, Strafella AP, et al., 2017. Imaging biomarkers in Parkinson's disease and Parkinsonian syndromes: current and emerging concepts. *Transl Neurodegener* 6, 1–25.
- Saiki M, Baker A, Williams-Gray CH, Foltynie T, et al., 2010. Association of the human leucocyte antigen region with susceptibility to Parkinson's disease. *J Neurol Neurosurg Psychiatry* 81, 890–891.
- Sainio K, Suvanto P, Davies J, Wartiovaara J, et al., 1997. Glial-cell-line-derived neurotrophic factor is required for bud initiation from ureteric epithelium. *Development* 124, 4077–4087.
- Salvatore MF, Ai Y, Fischer B, Zhang AM, et al., 2006. Point source concentration of GDNF may explain failure of phase II clinical trial. *Exp Neurol* 202, 497–505.



- Salvatore MF, Gerhardt GA, Dayton RD, Klein RL, et al., 2009. Bilateral effects of unilateral GDNF administration on dopamine- and GABA-regulating proteins in the rat nigrostriatal system. *Exp Neurol* 219, 197–207.
- Salvatore MF, Zhang J-L, Large DM, Wilson PE, et al., 2004. Striatal GDNF administration increases tyrosine hydroxylase phosphorylation in the rat striatum and substantia nigra. *J Neurochem* 90, 245–254.
- Sánchez MP, Silos-Santiago I, Frisén J, He B, et al., 1996. Renal agenesis and the absence of enteric neurons in mice lacking GDNF. *Nature* 382, 70–73.
- Sanicola M, Hession C, Worley D, Carmillo P, et al., 1997. Glial cell line-derived neurotrophic factor-dependent RET activation can be mediated by two different cell-surface accessory proteins. *Proc Natl Acad Sci U S A* 94, 6238–43.
- Sano I, Gamo T, Kakimoto Y, Taniguchi K, et al., 1959. Distribution of catechol compounds in human brain. *Biochim Biophys Acta* 32, 586–587.
- Sarabi A, Hoffer BJ, Olson L, and Morales M, 2003. Glial cell line neurotrophic factor-family receptor  $\alpha$ -1 is present in central neurons with distinct phenotypes. *Neuroscience* 116.
- Sarabi A, Hoffer BJ, Olson L, and Morales M, 2001. GFR $\alpha$ -1 mRNA in dopaminergic and nondopaminergic neurons in the substantia nigra and ventral tegmental area. *J Comp Neurol* 441, 106–117.
- Sauer H, and Oertel WH, 1994. Progressive degeneration of nigrostriatal dopamine neurons following intrastriatal terminal lesions with 6-hydroxydopamine: A combined retrograde tracing and immunocytochemical study in the rat. *Neuroscience* 59, 401–415.
- Savitt J, Singh D, Zhang C, Chen L-C, et al., 2012. The In Vivo Response of Stem and Other Undifferentiated Spermatogonia to the Reversible Inhibition of Glial Cell Line-Derived Neurotrophic Factor Signaling in the Adult. *Stem Cells* 30, 732–740.
- Sawada M, Sun W, Hayes P, Leskov K, et al., 2003. Ku70 suppresses the apoptotic translocation of Bax to mitochondria. *Nat Cell Biol* 5, 320–329.
- Schallert T, Fleming SM, Leasure JL, Tillerson JL, et al., 2000. CNS plasticity and assessment of forelimb sensorimotor outcome in unilateral rat models of stroke, cortical ablation, parkinsonism and spinal cord injury. *Neuropharmacology* 39, 777–787.
- Schallert T, and Woodlee M, 2005. Orienting and placing, in: *The Behavior of the Laboratory Rat: A Handbook with Tests*. Oxford University Press Oxford, pp. 129–140.
- Schapira AH V, Cooper JM, Dexter D, Jenner P, et al., 1989. MITOCHONDRIAL COMPLEX I DEFICIENCY IN PARKINSON'S DISEASE. *Lancet* 333, 1269.
- Scharr DG, Sieber B-A, Dreyfus CF, and Black IB, 1993. Regional and Cell-Specific Expression of GDNF in Rat Brain. *Exp Neurol* 124, 368–371.
- Schendzielorz N, Männistö PT, Karayiorgou M, Gogos JA, et al., 2012. A Transient Inhibition and Permanent Lack of Catechol-O-Methyltransferase have Minor Effects on Feeding Pattern of Female Rodents. *Basic Clin Pharmacol Toxicol* 110, 307–313.
- Schlösser B, Kudernatsch MB, Sutor B, and ten Bruggencate G, 1995.  $\delta$ ,  $\mu$ , and  $\kappa$  opioid receptor agonists inhibit dopamine overflow in rat neostriatal slices. *Neurosci Lett* 191, 126–130.
- Schmitz Y, Benoit-Marand M, Gonon F, and Sulzer D, 2003. Presynaptic regulation of dopaminergic neurotransmission. *J Neurochem* 87, 273–289.
- Schrag A, and Schott JM, 2006. Epidemiological, clinical, and genetic characteristics of early-onset parkinsonism. *Lancet Neurol* 5, 355–363.
- Schuchardt A, D'Agati V, Larsson-Blomberg L, Costantini F, et al., 1994. Defects in the kidney and enteric nervous system of mice lacking the tyrosine kinase receptor Ret. *Nature* 367, 380–383.
- Schuchardt A, D'Agati V, Pachnis V, and Costantini F, 1996. Renal agenesis and hypodysplasia in ret-k- mutant mice result from defects in ureteric bud development. *Development* 122, 1919–1929.
- Schultz W, 2007. Multiple Dopamine Functions at Different Time Courses. *Annu Rev Neurosci* 30, 259–288.
- Scott DA, Tabarean I, Tang Y, Cartier A, et al., 2010. A Pathologic Cascade Leading to Synaptic Dysfunction in  $\alpha$ -Synuclein-Induced Neurodegeneration. *J Neurosci* 30, 8083–8095.
- Scott RP, Eketjäll S, Aineskog H, and Ibáñez CF, 2005. Distinct turnover of alternatively spliced isoforms of the RET kinase receptor mediated by differential recruitment of the Cbl ubiquitin ligase. *J Biol Chem* 280, 13442–13449.
- Shahmoradian SH, Lewis AJ, Genoud C, Hench J, et al., 2019. Lewy pathology in Parkinson's disease consists of crowded organelles and lipid membranes. *Nat Neurosci* 22.
- Shakya R, Watanabe T, and Costantini F, 2005. The Role of GDNF/Ret Signaling in Ureteric Bud Cell Fate and Branching Morphogenesis. *Dev Cell* 8, 65–74.
- Shen Y, Sun A, Wang Y, Cha D, et al., 2012. Upregulation of mesencephalic astrocyte-derived neurotrophic factor in glial cells is associated with ischemia-induced glial activation. *J Neuroinflammation* 9, 254.
- Sherer TB, Fiske BK, Svendsen CN, Lang AE, et al., 2006. Crossroads in GDNF therapy for Parkinson's disease. *Mov Disord* 21, 136–141.
- Shimizu K, Ohtaki K, Matsubara K, Aoyama K, et al., 2001. Carrier-mediated processes in blood–brain barrier penetration and neural uptake of paraquat. *Brain Res* 906, 135–142.
- Shimozawa A, Ono M, Takahara D, Tarutani A, et al., 2017. Propagation of pathological  $\alpha$ -synuclein in marmoset brain. *Acta Neuropathol Commun* 5, 1–14.
- Sian-Hülsman J, Mandel S, Youdim MBH, and Riederer P, 2011. The relevance of iron in the pathogenesis of Parkinson's disease. *J Neurochem* 118, 939–957.
- Sidorova YA, Beshpalov MM, Wong AW, Kambur O, et al., 2017. A Novel Small Molecule GDNF Receptor RET Agonist, BT13, Promotes Neurite Growth from Sensory Neurons in Vitro and Attenuates Experimental Neuropathy in the Rat. *Front Pharmacol* 8, 365.
- Sidorova YA, and Saarna M, 2020. Small Molecules and Peptides Targeting Glial Cell Line-Derived Neurotrophic Factor Receptors for the Treatment of Neurodegeneration. *Int J Mol Sci* 21, 6575.
- Sidransky E, and Lopez G, 2012. The link between the GBA gene and parkinsonism. *Lancet Neurol* 11, 986–998.
- Sierra A, Encinas JM, Deudero JJP, Chancey JH, et al., 2010. Microglia Shape Adult Hippocampal Neurogenesis through Apoptosis-Coupled Phagocytosis. *Cell Stem Cell* 7, 483–495.
- Simola N, Morelli M, and Carta AR, 2007. The 6-Hydroxydopamine model of Parkinson's disease. *Neurotox Res* 11, 151–167.



- Simón-Sánchez J, Schulte C, Bras JM, Sharma M, et al., 2009. Genome-wide association study reveals genetic risk underlying Parkinson's disease. *Nat Genet* 41, 1308–1312.
- Sjöstrand D, Carlsson J, Paratcha G, Persson B, et al., 2007. Disruption of the GDNF binding site in NCAM dissociates ligand binding and homophilic cell adhesion. *J Biol Chem* 282, 12734–12740.
- Skagerberg G, and Lindvall O, 1985. Organization of diencephalic dopamine neurones projecting to the spinal cord in the rat. *Brain Res* 342, 340–351.
- Slevin JT, Gash DM, Smith CD, Gerhardt GA, et al., 2007. Unilateral intraputamenal glial cell line-derived neurotrophic factor in patients with Parkinson disease: response to 1 year of treatment and 1 year of withdrawal. *J Neurosurg* 106, 614–620.
- Slevin JT, Gerhardt GA, Smith CD, Gash DM, et al., 2005. Improvement of bilateral motor functions in patients with Parkinson disease through the unilateral intraputamenal infusion of glial cell line-derived neurotrophic factor. *J Neurosurg* 102, 216–222.
- Smith RC, O'Bryan LM, Mitchell PJ, Leung D, et al., 2015. Increased brain bio-distribution and chemical stability and decreased immunogenicity of an engineered variant of GDNF. *Exp Neurol* 267, 165–176.
- Smith SB, Reenilä I, Männistö PT, Slade GD, et al., 2014. Epistasis between polymorphisms in COMT, ESR1, and GCH1 influences COMT enzyme activity and pain. *Pain* 155, 2390–2399.
- Smith Y, Lavoie B, Dumas J, and Parent A, 1989. Evidence for a distinct nigropallidal dopaminergic projection in the squirrel monkey. *Brain Res* 482, 381–386.
- Smolders I, Klippel N De, Sarre S, Ebinger G, et al., 1995. Tonic GABA-ergic modulation of striatal dopamine release studied by in vivo microdialysis in the freely moving rat. *Eur J Pharmacol* 284, 83–91.
- Snead D, and Eliezer D, 2014. Alpha-synuclein function and dysfunction on cellular membranes. *Exp Neurol* 23, 292–313.
- Snyder GL, Allen PB, Fienberg AA, Valle CG, et al., 2000. Regulation of Phosphorylation of the GluR1 AMPA Receptor in the Neostriatum by Dopamine and Psychostimulants In Vivo. *J Neurosci* 20, 4480–4488.
- Sousa-Victor P, Jasper H, and Neves J, 2018. Trophic Factors in Inflammation and Regeneration: The Role of MANF and CDNF. *Front Physiol* 9, 1629.
- Spalding KL, Bergmann O, Alkass K, Bernard S, et al., 2013. Dynamics of Hippocampal Neurogenesis in Adult Humans. *Cell* 153, 1219–1227.
- Spalding KL, Bhardwaj RD, Buchholz BA, Druid H, et al., 2005. Retrospective Birth Dating of Cells in Humans. *Cell* 122, 133–143.
- Spillantini MG, Crowther RA, Jakes R, Hasegawa M, et al., 1998.  $\alpha$ -Synuclein in filamentous inclusions of Lewy bodies from Parkinson's disease and dementia with Lewy bodies. *Proc Natl Acad Sci* 95, 6469–6473.
- Spillantini MG, Schmidt ML, Lee VM-Y, Trojanowski JQ, et al., 1997.  $\alpha$ -Synuclein in Lewy bodies. *Nature* 388, 839–840.
- Sprenkle NT, Sims SG, Sánchez CL, and Meares GP, 2017. Endoplasmic reticulum stress and inflammation in the central nervous system. *Mol Neurodegener* 12, 42.
- Springer JE, Mu X, Bergmann LW, and Trojanowski JQ, 1994. Expression of GDNF mRNA in Rat and Human Nervous Tissue. *Exp Neurol* 127, 167–170.
- Springer JE, Seeburger JL, Jin HE, Gabrea A, et al., 1995. cDNA sequence and differential mRNA regulation of two forms of glial cell line-derived neurotrophic factor in Schwann cells and rat skeletal muscle. *Exp Neurol* 131, 47–52.
- Stanford JA, Salvatore MF, Joyce BM, Zhang H, et al., 2007. Bilateral effects of unilateral intrastriatal GDNF on locomotor-excited and nonlocomotor-related striatal neurons in aged F344 rats. *Neurobiol Aging* 28, 156–165.
- Stanic D, Finkelstein DI, Bourke DW, Drago J, et al., 2003a. Timecourse of striatal re-innervation following lesions of dopaminergic SNpc neurons of the rat. *Eur J Neurosci* 18, 1175–1188.
- Stanic D, Parish CL, Zhu WM, Krstew E V, et al., 2003b. Changes in function and ultrastructure of striatal dopaminergic terminals that regenerate following partial lesions of the SNpc. *J Neurochem* 86, 329–343.
- Stuber GD, Hnasko TS, Britt JP, Edwards RH, et al., 2010. Dopaminergic Terminals in the Nucleus Accumbens But Not the Dorsal Striatum Corelease Glutamate. *J Neurosci* 30, 8229–8233.
- Su X, Fischer DL, Li X, Bankiewicz K, et al., 2017. Alpha-Synuclein mRNA Is Not Increased in Sporadic PD and Alpha-Synuclein Accumulation Does Not Block GDNF Signaling in Parkinson's Disease and Disease Models. *Mol Ther* 25, 2231–2235.
- Su X, Maguire-Zeiss KA, Giuliano R, Prifti L, et al., 2008. Synuclein activates microglia in a model of Parkinson's disease. *Neurobiol Aging* 29, 1690–1701.
- Sulzer D, 2011. How Addictive Drugs Disrupt Presynaptic Dopamine Neurotransmission. *Neuron* 69, 628–649.
- Sulzer D, Cragg SJ, and Rice ME, 2016. Chapter 19 - Regulation of Extracellular Dopamine: Release and Uptake, in: Steiner, H., Tseng, K.Y. (Eds.), *Handbook of Basal Ganglia Structure and Function, Second Edition, Handbook of Behavioral Neuroscience*. Elsevier, pp. 373–402.
- Sulzer D, and Rayport S, 1990. Amphetamine and other psychostimulants reduce pH gradients in midbrain dopaminergic neurons and chromaffin granules: a mechanism of action. *Neuron* 5, 797–808.
- Sulzer D, and Zecca L, 1999. Intraneuronal dopamine-quinone synthesis: A review. *Neurotox Res* 1, 181–195.
- Sun X, Zhao Y, and Wolf ME, 2005. Dopamine Receptor Stimulation Modulates AMPA Receptor Synaptic Insertion in Prefrontal Cortex Neurons. *J Neurosci* 25, 7342–7351.
- Surmeier DJ, Ding J, Day M, Wang Z, et al., 2007. D1 and D2 dopamine-receptor modulation of striatal glutamatergic signaling in striatal medium spiny neurons. *Trends Neurosci* 30, 228–235.
- Surmeier DJ, and Schumacker PT, 2013. Calcium, Bioenergetics, and Neuronal Vulnerability in Parkinson's Disease. *J Biol Chem* 288, 10736–10741.
- Suvanto P, Hiltunen JO, Arumäe U, Moshnyakov M, et al., 1996. Localization of glial cell line-derived neurotrophic factor (GDNF) mRNA in embryonic rat by in situ hybridization. *Eur J Neurosci* 8, 816–822.
- Suvanto P, Wartiovaara K, Lindahl M, Arumäe U, et al., 1997. Cloning, mRNA distribution and chromosomal localisation of the gene for glial cell line-derived neurotrophic factor receptor beta, a homologue to GDNFR-alpha. *Hum Mol Genet* 6, 1267–1273.
- Svensson E, Horváth-Puhó E, Thomsen RW, Djurhuus JC, et al., 2015. Vagotomy and subsequent risk of Parkinson's disease. *Ann Neurol* 78,

- 522–529.
- Svingos AL, Chavkin C, Colago EEO, and Pickel VM, 2001. Major coexpression of  $\kappa$ -opioid receptors and the dopamine transporter in nucleus accumbens axonal profiles. *Synapse* 42, 185–192.
- Svingos AL, Clarke CL, and Pickel VM, 1999. Localization of the  $\delta$ -opioid receptor and dopamine transporter in the nucleus accumbens shell: Implications for opiate and psychostimulant cross-sensitization. *Synapse* 34, 1–10.
- Swahn C-G, and Wiesel F-A, 1976. Determination of conjugated monoamine metabolites in brain tissue. *J Neural Transm* 39, 281–290.
- Swanson LW, 1982. The projections of the ventral tegmental area and adjacent regions: a combined fluorescent retrograde tracer and immunofluorescence study in the rat. *Brain Res Bull* 9, 321–53.
- Szewczyk-Krolkowski K, Menke RAL, Rolinski M, Duff E, et al., 2014. Functional connectivity in the basal ganglia network differentiates PD patients from controls. *Neurology* 83, 208–214.
- Taber KH, and Hurley R a, 2014. Volume transmission in the brain: beyond the synapse. *J Neuropsychiatry Clin Neurosci* 26, 1–4.
- Tadimalla A, Belmont PJ, Thuerauf DJ, Glassy MS, et al., 2008. Mesencephalic astrocyte-derived neurotrophic factor is an ischemia-inducible secreted endoplasmic reticulum stress response protein in the heart. *Circ Res* 103, 1249–1258.
- Tagliaferro P, and Burke RE, 2016. Retrograde Axonal Degeneration in Parkinson Disease. *J Parkinsons Dis*.
- Tahira T, Ishizaka Y, Itoh F, Sugimura T, et al., 1990. Characterization of ret proto-oncogene mRNAs encoding two isoforms of the protein product in a human neuroblastoma cell line. *Oncogene* 5, 97–102.
- Takahashi M, 2001. The GDNF/RET signaling pathway and human diseases. *Cytokine Growth Factor Rev* 12, 361–73.
- Takahashi M, Ritz J, and Cooper GM, 1985. Activation of a novel human transforming gene, ret, by DNA rearrangement. *Cell* 42, 581–588.
- Tanner CM, Freya K, Webster RG, Hoppin J, et al., 2011. Rotenone, Paraquat, and Parkinson's Disease. *Environ Health Perspect* 119, 866–872.
- Tansey MG, Baloh RH, Milbrandt J, and Johnson EM, 2000. GFR $\alpha$ -mediated localization of RET to lipid rafts is required for effective downstream signaling, differentiation, and neuronal survival. *Neuron* 25, 611–23.
- Taraviras S, Marcos-Gutierrez C V, Durbec P, Jani H, et al., 1999. Signalling by the RET receptor tyrosine kinase and its role in the development of the mammalian enteric nervous system. *Development* 126, 2785–97.
- Tatarewicz SM, Wei X, Gupta S, Masterman D, et al., 2007. Development of a Maturing T-Cell-Mediated Immune Response in Patients with Idiopathic Parkinson's Disease Receiving r-metHuGDNF Via Continuous Intrapaternal Infusion. *J Clin Immunol* 27, 620–627.
- Tecuapetla F, Patel JC, Xenias H, English D, et al., 2010. Glutamatergic Signaling by Mesolimbic Dopamine Neurons in the Nucleus Accumbens. *J Neurosci* 30, 7105–7110.
- Tepper JM, Wilson CJ, and Koós T, 2008. Feedforward and feedback inhibition in neostriatal GABAergic spiny neurons. *Brain Res Rev* 58, 272–281.
- Thakur P, Breger LS, Lundblad M, Wan OW, et al., 2017. Modeling Parkinson's disease pathology by combination of fibril seeds and  $\alpha$ -synuclein overexpression in the rat brain. *Proc Natl Acad Sci*.
- Thompson J, Doxakis E, Piñón LGP, Strachan P, et al., 1998. GFR $\alpha$ -4, a New GDNF Family Receptor. *Mol Cell Neurosci* 11, 117–126.
- Threlfell S, Lalic T, Platt NJ, Jennings KA, et al., 2012. Striatal dopamine release is triggered by synchronized activity in cholinergic interneurons. *Neuron* 75, 58–64.
- Tillerson JL, Cohen AD, Philhower J, Miller GW, et al., 2001. Forced limb-use effects on the behavioral and neurochemical effects of 6-hydroxydopamine. *J Neurosci* 21, 4427–4435.
- Timmerman W, and Abercrombie ED, 1996. Amphetamine-induced release of dendritic dopamine in substantia nigra pars reticulata: D1-mediated behavioral and electrophysiological effects. *Synapse* 23, 280–291.
- Tokugawa K, Yamamoto K, Nishiguchi M, Sekine T, et al., 2003. XIB4035, a novel nonpeptidyl small molecule agonist for GFR $\alpha$ -1. *Neurochem Int* 42, 81–86.
- Tolosa E, Wenning G, and Poewe W, 2006. The diagnosis of Parkinson's disease. *Lancet Neurol* 5, 75–86.
- Tomac A, Lindqvist E, Lin L-FH, Ögren SO, et al., 1995. Protection and repair of the nigrostriatal dopaminergic system by GDNF in vivo. *Nature* 373, 335–339.
- Tomac A, Widenfalk J, Lin LF, Kohno T, et al., 1995b. Retrograde axonal transport of glial cell line-derived neurotrophic factor in the adult nigrostriatal system suggests a trophic role in the adult. *Proc Natl Acad Sci U S A* 92, 8274–8.
- Tomac AC, Agulnick AD, Haughey N, Chang C-F, et al., 2002. Effects of cerebral ischemia in mice deficient in Persephin. *Proc Natl Acad Sci U S A* 99, 9521–9526.
- Tozzi A, de Iure A, Di Filippo M, Tantucci M, et al., 2011. The Distinct Role of Medium Spiny Neurons and Cholinergic Interneurons in the D2/A2A Receptor Interaction in the Striatum: Implications for Parkinson's Disease. *J Neurosci* 31, 1850–1862.
- Treanor JJS, Goodman L, de Sauvage F, Stone DM, et al., 1996. Characterization of a multicomponent receptor for GDNF. *Nature* 382, 80–83.
- Tritsch NX, Ding JB, and Sabatini BL, 2012. Dopaminergic neurons inhibit striatal output through non-canonical release of GABA. *Nature* 490, 262–266.
- Tritsch NX, Oh WJ, Gu C, and Sabatini BL, 2014. Midbrain dopamine neurons sustain inhibitory transmission using plasma membrane uptake of GABA, not synthesis. *Elife* 3, e01936.
- Tronci E, Shin E, Björklund A, and Carta M, 2012. Amphetamine-induced rotation and l-DOPA-induced dyskinesia in the rat 6-OHDA model: A correlation study. *Neurosci Res* 73, 168–172.
- Trupp M, Arenas E, Fainzilber M, Nilsson A-S, et al., 1996. Functional receptor for GDNF encoded by the c-ret proto-oncogene. *Nature* 381, 785–789.
- Trupp M, Belluardo N, Funakoshi H, and Ibáñez CF, 1997. Complementary and overlapping expression of glial cell line-derived neurotrophic factor (GDNF), c-ret proto-oncogene, and GDNF receptor- $\alpha$  indicates multiple mechanisms of trophic actions in the adult rat CNS. *J Neurosci* 17, 3554–67.
- Trupp M, Jonsson P, Öhrfelt A, Zetterberg H, et al., 2014. Metabolite and Peptide Levels in Plasma and CSF Differentiating Healthy Controls

- from Patients with Newly Diagnosed Parkinson's Disease. *J Parkinsons Dis* 4, 549–560.
- Trupp M, Raynoschek C, Belluardo N, and Ibáñez CF, 1998. Multiple GPI-Anchored Receptors Control GDNF-Dependent and Independent Activation of the c-Ret Receptor Tyrosine Kinase. *Mol Cell Neurosci* 11, 47–63.
- Trupp M, Rydén M, Jörnvall H, Funakoshi H, et al., 1995. Peripheral expression and biological activities of GDNF, a new neurotrophic factor for avian and mammalian peripheral neurons. *J Cell Biol* 130, 137–148.
- Tseng K-Y, Anttila JE, Khodosevich K, Tuominen RK, et al., 2018. MANF Promotes Differentiation and Migration of Neural Progenitor Cells with Potential Neural Regenerative Effects in Stroke. *Mol Ther* 26, 238–255.
- Tseng K-Y, Danilova T, Domanskyi A, Saarma M, et al., 2017. MANF Is Essential for Neurite Extension and Neuronal Migration in the Developing Cortex. *eNeuro* 4.
- Tsui CC, Gabreski NA, Hein SJ, and Pierchala BA, 2015. Lipid Rafts Are Physiologic Membrane Microdomains Necessary for the Morphogenic and Developmental Functions of Glial Cell Line-Derived Neurotrophic Factor In Vivo. *J Neurosci* 35, 13233–43.
- Tsui CC, and Pierchala BA, 2010. The Differential Axonal Degradation of Ret Accounts for Cell-Type-Specific Function of Glial Cell Line-Derived Neurotrophic Factor as a Retrograde Survival Factor. *J Neurosci* 30, 5149–5158.
- Tsuzuki T, Takahashi M, Asai N, Iwashita T, et al., 1995. Spatial and temporal expression of the ret proto-oncogene product in embryonic, infant and adult rat tissues. *Oncogene* 10, 191–198.
- Turconi G, Kopra J, Vöikar V, Kuleshkaya N, et al., 2020. Chronic 2-Fold Elevation of Endogenous GDNF Levels Is Safe and Enhances Motor and Dopaminergic Function in Aged Mice. *Mol Ther - Methods Clin Dev* 17, 831–842.
- Tyson T, Steiner JA, and Brundin P, 2016. Sorting out release, uptake and processing of alpha-synuclein during prion-like spread of pathology. *J Neurochem*.
- Uesaka T, Nagashimada M, Yonemura S, and Enomoto H, 2008. Diminished Ret expression compromises neuronal survival in the colon and causes intestinal aganglionosis in mice. *J Clin Invest* 118, 1890–1898.
- Um JW, Im E, Lee HJ, Min B, et al., 2010. Parkin Directly Modulates 26S Proteasome Activity. *J Neurosci* 30, 11805–11814.
- Ungerstedt U, 1971. Postsynaptic Supersensitivity after 6-Hydroxy-dopamine Induced Degeneration of the Nigro-striatal Dopamine System. *Acta Physiol Scand* 82, 69–93.
- Ungerstedt U, 1968. 6-hydroxy-dopamine induced degeneration of central monoamine neurons. *Eur J Pharmacol* 5, 107–110.
- Ungerstedt U, and Arbuthnott GW, 1970a. Quantitative recording of rotational behavior in rats after 6-hydroxy-dopamine lesions of the nigrostriatal dopamine system. *Brain Res* 24, 485–493.
- Uutela P, Karhu L, Piepponen P, Käenmäki M, et al., 2009. Discovery of Dopamine Glucuronide in Rat and Mouse Brain Microdialysis Samples Using Liquid Chromatography Tandem Mass Spectrometry. *Anal Chem* 81, 427–434.
- Vallee RB, Shpetner HS, and Paschal BM, 1989. The role of dynein in retrograde axonal transport. *Trends Neurosci* 12, 66–70.
- Vallee RB, Williams JC, Varma D, and Barnhart LE, 2004. Dynein: An Ancient Motor Protein Involved in Multiple Modes of Transport. *J Neurobiol* 58, 189–200.
- Valros A, Palander P, Heinonen M, Munsterhjelm C, et al., 2015. Evidence for a link between tail biting and central monoamine metabolism in pigs (*Sus scrofa domestica*). *Physiol Behav* 143, 151–157.
- Vega QC, Worby CA, Lechner MS, Dixon JE, et al., 1996. Glial cell line-derived neurotrophic factor activates the receptor tyrosine kinase RET and promotes kidney morphogenesis. *Proc Natl Acad Sci U S A* 93, 10657–10661.
- Venton BJ, Zhang H, Garris PA, Phillips PEM, et al., 2003. Real-time decoding of dopamine concentration changes in the caudate–putamen during tonic and phasic firing. *J Neurochem* 87, 1284–1295.
- Vergo S, Johansen JL, Leist M, and Lotharius J, 2007. Vesicular monoamine transporter 2 regulates the sensitivity of rat dopaminergic neurons to disturbed cytosolic dopamine levels. *Brain Res* 1185, 18–32.
- Verity AN, Wyatt TL, Hajos B, Eglen RM, et al., 1998. Regulation of Glial Cell Line-Derived Neurotrophic Factor Release from Rat C6 Glioblastoma Cells. *J Neurochem* 70, 531–539.
- Verity AN, Wyatt TL, Lee W, Hajos B, et al., 1999. Differential regulation of glial cell line–derived neurotrophic factor (GDNF) expression in human neuroblastoma and glioblastoma cell lines. *J Neurosci Res* 55, 187–197.
- Vihavainen T, Relander TRA, Leiviskä R, Airavaara M, et al., 2008. Chronic nicotine modifies the effects of morphine on extracellular striatal dopamine and ventral tegmental GABA. *J Neurochem* 107, 844–854.
- Viisanen H, Nuotio U, Kambur O, Mahato AK, et al., 2020. Novel RET agonist for the treatment of experimental neuropathies. *Mol Pain* 16.
- Vila M, Jackson-Lewis V, Vukosavic S, Djaldetti R, et al., 2001. Bax ablation prevents dopaminergic neurodegeneration in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine mouse model of Parkinson's disease. *Proc Natl Acad Sci* 98, 2837–2842.
- Visanji NP, Brooks PL, Hazrati L-N, and Lang AE, 2013. The prion hypothesis in Parkinson's disease: Braak to the future. *Acta Neuropathol Commun* 1.
- Visanji NP, Brochie JM, Kalia L V., Koprach JB, et al., 2016.  $\alpha$ -Synuclein-Based Animal Models of Parkinson's Disease: Challenges and Opportunities in a New Era. *Trends Neurosci*.
- Visanji NP, Mollenhauer B, Beach TG, Adler CH, et al., 2017. The Systemic Synuclein Sampling Study: toward a biomarker for Parkinson's disease. *Biomark Med* 11, 359–368.
- Vlaar AMM, van Kroonenburgh MJPG, Kessels AGH, and Weber WEJ, 2007. Meta-analysis of the literature on diagnostic accuracy of SPECT in parkinsonian syndromes. *BMC Neurol* 7, 27.
- Volkman J, Albanese A, Antonini A, Chaudhuri KR, et al., 2013. Selecting deep brain stimulation or infusion therapies in advanced Parkinson's disease: an evidence-based review. *J Neurol* 260, 2701–2714.
- Volpicelli-Daley LA, Gamble KL, Schultheiss CE, Riddle DM, et al., 2014. Formation of  $\alpha$ -synuclein Lewy neurite-like aggregates in axons impedes the transport of distinct endosomes. *Mol Biol Cell* 25, 4010–4023.
- Voutilainen MH, Arumäe U, Airavaara M, and Saarma M, 2015. Therapeutic potential of the endoplasmic reticulum located and secreted CDNF/MANF family of neurotrophic factors in Parkinson's disease. *FEBS Lett* 589, 3739–3748.
- Voutilainen MH, Bäck S, Peränen J, Lindholm P, et al., 2011. Chronic infusion of CDNF prevents 6-OHDA-induced deficits in a rat model of

- Parkinson's disease. *Exp Neurol* 228, 99–108.
- Voutilainen MH, Bäck S, Pörsti E, Toppinen L, et al., 2009. Mesencephalic astrocyte-derived neurotrophic factor is neurorestorative in rat model of Parkinson's disease. *J Neurosci* 29, 9651–9659.
- Voutilainen MH, De Lorenzo F, Stepanova P, Bäck S, et al., 2017. Evidence for an Additive Neurorestorative Effect of Simultaneously Administered CDNF and GDNF in Hemiparkinsonian Rats: implications for Different Mechanism of Action. *eNeuro* 4, ENEURO0117-162017.
- Wakabayashi K, and Takahashi H, 1997. Neuropathology of autonomic nervous system in Parkinson's disease. *Eur Neurol* 38, 2–7.
- Wakabayashi K, Tanji K, Odagiri S, Miki Y, et al., 2013. The Lewy Body in Parkinson's Disease and Related Neurodegenerative Disorders. *Mol Neurobiol* 47, 495–508.
- Wake H, Moorhouse AJ, Miyamoto A, and Nabekura J, 2013. Microglia: actively surveying and shaping neuronal circuit structure and function. *Trends Neurosci* 36, 209–217.
- Walker DG, Beach TG, Xu R, Lile J, et al., 1998. Expression of the proto-oncogene Ret, a component of the GDNF receptor complex, persists in human substantia nigra neurons in Parkinson's disease. *Brain Res* 792, 207–217.
- Wang D, Hou C, Cao Y, Cheng Q, et al., 2018. XBP1 activation enhances MANF expression via binding to endoplasmic reticulum stress response elements within MANF promoter region in hepatitis B. *Int J Biochem Cell Biol* 99.
- Wang H, Hughes I, Planer W, Parsadanian A, et al., 2010. The Timing and Location of Glial Cell Line-Derived Neurotrophic Factor Expression Determine Enteric Nervous System Structure and Function. *J Neurosci* 30, 1523–1538.
- Wang H, Ke Z, Alimov A, Xu M, et al., 2014. Spatiotemporal Expression of MANF in the Developing Rat Brain. *PLoS One* 9, e90433.
- Wang J, Chen G, Lu B, and Wu CP, 2003. GDNF acutely potentiates Ca<sup>2+</sup> channels and excitatory synaptic transmission in midbrain dopaminergic neurons. *NeuroSignals* 12, 78–88.
- Wang P-C, Kuchel O, Buu NT, and Genest J, 1983. Catecholamine Glucuronidation: An Important Metabolic Pathway for Dopamine in the Rat. *J Neurochem* 40, 1435–1440.
- Wang S-Y, Ellitt CM, Malin SA, and Albers KM, 2008. Effects of the neurotrophic factor artemin on sensory afferent development and sensitivity. *Sheng Li Xue Bao* 60, 565–570.
- Warren OC, Bartus RT, Baumann TL, Factor S, et al., 2015a. Gene delivery of neurturin to putamen and substantia nigra in Parkinson disease: A double-blind, randomized, controlled trial. *Ann Neurol* 78, 248–257.
- Weinreb O, Amit T, Sagi Y, Drigues N, et al., 2009. Genomic and proteomic study to survey the mechanism of action of the anti-Parkinson's disease drug, rasagiline compared with selegiline, in the rat midbrain. *J Neural Transm* 116, 1457–1472.
- West AB, Moore DJ, Biskup S, Bugayenko A, et al., 2005. Parkinson's disease-associated mutations in leucine-rich repeat kinase 2 augment kinase activity. *Proc Natl Acad Sci* 102, 16842–16847.
- West MJ, Slomianka L, and Gundersen HJG, 1991. Unbiased stereological estimation of the total number of neurons in the subdivisions of the rat hippocampus using the optical fractionator. *Anat Rec* 231, 482–497.
- Westerink BHC, Hofsteede RM, Tuntler J, and de Vries JB, 1989. Use of Calcium Antagonism for the Characterization of Drug-Evoked Dopamine Release from the Brain of Conscious Rats Determined by Microdialysis. *J Neurochem* 52, 722–729.
- Whone AL, Luz M, Boca M, Woolley M, et al., 2019. Randomized trial of intermittent intraputamenal glial cell line-derived neurotrophic factor in Parkinson's disease. *Brain* 142, 512–525.
- Whone AL, Boca M, Luz M, Woolley M, et al., 2019b. Extended Treatment with Glial Cell Line-Derived Neurotrophic Factor in Parkinson's Disease. *J Parkinsons Dis* 9, 301–313.
- Whone AL, Watts RL, Stoessl AJ, Davis M, et al., 2003. Slower progression of Parkinson's disease with ropinirole versus levodopa: The REAL-PET study. *Ann Neurol* 54, 93–101.
- Widenfalk J, Nosrat C, Tomac A, Westphal H, et al., 1997. Neurturin and glial cell line-derived neurotrophic factor receptor-beta (GDNFR-beta), novel proteins related to GDNF and GDNFR-alpha with specific cellular patterns of expression suggesting roles in the developing and adult nervous system and in periphery. *J Neurosci* 17, 8506–8519.
- Widenfalk J, Tomac A, Lindqvist E, Hoffer B, et al., 1998. GFR $\alpha$ -3, a protein related to GFR $\alpha$ -1, is expressed in developing peripheral neurons and ensheathing cells. *Eur J Neurosci* 10, 1508–1517.
- Wieler M, Gee M, and Martin WRW, 2015. Longitudinal midbrain changes in early Parkinson's disease: Iron content estimated from R2\*/MRI. *Parkinsonism Relat Disord* 21, 179–183.
- Williams-Gray CH, Wijeyekoon R, Yarnall AJ, Lawson RA, et al., 2016. Serum immune markers and disease progression in an incident Parkinson's disease cohort (ICICLE-PD). *Mov Disord* 31, 995–1003.
- Williams LR, Inoué G, Cummins V, and Pelleymounter MA, 1996. Glial cell line-derived neurotrophic factor sustains axotomized basal forebrain cholinergic neurons in vivo: dose-response comparison to nerve growth factor and brain-derived neurotrophic factor. *J Pharmacol Exp Ther* 277, 1140–51.
- Williams SM, and Goldman-Rakic PS, 1998. Widespread origin of the primate mesofrontal dopamine system. *Cereb Cortex* 8, 321–345.
- Willkommen D, Lucio M, Moritz F, Forcisi S, et al., 2018. Metabolomic investigations in cerebrospinal fluid of Parkinson's disease. *PLoS One* 13, e0208752.
- Wiltshcko AB, Pettibone JR, and Berke JD, 2010. Opposite Effects of Stimulant and Antipsychotic Drugs on Striatal Fast-Spiking Interneurons. *Neuropsychopharmacology* 35, 1261–1270.
- Windle WF, and Cammermeyer J, 1958. Functional and Structural Observations on Chronically Reserpinized Monkeys. *Science (80- )* 127, 1503–1504.
- Winkler C, Sauer H, Lee CS, and Björklund A, 1996. Short-term GDNF treatment provides long-term rescue of lesioned nigral dopaminergic neurons in a rat model of Parkinson's disease. *J Neurosci* 16, 7206–15.
- Winner B, Jappelli R, Maji SK, Desplats PA, et al., 2011. In vivo demonstration that  $\alpha$ -synuclein oligomers are toxic. *Proc Natl Acad Sci* 108, 4194–4199.
- Wolter KG, Hsu YT, Smith CL, Nechushtan A, et al., 1997. Movement of Bax from the cytosol to mitochondria during apoptosis. *J Cell Biol* 139, 1281–1292.

- Wong LE, Gibson ME, Arnold HM, Pepinsky B, et al., 2015. Artemin promotes functional long-distance axonal regeneration to the brainstem after dorsal root crush. *Proc Natl Acad Sci U S A* 112, 6170–6175.
- Wonnacott S, Kaiser S, Mogg A, Soliakov L, et al., 2000. Presynaptic nicotinic receptors modulating dopamine release in the rat striatum. *Eur J Pharmacol* 393, 51–58.
- Worby CA, Vega QC, Chao HH, Seasholtz AF, et al., 1998. Identification and characterization of GFRalpha-3, a novel Co-receptor belonging to the glial cell line-derived neurotrophic receptor family. *J Biol Chem* 273, 3502–3508.
- Worby CA, Vega QC, Zhao Y, Chao HH, et al., 1996. Glial cell line-derived neurotrophic factor signals through the RET receptor and activates mitogen-activated protein kinase. *J Biol Chem* 271, 23619–23622.
- Worley DS, Pisano JM, Choi ED, Walus L, et al., 2000. Developmental regulation of GDNF response and receptor expression in the enteric nervous system. *Development* 127, 4383–4393.
- Wu DC, Jackson-Lewis V, Vila M, Tieu K, et al., 2002. Blockade of Microglial Activation Is Neuroprotective in the 1-Methyl-4-Phenyl-1,2,3,6-Tetrahydropyridine Mouse Model of Parkinson Disease. *J Neurosci* 22, 1763–1771.
- Xia Q, Liao L, Cheng D, Duong DM, et al., 2008. Proteomic identification of novel proteins associated with Lewy bodies. *Front Biosci* 13, 3850–3856.
- Xilouri M, Brekk OR, and Stefanis L, 2013. Alpha-synuclein and Protein Degradation Systems: a Reciprocal Relationship. *Mol Neurobiol* 47, 537–551.
- Xu K, and Dluzen DE, 2000. The effect of GDNF on nigrostriatal dopaminergic function in response to a two-pulse K(+) stimulation. *Exp Neurol* 166, 450–7.
- Yamamoto K, Yoshida H, Kokame K, Kaufman RJ, et al., 2004. Differential Contributions of ATF6 and XBP1 to the Activation of Endoplasmic Reticulum Stress-Responsive cis-Acting Elements ERSE, UPRE and ERSE-II. *J Biochem* 136, 343–350.
- Yan Q, Matheson C, and Lopez OT, 1995. In vivo neurotrophic effects of GDNF on neonatal and adult facial motor neurons. *Nature* 373, 341–344.
- Yan Y, Rato C, Rohland L, Preissler S, et al., 2019. MANF antagonizes nucleotide exchange by the endoplasmic reticulum chaperone BiP. *Nat Commun* 10, 541.
- Yan Z, and Surmeier DJ, 1996. Muscarinic (m2/m4) receptors reduce N- and P-type Ca2+ currents in rat neostriatal cholinergic interneurons through a fast, membrane- delimited, G-protein pathway. *J Neurosci* 16, 2592–2604.
- Yang F, Feng L, Zheng F, Johnson SW, et al., 2001. GDNF acutely modulates excitability and A-type K(+) channels in midbrain dopaminergic neurons. *Nat Neurosci* 4, 1071–8.
- Yang F, Pedersen NL, Ye W, Liu Z, et al., 2017. Moist smokeless tobacco (Snus) use and risk of Parkinson's disease. *Int J Epidemiol* 46, 872–880.
- Yang L, Chang CC, Sun Z, Madsen D, et al., 2017. GFRAL is the receptor for GDF15 and is required for the anti-obesity effects of the ligand. *Nat Med* 23, 1158–1166.
- Yang W, Shen Yujun, Chen Y, Chen L, et al., 2014. Mesencephalic astrocyte-derived neurotrophic factor prevents neuron loss via inhibiting ischemia-induced apoptosis. *J Neurol Sci* 344, 129–138.
- Ye H, Kuruvilla R, Zweifel LS, and Ginty DD, 2003. Evidence in support of signaling endosome-based retrograde survival of sympathetic neurons. *Neuron* 39, 57–68.
- Yilmaz R, Hopfner F, van Eimeren T, and Berg D, 2019. Biomarkers of Parkinson's disease: 20 years later. *J Neural Transm* 126, 803–813.
- Young HM, Hearn CJ, Farlie PG, Canty AJ, et al., 2001. GDNF Is a Chemoattractant for Enteric Neural Cells. *Dev Biol* 229, 503–516.
- Yu T, Scully S, Yu Y, Fox GM, et al., 1998a. Expression of GDNF family receptor components during development: implications in the mechanisms of interaction. *J Neurosci* 18, 4684–4696.
- Yue X, Hariri DJ, Caballero B, Zhang S, et al., 2014a. Comparative study of the neurotrophic effects elicited by VEGF-B and GDNF in preclinical in vivo models of Parkinson's disease. *Neuroscience* 258, 385–400.
- Yue Z, and Yang XW, 2013. Dangerous duet: LRRK2 and  $\alpha$ -synuclein jam at CMA. *Nat Neurosci* 16, 375–377.
- Yung KK, Bolam JP, Smith AD, Hersch SM, et al., 1995. Immunocytochemical localization of D1 and D2 dopamine receptors in the basal ganglia of the rat: light and electron microscopy. *Neuroscience* 65, 709–730.
- Yurek DM, and Fletcher-Turner A, 2002. Temporal changes in the neurotrophic environment of the denervated striatum as determined by the survival and outgrowth of grafted fetal dopamine neurons. *Brain Res* 931, 126–134.
- Yurek DM, and Fletcher-Turner A, 2001. Differential expression of GDNF, BDNF, and NT-3 in the aging nigrostriatal system following a neurotoxic lesion. *Brain Res* 891, 228–235.
- Zacchigna S, Lambrechts D, and Carmeliet P, 2008. Neurovascular signalling defects in neurodegeneration. *Nat Rev Neurosci* 9, 169–181.
- Zhang GL, Wang LH, Liu XY, Zhang YX, et al., 2018. Cerebral dopamine neurotrophic factor (CDNF) has neuroprotective effects against cerebral ischemia that may occur through the endoplasmic reticulum stress pathway. *Int J Mol Sci* 19.
- Zhang H, and Sulzer D, 2003. Glutamate Spillover in the Striatum Depresses Dopaminergic Transmission by Activating Group I Metabotropic Glutamate Receptors. *J Neurosci* 23, 10585–10592.
- Zhang K, and Kaufman RJ, 2008. From endoplasmic-reticulum stress to the inflammatory response. *Nature* 454, 455–462.
- Zhang W, Wang T, Pei Z, Miller DS, et al., 2005. Aggregated  $\alpha$ -synuclein activates microglia: a process leading to disease progression in Parkinson's disease. *FASEB J* 19, 533–542.
- Zhang Z, Shen Y, Luo H, Zhang F, et al., 2018. MANF protects dopamine neurons and locomotion defects from a human  $\alpha$ -synuclein induced Parkinson's disease model in *C. elegans* by regulating ER stress and autophagy pathways. *Exp Neurol* 308.
- Zhao H, Cheng L, Du X, Hou Y, et al., 2016. Transplantation of Cerebral Dopamine Neurotrophic Factor Transduced BMSCs in Contusion Spinal Cord Injury of Rats: Promotion of Nerve Regeneration by Alleviating Neuroinflammation. *Mol Neurobiol* 53, 187–199.
- Zhao H, Cheng L, Liu Y, Zhang W, et al., 2014a. Mechanisms of Anti-inflammatory Property of Conserved Dopamine Neurotrophic Factor: Inhibition of JNK Signaling in Lipopolysaccharide-Induced Microglia. *J Mol Neurosci* 52, 186–192.
- Zhao H, Liu Y, Cheng L, Liu B, et al., 2013. Mesencephalic Astrocyte-Derived Neurotrophic Factor Inhibits Oxygen–Glucose Deprivation-

- Induced Cell Damage and Inflammation by Suppressing Endoplasmic Reticulum Stress in Rat Primary Astrocytes. *J Mol Neurosci* 51, 671–678.
- Zhou C, Xiao C, Commissiong JW, Krnjevic K, et al., 2006. Mesencephalic astrocyte-derived neurotrophic factor enhances nigral gamma-aminobutyric acid release. *Neuroreport* 17, 293–297.
- Zhou F-M, Liang Y, and Dani JA, 2001. Endogenous nicotinic cholinergic activity regulates dopamine release in the striatum. *Nat Neurosci* 4, 1224–1229.
- Zhou F-W, Jin Y, Matta SG, Xu M, et al., 2009. An Ultra-Short Dopamine Pathway Regulates Basal Ganglia Output. *J Neurosci* 29, 10424–10435.
- Zhou W, Chang L, Fang Y, Du Z, et al., 2016. Cerebral dopamine neurotrophic factor alleviates A $\beta$ 25-35-induced endoplasmic reticulum stress and early synaptotoxicity in rat hippocampal cells. *Neurosci Lett*.
- Zhu S, Zhao C, Wu Y, Yang Q, et al., 2015. Identification of a Vav2-dependent mechanism for GDNF/Ret control of mesolimbic DAT trafficking. *Nat Neurosci* 18, 1084–1093.
- Zhu W, Li J, Liu Y, Xie K, et al., 2016. Mesencephalic astrocyte-derived neurotrophic factor attenuates inflammatory responses in lipopolysaccharide-induced neural stem cells by regulating NF- $\kappa$ B and phosphorylation of p38-MAPKs pathways. *Immunopharmacol Immunotoxicol* 38, 205–213.
- Zhu W, Xie W, Pan T, Jankovic J, et al., 2008. Comparison of neuroprotective and neurorestorative capabilities of rasagiline and selegiline against lactacystin-induced nigrostriatal dopaminergic degeneration. *J Neurochem* 105, 1970–1978.
- Zigmond MJ, and Smeyne RJ, 2014. Exercise: Is it a neuroprotective and if so, how does it work? *Parkinsonism Relat Disord* 20, S123–S127.
- Zimprich A, Biskup S, Leitner P, Lichtner P, et al., 2004. Mutations in LRRK2 Cause Autosomal-Dominant Parkinsonism with Pleomorphic Pathology. *Neuron* 44, 601–607.
- Zou L-L, Xu J, Jankovic J, He Y, et al., 2000. Pramipexole inhibits lipid peroxidation and reduces injury in the substantia nigra induced by the dopaminergic neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine in C57BL/6 mice. *Neurosci Lett* 281, 167–170.
- Åkerud P, Alberch J, Eketjäll S, Wagner J, et al., 1999. Differential effects of glial cell line-derived neurotrophic factor and neurturin on developing and adult substantia nigra dopaminergic neurons. *J Neurochem* 73, 70–78.
- Åkerud P, Holm PC, Castelo-Branco G, Sousa K, et al., 2002. Persephin-Overexpressing Neural Stem Cells Regulate the Function of Nigral Dopaminergic Neurons and Prevent Their Degeneration in a Model of Parkinson's Disease. *Mol Cell Neurosci* 21, 205–222.
- Websites:
- The Human Protein Atlas, 2020, [www.proteinatlas.org](http://www.proteinatlas.org), visited 4.2.2020
- Herantis Pharma Plc, Company press release 27 August 2020, [www.herantis.com](http://www.herantis.com), visited 12.10.2020